



<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each rectal suppository)</b> Acetaminophen, USP 120 mg. Pain reliever/fever reducer</p> <p><b>Purposes</b></p> <p><b>Uses</b> temporarily • reduces fever • relieves minor aches, pains, and headache</p> <p><b>Warnings</b></p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen.</p> <p><b>Allergy alert:</b> acetaminophen may cause severe skin reactions. Symptoms may include: skin redness • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>For rectal use only.</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• if you are allergic to acetaminophen</li> <li>• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>• the child has liver disease.</li> <li>• the child is taking the blood thinning drug warfarin.</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>• fever lasts more than 3 days (72 hours), or recurs.</li> <li>• pain lasts more than 5 days or gets worse.</li> <li>• redness or swelling is present in the painful area.</li> </ul> <p>These may be signs of a serious condition.</p>													
<p><b>Keep out of reach of children.</b> If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>• this product does not contain directions or warnings for adult use</li> <li>• do not use more than directed</li> <li>• remove wrapper</li> <li>• carefully insert suppository well up into the rectum</li> </ul> <p><b>Dosing Chart</b></p> <table border="1"> <thead> <tr> <th>Age</th> <th>Dose</th> </tr> </thead> <tbody> <tr> <td>under 3 years</td> <td>Do not use unless directed by a doctor</td> </tr> <tr> <td>3 to 6 years</td> <td>Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)</td> </tr> </tbody> </table> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• store at 2°-27°C (35°-80°F)</li> <li>• do not use if imprinted suppository wrapper is opened or damaged</li> </ul> <p><b>Inactive ingredients</b></p> <p>glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80</p> <p><b>Questions?</b> call 1-866-923-4914</p>	Age	Dose	under 3 years	Do not use unless directed by a doctor	3 to 6 years	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)	<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b> If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>• this product does not contain directions or warnings for adult use</li> <li>• do not use more than directed</li> <li>• remove wrapper</li> <li>• carefully insert suppository well up into the rectum</li> </ul> <p><b>Dosing Chart</b></p> <table border="1"> <thead> <tr> <th>Age</th> <th>Dose</th> </tr> </thead> <tbody> <tr> <td>under 3 years</td> <td>Do not use unless directed by a doctor</td> </tr> <tr> <td>3 to 6 years</td> <td>Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)</td> </tr> </tbody> </table> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• store at 2°-27°C (35°-80°F)</li> <li>• do not use if imprinted suppository wrapper is opened or damaged</li> </ul> <p><b>Inactive ingredients</b></p> <p>glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80</p> <p><b>Questions?</b> call 1-866-923-4914</p>	Age	Dose	under 3 years	Do not use unless directed by a doctor	3 to 6 years	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)
Age	Dose												
under 3 years	Do not use unless directed by a doctor												
3 to 6 years	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)												
Age	Dose												
under 3 years	Do not use unless directed by a doctor												
3 to 6 years	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)												

Laser Ablation Lot # Ex .

(2plcs) Not actual code Code bars to be changed to match t is product



NDC 51672-2114-2



### Doctor Recommended Pain Reliever/Fever Reducer

- ✓ No Parabens or Any Other Preservatives
- ✓ No Artificial Colors

\*actual size



INFANTS' ages 6-36 months

# FeverAll

ACETAMINOPHEN SUPPOSITORIES

Pain Reliever/Fever Reducer

**6 Rectal Suppositories** | **80 mg each**



Dist. by: Taro Pharmaceuticals U.S.A., Inc.  
Hawthorne, NY 10532 Made in USA  
GW 7303 Rev. 07/2019

Tamper-evident: Suppositories are individually wrapped. Do not use if imprinted wrapper is opened or damaged.



**Drug Facts**

**Active ingredient (in each rectal suppository)** Acetaminophen, USP 80 mg. Pain reliever/fever reducer

**Uses** temporarily • reduces fever • relieves minor aches, pains, and headache

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes  
• more than 4 doses in 24 hours, which is the maximum daily amount for ages 6 to 11 months  
• more than 5 doses in 24 hours, which is the maximum daily amount for ages 12 to 36 months  
• with other drugs containing acetaminophen  
**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
• skin redness • rash • blisters • hives  
If a skin reaction occurs, stop use and seek medical help right away.  
**Do not use**  
• if you are allergic to acetaminophen.  
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
**Ask a doctor before use if**  
• the child has liver disease.  
• the child is taking the blood thinning drug warfarin.  
**Stop use and ask a doctor if**  
• fever lasts more than 3 days (72 hours), or recurs.  
• pain lasts more than 3 days or gets worse.  
• new symptoms occur.  
• redness or swelling is present in the painful area.  
These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

**Directions • this product does not contain directions or warnings for adult use**  
• do not use more than directed  
• remove wrapper  
• carefully insert suppository well up into the rectum

**Dosing Chart**

Age	Dose
6 months under	Do not use unless directed by a doctor
6 to 11 months	Use 1 suppository every 6 hours
12 to 36 months	Use 1 suppository every 4 to 6 hours (maximum of 4 doses in 24 hours)
12 to 36 months	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)

**Other information**  
• store at 20°-27°C (68°-80°F)  
• do not use if imprinted suppository wrapper is opened or damaged

**Inactive ingredients**  
glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polyсорate 80

**Questions?** call 1-866-923-4914

Laser Ablation  
Lot #  
Ex .

(2plcs) Not actual code  
Code bars to be changed  
to match t is product

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

VALERIE S PRATT  
02/07/2020 04:34:55 PM  
Signing on behalf of Dr. Karen Mahoney