

PRODUCT SHOULD BE DISCARDED
2 MONTHS AFTER INITIAL USE.

Barcode
3 68791-103-20 2

NDC 68791-103-20

Rx only

DermOtic® Oil
(fluocinolone acetonide oil) 0.01%
Ear Drops

For Otic Use Only
Not for Ophthalmic Use

Net contents: 20 mL



GTIN: 00368791103202
EXP/LOT/SN



Non Coating Area



**For Otic Use Only
Not for Ophthalmic Use**
Dosage and Administration:
For the treatment of chronic exogenous external otitis, itching, edema, and redness of the ear. Using the supplied ear-dropper, apply 5 drops of DermOtic® Oil (fluocinolone acetonide oil) 0.01% to the ear canal of the affected ear 2 times daily. Do not use if you are allergic to any of the ingredients. Gently pull the earlobe backward and upward and apply 5 drops of DermOtic® Oil into the ear canal. Gently pat excess material dripping out of the ear using a clean cotton ball. Follow these instructions twice each day for 7 to 14 days.
See package insert for full prescribing information

CODE NO. 171A237-3

NDC 68791-103-20

Rx only

DermOtic® Oil
(fluocinolone acetonide oil) 0.01%
Ear Drops

For Otic Use Only
Not for Ophthalmic Use

Net contents: 20 mL



CONTAINS: Active: Fluocinolone Acetonide USP, (0.01%)
Inactive: Isopropyl Alcohol USP, (1.0%), in a base containing, Refined Peanut Oil NF,
Light Mineral Oil NF, Isopropyl Myristate NF, Oleth-2, Cream Fragrance, Balsam Pine
CONTENTS OF PACKAGE: 20mL in 1 ea bottle, 1 Dropper
STORAGE: Keep tightly closed. Store at 25°C (77°F); excursions permitted to 15°-30°C
(59°-86°F).

Date of first opening / / Manufactured by: Hill Demaceuticals, Inc.*
Discard unused portion 2 months after first opening. Skidport, FL 32773
for: Royal Pharmaceutical® Manassas, VA 20108
KEEP OUT OF REACH OF CHILDREN. 1-800-510-3403

NDC 68791-103-20
Rx only
DermOtic® Oil
(flucinolone acetonide oil) 0.01%
Ear Drops
For Otic Use Only
Not for Ophthalmic Use

Net contents: 20 mL



CONTAINS:
Active: Fluocinolone Acetonide
USP, (0.01%)
Contains Refined Peanut Oil

STORAGE:
Keep tightly closed. Store at 25°C
(77°F); excursions permitted to
15°- 30°C (59°- 86°F).

Date of first opening / /
Discard unused portion 2 months
after first opening.

**KEEP OUT OF REACH OF CHILDREN.
PRODUCT SHOULD BE DISCARDED
2 MONTHS AFTER INITIAL USE.**

Manufactured by:
Hill Dermaceuticals, Inc.®
Sanford, FL 32773
for: Royal Pharmaceuticals®
Manasquan, NJ 08736
1-800-510-3401
CODE NO. 171A238-2

**For Otic Use Only
Not for Ophthalmic Use**
Dosage and Administration:
For the treatment of chronic
eczematous external otitis: itching,
dryness, and redness of the ear. Using
the supplied ear-dropper, apply 5 drops
of DermOtic® Oil into the affected ear.
To apply, tilt head to one side so that the
ear is facing up. Then gently pull the ear
lobe backward and upward and apply
5 drops of DermOtic® Oil into the ear.
Keep head tilted for about a minute to
allow DermOtic® Oil to penetrate lower
into the ear canal. Gently pat excess
material dripping out of the ear using a
clean cotton ball. Follow these
instructions twice each day for 7 to 14 days.

See package insert for full prescribing
information.

LOT No.:
EXP. DATE:



3 DRAFT LABELING TEXT

The Packaging Insert is being revised per the FDA Information Request dated November 21, 2019 to add the statement, “PRODUCT SHOULD BE DISCARDED 2 MONTHS AFTER INITIAL USE.”, based on In-Use stability data (see 3.2.P.8.1 and 3.2.P.8.3). Two administrative changes are also being made. The first is an update to the revision number of Hill Dermaceuticals’ internal insert code number and the second is the omission of the revision date previously listed next to the insert code number as the internal insert code number is adequate to identify the insert revision. The Brand name package insert has been revised as indicated in 1.14.1.2.1.

3.1 Draft Package Insert

BRAND NAME PACKAGE INSERT

Front Page of Insert


DermOtic[®] Oil
(fluocinolone acetonide oil) 0.01%
Ear Drops

For Otic Use Only-
Not for Ophthalmic Use

NDC 68791-103-20

DESCRIPTION

DermOtic[®] Oil Ear Drops contain fluocinolone acetonide {(6 α ,11 β ,16 α)-6,9-difluoro-11,21-dihydroxy-16,17[(1-methylethylidene)bis(oxy)]-pregna-1,4-diene-3,20-dione, cyclic 16,17 acetal with acetone}, a synthetic corticosteroid. This formulation is also marketed as Derma-Smoother/FS[®] Body Oil for the treatment of atopic dermatitis and Derma-Smoother/FS[®] Scalp Oil for the treatment of psoriasis of the scalp. Chemically, fluocinolone acetonide is C₂₄H₃₀F₂O₆. It has the following structural formula:



Fluocinolone acetonide in DermOtic[®] Oil has a molecular weight of 452.50. It is a white crystalline powder that is odorless, stable in light, and melts at 270°C with decomposition; soluble in alcohol, acetone and methanol; slightly soluble in chloroform; insoluble in water.

Each gram of DermOtic[®] Oil contains approximately 0.11 mg of fluocinolone acetonide in a blend of oils, which contains isopropyl alcohol, isopropyl myristate, light mineral oil, oleth-2, refined peanut oil NF and fragrances.

CLINICAL PHARMACOLOGY

Like other topical corticosteroids, fluocinolone acetonide has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear. However, corticosteroids are thought to act by the induction of phospholipase A2 inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A2.

Pharmacokinetics: The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle and the integrity of the epidermal barrier. Occlusion of topical corticosteroids can enhance penetration. Topical corticosteroids can be absorbed from normal intact skin. Also, inflammation and/or other disease processes in the skin can increase percutaneous absorption.

DermOtic[®] Oil is in the low to medium range of potency as compared with other topical corticosteroids.

CLINICAL STUDIES

Efficacy in a placebo-controlled study for the treatment of chronic eczematous external otitis on 154 patients (adults and children 2 years of age and older) treated with five drops per ear of DermOtic[®] Oil twice daily, after 7 days of treatment, showed DermOtic[®] Oil to be superior to placebo in clearing the signs and symptoms of eczematous external otitis.

Clinical safety studies were conducted on the same formulation of fluocinolone acetonide oil 0.01%, marketed as Derma-Smoothe/FS[®] Topical Oil. Open-label safety studies on 33 children (20 subjects ages 2 to 6 years, 13 subjects ages 7 to 12 years) with moderate to severe stable atopic dermatitis, and baseline body surface area involvement greater than 75% in 18 patients, and 50% to 75% in 15 patients, were treated with Derma-Smoothe/FS[®] Topical Oil twice daily for 4 weeks. Morning pre-stimulation cortisol level and post-Cortrosyn stimulation cortisol level were obtained in each subject at the beginning of the trial and at the end of 4 weeks of treatment. At the end of treatment, 4 out of 18 subjects aged 2 to 5 years showed low pre-stimulation cortisol levels (3.2 to 6.6 µg/dL; normal: cortisol > 7 µg/dL) but all had normal responses to 0.25 mg of Cortrosyn stimulation (cortisol > 18 µg/dL).

A clinical study was conducted to assess the safety of Derma-Smoothe/FS[®] Topical Oil, which contains refined peanut oil, on subjects with known peanut allergies. The study enrolled 13 patients with atopic dermatitis, 6 to 17 years of age. Of the 13 patients, 9 were Radioallergosorbent Test (RAST) positive to peanuts and 4 had no peanut sensitivity (controls). The study evaluated the responses to both prick test and patch test utilizing peanut oil NF, Derma-Smoothe/FS[®] Topical Oil and histamine/ saline controls on the 13 individuals.

These subjects were also treated with Derma-Smoothe/FS[®] Topical Oil twice daily for 7 days. Prick test and patch test results for all 13 patients were negative to Derma-Smoothe/FS[®] Topical Oil and the refined peanut oil. One of the 9 peanut-sensitive patients experienced an exacerbation of atopic dermatitis after 5 days of Derma-Smoothe/FS[®] Topical Oil use. Importantly, the bulk peanut oil NF, used in Derma-Smoothe/FS[®] Topical Oil is heated at 475°F for at least 15 minutes, which should provide for adequate decomposition of allergenic proteins.

INDICATION AND USAGE

DermOtic[®] Oil is a low to medium potency corticosteroid indicated for the treatment of chronic eczematous external otitis in adults and pediatric patients 2 years and older.

CONTRAINDICATIONS

DermOtic[®] Oil is contraindicated in those patients with a history of hypersensitivity to any of the components of the preparation.

This product contains refined peanut oil NF (See PRECAUTIONS).

PRECAUTIONS

General: Systemic absorption of topical corticosteroids can produce reversible hypothalamic-pituitary-adrenal (HPA) axis suppression with the potential for glucocorticosteroid insufficiency after withdrawal of treatment. Manifestations of Cushing's syndrome, hyperglycemia, and glucosuria can also be produced in some patients by systemic absorption of topical corticosteroids while on treatment.

Patients applying a topical steroid to a large surface area or to areas under occlusion should be evaluated periodically for evidence of HPA axis suppression. This may be done by using the ACTH stimulation, A.M. plasma cortisol, and urinary free cortisol tests.

If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent corticosteroid. Infrequently, signs and symptoms of glucocorticoid insufficiency may occur requiring supplemental systemic corticosteroids. For information on systemic supplementation, see prescribing information for those products.

Children may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios. (See PRECAUTIONS-Pediatric use).

Allergic contact dermatitis to any component of topical corticosteroids is usually diagnosed by a failure to heal rather than noting a clinical exacerbation, which may occur with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic testing. One peanut-sensitive child experienced a flare of his atopic dermatitis after 5 days of twice daily treatment with Derma-Smoothe/FS[®] Topical Oil (see CLINICAL STUDIES section).

If wheal and flare type reactions (which may be limited to pruritus) or other manifestations of hypersensitivity develop, DermOtic[®] Oil should be discontinued immediately and appropriate therapy instituted.

If concomitant skin infections are present or develop, an appropriate antifungal or antibacterial agent should be used. If a favorable response does not occur promptly, use of DermOtic[®] Oil should be discontinued until the infection has been adequately controlled.

Back Page of Insert:

DermOtic® Oil is formulated with 48% refined peanut oil NF. Peanut oil used in this product is routinely tested for peanut proteins through amino acid analysis; the quantity of amino acids is below 0.5 parts per million (ppm). Physicians should use caution in prescribing DermOtic® Oil for peanut-sensitive individuals.

Information for Patients: Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external ear use only. Do not use occlusive dressings.
2. Avoid contact with the eyes. In case of contact, wash eyes liberally with water.
3. This medication should not be used for any disorder other than that for which it was prescribed.
4. Patients should promptly report to their physician any worsening of their skin condition.
5. As with other corticosteroids, therapy should be discontinued when control is achieved. If no improvement is seen within 2 weeks, contact the physician.

Laboratory Tests: The following tests may be helpful in evaluating patients for HPA axis suppression:

ACTH stimulation test

A.M. plasma cortisol test

Urinary free cortisol test

Carcinogenesis, mutagenesis, and impairment of fertility: Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of DermOtic® Oil. Studies have not been performed to evaluate the mutagenic potential of fluocinolone acetonide, the active ingredient in DermOtic® Oil. Some corticosteroids have been found to be genotoxic in various genotoxicity tests (i.e. the in vitro human peripheral blood lymphocyte chromosome aberration assay with metabolic activation, the in vivo mouse bone marrow micronucleus assay, the Chinese hamster micronucleus test and the in vitro mouse lymphoma gene mutation assay).

Pregnancy: Teratogenic effects: Pregnancy category C: Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Some corticosteroids have been shown to be teratogenic after dermal application in laboratory animals.

There are no adequate and well-controlled studies in pregnant women on teratogenic effects from DermOtic® Oil. Therefore, DermOtic® Oil should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers: Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when DermOtic® Oil is administered to a nursing woman.

Pediatric Use: DermOtic® Oil may be used twice daily for up to 2 weeks in pediatric patients 2 years of age and older with chronic eczematous external otitis.

DermOtic® Oil is not recommended for use on the face (See ADVERSE REACTIONS section).

Because of a higher ratio of skin surface area to body mass, children are at a greater risk than adults of HPA-axis-suppression when they are treated with topical corticosteroids. They are therefore also at greater risk of glucocorticosteroid insufficiency after withdrawal of treatment and of Cushing's syndrome while on treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in infants and children. (See PRECAUTIONS).

HPA axis suppression, Cushing's syndrome, and intracranial hypertension have been reported in children receiving topical corticosteroids. Children may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios. Manifestations of adrenal suppression in children include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

DermOtic® Oil is formulated with 48% refined peanut oil NF. Peanut oil used in this product is routinely tested for peanut proteins through amino acid analysis; the quantity of amino acids is below 0.5 parts per million (ppm). Physicians should use caution in prescribing DermOtic® Oil for peanut-sensitive individuals.

ADVERSE REACTIONS

The following local adverse reactions have been reported infrequently with topical corticosteroids. They may occur more frequently with the use of occlusive dressings, especially with higher potency corticosteroids. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, skin atrophy, striae, and miliaria. One peanut sensitive child experienced a flare of his atopic dermatitis after 5 days of twice daily treatment with Derma- Smothe/FS® Topical Oil.

OVERDOSAGE

Topically applied DermOtic® Oil can be absorbed in sufficient amounts to produce systemic effects (See PRECAUTIONS).

DOSAGE AND ADMINISTRATION

For the treatment of chronic eczematous external otitis, using the supplied ear-dropper, apply 5 drops of DermOtic® Oil into the affected ear. To apply, tilt head to one side so that the ear is facing up. Then gently pull the ear lobe backward and upward and apply 5 drops of DermOtic® Oil into the ear. Keep head tilted for about a minute to allow DermOtic® Oil to penetrate lower into the ear canal. Gently pat excess material dripping out of the ear using a clean cotton ball. Follow these instructions twice each day for 7 to 14 days.

HOW SUPPLIED

DermOtic[®] Oil (fluocinolone acetonide oil) 0.01% Ear Drops is supplied in 1 fluid ounce bottles containing 20 mL, net content (Dropper Included) (NDC # 68791-103-20).

Keep tightly closed. Store at 20°-25° C (68° to 77° F); excursions permitted to 15°-30°C (59°-86°F.) [see USP Controlled Room Temperature].

Discard DermOtic Oil 2 months after initial use.

CAUTION: Rx only