




FRONT



BACK

Actual Size 100%

FDA00305-2

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	8.1 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	12 pt
Leading	≥0.5 pt	0-0.5 pt
Hair lines	0.5 pt rule	0.5 pt
Box and Borderline†	>0.5 pt rule	2.5 pt
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in ODP
 † FDA recommends 2.5pt rule

ALL DAY STRONG®
ALEVE®
 naproxen sodium tablets, 220 mg (NSAID)
 Pain reliever/fever reducer

STRENGTH TO LAST 12 HOURS 10 TABLETS

Read Vial Blister Card Before Use.
Uses temporarily relieves minor aches and pains and temporarily reduces fever. *Warnings Allergy alert:* Naproxen sodium may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing) or shock. *Stomach bleeding warning:* This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. Ask a doctor or pharmacist before use if you are taking any other drug. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), or if pain or fever gets worse. If pregnant or breast-feeding, ask a health professional before use. *Directions* Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 tablet every 8 to 12 hours while symptoms last. For the first dose you may take 2 tablets within the first hour. Do not exceed 2 tablets in any 8- to 12-hour period. Do not exceed 3 tablets in a 24-hour period. Children under 12 years: ask a doctor. Store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F). *Questions or comments?* 1-800-395-0689 (Mon-Fri 9AM - 5PM EST)
 Dist. by:
 Bayer HealthCare LLC
 Whippany, NJ 07981 00000

Lot/Exp Date Area

200%

ALL DAY STRONG®
ALEVE®
 naproxen sodium tablets, 220 mg (NSAID)
 Pain reliever/fever reducer

STRENGTH TO LAST 12 HOURS 10 TABLETS

Read Vial Blister Card Before Use.
Uses temporarily relieves minor aches and pains and temporarily reduces fever. *Warnings Allergy alert:* Naproxen sodium may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing) or shock. *Stomach bleeding warning:* This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. Ask a doctor or pharmacist before use if you are taking any other drug. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), or if pain or fever gets worse. If pregnant or breast-feeding, ask a health professional before use. *Directions* Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 tablet every 8 to 12 hours while symptoms last. For the first dose you may take 2 tablets within the first hour. Do not exceed 2 tablets in any 8- to 12-hour period. Do not exceed 3 tablets in a 24-hour period. Children under 12 years: ask a doctor. Store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F). *Questions or comments?* 1-800-395-0689 (Mon-Fri 9AM - 5PM EST)
 Dist. by:
 Bayer HealthCare LLC
 Whippany, NJ 07981 00000

Lot/Exp Date Area

Actual Size 100%

FDA00295-1

ICL (updated 2/27/19)		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold italic*	4.5 pt
Subheadings	6 pt*	4.5 pt
Body Text	6 pt*	4.5 pt
Bullets	5 pt	6 pt
Hairlines	0.5 pt rule	-
Barlines†	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
 *ICL minimum font size is 4.5 pt.
 † 0.5 pt leading may be used as long as ascenders and descenders do not touch.
 † FDA recommends size 2.5 pt type




Actual Size 100%

FDA00308-1

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0 - 0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 30	≤30

* Headings are to be at least 2pts. larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in ODP
 † FDA Recommends 2.5pt. rule






200%



Actual Size 100%

FDA00403-2



ICL (updated 2/27/19)		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	NA
Drug Facts Continued	7 pt*	NA
Headings	7 pt- bold italic *	4.5 pt
Subheadings	8 pt*	4.5 pt
Body Text	8 pt*	4.5 pt
Bullets	5 pt	8 pt
Hairlines	0.5 pt rule	NA
Barlines†	> 0.5 pt rule	NA
Characters per inch rule is not applicable to ICL. *ICL minimum font size is 4.5 pt. † 0.5 pt leading may be used as long as ascenders and descenders do not touch. † FDA recommends size 2,5 pt type		




Actual Size 100%

FDA00301-2

DRUG FACTS INFO updated 5/19/18		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0 pt. - 0.5 pt.
Hair lines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type sizes used in ODP.
 † FDA recommends 2.5pt rule.






200%



Actual Size 100%

FDA00404-3



ICL (updated 2/27/19)		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	NA
Drug Facts Continued	7 pt*	NA
Headings	7 pt- bold italic *	4.5 pt
Subheadings	8 pt*	4.5 pt
Body Text	8 pt*	4.5 pt
Bullets	5 pt	8 pt
Hairlines	0.5 pt rule	NA
Barlines†	> 0.5 pt rule	NA
Characters per inch rule is not applicable to ICL. *ICL minimum font size is 4.5 pt. † 0.5 pt leading may be used as long as ascenders and descenders do not touch. † FDA recommends size 2,5 pt type		



Actual Size 100%

FDA00303-4

DRUG FACTS INFO Updated 5/18/19		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	6.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 30	≤ 30

* Headings are to be at least 2 pts. larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommendations: 2.5 pt. rule



200%



Actual Size 100%

FDA00293-1

Don't **Car** **for**
Full **Information** **Warnings**
Allergic **Alert:** Naproxen sodium may cause a
severe **allergic** **reaction** **which** **may** **include:** **hives,** **itching,** **swelling,** **asthma** **(wheezing)** **or** **shock.** **Stomach** **bleeding**
warning: This product contains a nonsteroidal anti-inflammatory
drug **(NSAID),** **which** **may** **cause** **severe** **stomach** **bleeding.** **The** **chance**
is **higher** **if** **you:** • **are** **age** **60** **or** **older** • **have** **had** **stomach** **ulcers** **or**
bleeding **problems** • **take** **a** **blood** **thinning** **(anticoagulant)** **or** **steroid** **drug**
• **take** **other** **drugs** **containing** **prescription** **or** **nonprescription** **NSAIDs**
(aspirin, **ibuprofen,** **naproxen,** **or** **others)** • **have** **3** **or** **more** **alcoholic** **drinks**
every **day** **while** **using** **this** **product** • **take** **more** **or** **for** **a** **longer** **time** **than** **directed.**
Heart **attack** **and** **stroke** **warning:** NSAIDs, **except** **aspirin,** **increase** **the** **risk** **of** **heart**
attack, **heart** **failure,** **and** **stroke.** **These** **can** **be** **fatal.** **The** **risk** **is** **higher** **if** **you** **use** **more**
than **directed** **or** **for** **longer** **than** **directed.** **Do** **not** **use** **if** **you** **have** **ever** **had** **an** **allergic**
reaction **to** **any** **other** **pain** **reliever/fever** **reducer.** **Stop** **use** **and** **ask** **a** **doctor** **if** **you**
experience **any** **of** **the** **following** **signs** **of** **stomach** **bleeding** **(red** **flaxy,** **vomit** **blood,** **have**
stool **that** **is** **black** **or** **clay,** **have** **stomach** **pain** **that** **does** **not** **get** **better),** **have** **symptoms** **of**
heart **problems** **or** **stroke** **(chest** **pain,** **trouble** **breathing,** **weakness** **in** **one** **part** **or** **side** **of**
body, **altered** **speech,** **leg** **swelling),** **or** **if** **pain** **or** **fever** **gets** **worse.** **If** **pregnant** **or**
breast-feeding, **ask** **a** **health** **professional** **before** **use.** **Directions** **ADULTS** **and** **children**
12 **years** **and** **older:** **take** **1** **tablet** **with** **water** **every** **8** **to** **12** **hours** **while** **symptoms** **last.**
For **the** **first** **dose** **you** **may** **take** **2** **tablets** **within** **the** **first** **hour.** **Do** **not** **exceed**
2 **tablets** **in** **a** **24-hour** **period.** **Children** **under** **12** **years:** **ask** **a** **doctor,** **dentist**
or **pharmacist.** **1-800-368-0688**
(Mon-Fri 9AM-5PM EST)
Dist. by: Bayer HealthCare LLC
Whispery, NJ 07081

8598791

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	-
Drug Facts Continued	8 pt.	-
Headings*	8 pt- bold <i>italic</i> †	4,5 pt.
Subheadings	>6 pt.	4,5 pt.
Body Text	>6 pt.	4,5 pt.
Bullets	6 pt.	6 pt.
Leading	>0,5 pt.	4,5 pt.
Headlines	0,5 pt. rule	-
Box and Barlines	>0,5 pt. rule	-
Characters Per Inch:	No more than 39	> 39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2,5pt rule




200%



Actual Size 100%

FDA00274-5



Actual Size 100%

FDA00169-1

DRUG FACTS INFO (updated 5/18/16)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings [†]	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0.5 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barlines [†]	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 38	≤39

[†] Headings are to be at least 2 pts larger than Body Text.
^{**} Drug Facts Title to be larger than all other type sizes used in DFP.
[†] FDA Recommends 2.5pt rule.

Do not use this product if printed safety seal bearing "Bayer HealthCare" under cap is torn or missing.

OPEN TO READ DRUG FACTS BEFORE USE

LEFT SIDE

Drug Facts (continued)

Warnings or Precautions: ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless, definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years ● ask a doctor

Other Information

- each tablet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive Ingredients: FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-368-0898 (Plan - FIDAM - RPM EST)

Drug Facts

Active Ingredient (in each tablet)

Naproxen sodium (naproxen 200 mg) (NSAID).....Pain reliever/fever reducer

"nonsteroidal anti-inflammatory drug"

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
- the common cold
- temporarily reduces fever

Warnings

Allergic Reaction: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin redness
- blisters
- facial swelling
- stomach
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Use with caution in patients taking aspirin, NSAIDs, or aspirin. Increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before you use it

- if you are taking any other pain reliever/fever reducer
- if you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- if you are taking a diuretic
- if you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease the benefit of aspirin
- taking any other drug

What to watch for:

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - stomach pain
 - black stools
 - blood in stool
 - blood in vomit
 - black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear



200%



Actual Size 100%

FDA00300-1

LOT:
EXP:
MADE IN

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Bayer, the Bayer Cross, Aleve,
All Day Strong, and Soft Grip
are registered trademarks of Bayer.

Bayer

Actual Size 100%

FDA00170-1

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	>6 pt.	6 pt.
Body Text	>6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0.5 pt.
Headlines	0.5 pt.rule	0.5 pt.
Box and Bar lines†	>0.5 pt.rule	2.5 pt.
Characters Per Inch	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type sizes used in DFP.
† FDA Recommendation: 2.5pt rule.

Do not use this product if printed safety seal bearing "Bayer HealthCare" under cap is torn or missing.

OPEN TO READ DRUG FACTS BEFORE USE

Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 tablet every 8 to 12 hours while symptoms last
- do not take more than 2 tablets every 24 hours
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years • ask a doctor

Other Information

- each tablet contains sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive Ingredients FD&C Blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polydione, talc, titanium dioxide

Questions or comments? 1-800-868-8686 (Mon-Fri 9AM-6PM EST)

Drug Facts

Active Ingredient (in each tablet)

Naproxen sodium 220 mg (naproxen 200 mg)

Purposes

.....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

CLIP IT MEDIA

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - backache
 - headache
 - the common cold
- temporarily reduces fever

Warnings

Always alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- dizziness
- difficulty breathing

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause serious stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, including naproxen, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease the benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- you have a rash or hives
- any new symptoms appear

Drug Facts

Active Ingredient (in each tablet)

Naproxen sodium 220 mg (naproxen 200 mg)

Purposes

.....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug



Actual Size 100%

FDA00289-3

DRUG FACTS INFO (updated 5/19/19)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt-bold italic*	8 pt.
Subheadings	≥6 pt.	8 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	>0.5 pt.	6.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch	No more than 36	≤ 36

* Headings can be in at least 2x larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP.
 † FDA Recommends 2.5pt rule.




Actual Size 100%

FDA00306-1

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0 - 0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 30	≤30

* Headings are to be at least 2pts. larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP.
 † FDA Recommends 2.5pt. rule





200%



Actual Size 100%

FDA00408-2



ICL (updated 2/27/19)		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold (ital)c*	-
Subheadings	6 pt*	4,5 pt
Body Text	6 pt*	4,5 pt
Bullets	5 pt	8 pt
Headlines	0,5 pt rule	-
Barlines [†]	> 0,5 pt rule	-
Characters per inch rule is not applicable to ICL. *CL minimum font size is 4,5 pt. < 0,5 pt leading may be used as long as ascenders and descenders do not touch. † FDA recommends size 2,5 pt type		




Actual Size 100%

FDA00281-4

DRUG FACTS INFO <small>updated 5/19/18</small>		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0 pt. - 0.5 pt.
Hair lines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2x's larger than Body Text.
** Drug Facts Title to be larger than all other type sizes used in ODP.
† FDA recommends 2.5pt. rule





200%



Actual Size 100%

FDA00407-2



ICL (updated 2/27/19)		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold (ital)c*	-
Subheadings	6 pt*	4,5 pt
Body Text	6 pt*	4,5 pt
Bullets	5 pt	8 pt
Headlines	0,5 pt rule	-
Barlines [†]	> 0,5 pt rule	-
Characters per inch rule is not applicable to ICL. *ICL minimum font size is 4,5 pt. † < 0,5 pt leading may be used as long as ascenders and descenders do not touch. † FDA recommends size 2,5 pt type		



Actual Size 100%

FDA00304-3

DRUG FACTS INFO Updated 5/18/19		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	6.5 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barlines*	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 30	≤ 30

* Headings are to be at least 2 pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 * FDA Recommendations 2.5pt rule



200%



Actual Size 100%

FDA00284-1

See Carton For Full Information Warnings

Allergy alert: Naproxen sodium may cause severe allergic reactions which may include: hives, facial swelling, asthma (wheezing) or shock. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 2 or more alcoholic drinks every day while using this product • take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding (red hair, vomit blood, have bloody or black stools, have stomach pain that does not get better), have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), or if pain or fever gets worse. If pregnant or breast-feeding, ask a health professional before use. **Discontinue** Adults and children 12 years and older: take 1 capsule with water every 6 to 12 hours while symptoms last. For the first dose you may take 2 capsules within the first hour. Do not exceed 3 capsules in a 24-hour period. Children under 12 years: ask a doctor. **Overdose or comments? 1-800-396-0888 (Mon-Fri 8AM-8PM EST) Fax: Dept Health Care LLC, Westbury, NY 07981**

85669470

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	-
Drug Facts Continued	8 pt	-
Headings ^a	8 pt- bold italic ^a	-
Subheadings	≥6 pt	4,5 pt
Body Text	≥6 pt	4,5 pt
Bullets	5 pt	8 pt
Leading	≥0,5 pt	4,5 pt
Hairlines	0,5 pt rule	-
Box and Barlineart	>0,5 pt rule	-
Characters Per Inch:	No more than 39	>39

^a Headings are to be at least 2pts larger than Body Text.
^{**} Drug Facts Title to be larger than all other type size used in ODP
 1 FDA Recommends 2,2pt rule



LOT & EXP
CODE AREA

Actual Size 100%

FDA00282-3

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold (italic)*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0.8 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines*	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 39	≥39

* Headings are to be at least 2 pts. longer than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DPP.
 * FDA Recommendations 2.5pt. rule



200%



Actual Size 100%

FDA00409-3

See Carton for Full Information

Warnings: Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), stomach pain, redness, rash, or blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

• take more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. Ask a doctor before use if the stomach bleeding warning applies to you, you have a history of stomach problems (such as heartburn), you have high blood pressure, heart disease, liver cirrhosis, or kidney disease, asthma, or had a stroke, you are taking a diuretic, you have problems or serious side effects from taking pain relievers or fever reducers. Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition or taking any other drug. When using this product take with food or in 1/2 stomach upset occurs. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding (belly pain, vomit blood, have bloody or black stools, have stomach pain but does not get better), have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, 1 inch below the pill is stuck in your throat, redness or swelling is present in the painful area, or any new symptoms appear.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. **Directions:** Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 caplet every 8 to 12 hours while symptoms last. For the first dose you may take 2 caplets within the first hour. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period. Children under 12 years: ask a doctor. Store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Use of this or any component? 1-800-316-1888 (Mon-Fri 9AM - 5PM EST)

Mfg. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Actual Size 100%

FDA00250-1

ICL (updated 2/27/19)		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold italic*	4.5 pt.
Subheadings	6 pt*	4.5 pt.
Body Text	6 pt*	4.5 pt.
Bullets	5 pt.	8 pt.
Hair lines	0.5 pt. rule	-
Barlines†	> 0.5 pt. rule	-
Characters per inch rule is not applicable to ICL. *ICL minimum font size is 4.5 pt. † 0.5 pt. leading may be used as long as ascenders and descenders do not touch. † FDA recommends size 2.5 pt. type		




200%



Actual Size 100%

FDA00310-1



Actual Size 100%

FDA00183-1

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch	No more than 39	≤39

* Headings are to be at least 2 pts. larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DPP.
† FDA Recommends 2.5 pt. rule.

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - the common cold
 - temporarily reduces fever

Warnings

Allergic alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or blood drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take 8 or more alcohol drinks every day while using this product
- have a history of stomach ulcers or bleeding
- are on long-term therapy with aspirin, aspirin/acetaminophen, or other NSAIDs, except aspirin.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Do not use this product if printed safety seal bearing 'Bayer HealthCare' under cap is torn or missing.

OPEN TO READ DRUG FACTS BEFORE USE

LIFT HERE

0 00000 00000 0

Drug Facts

Active Ingredient (in each caplet)

Naproxen sodium (naproxen 200 mg) (NSAID)Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 7 caplets in a 24-hour period

Children under 12 years

- ask a doctor

Other information

- each caplet contains sodium 20 mg
- store at 25°C (77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients: FDAC Plus 42 lake, hypromellose, microcrystalline cellulose, polyethylene glycol, polydioxane, talc, titanium dioxide

Questions or comments? 1-800-395-0689 (Mon-Fri 9AM - 5PM EST)



200%



Actual Size 100%

FDA00273-2



Actual Size 100%

FDA00317-1

DRUG FACTS INFO Updated 5/19/18		
CATEGORY	REQUIRE	ACTUAL IN FILE
Drug Facts Title	>8 pt **	8 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings*	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	6 pt	12 pt
Leading	≥0.5 pt	0.5 pt
Headline	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	2.5 pt
Characters Per Inch:	No more than 36	≤ 36

* Headings are to be at least 2x6 larger than body text.
 ** Drug facts file to be larger than all other type size used in DFP
 † FDA recommends 2.5pt rule






200%



Actual Size 100%

FDA00285-3




DRUG FACTS INFO <small>Updated 5/18/18</small>		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	-
Drug Facts Continued	8 pt.	-
Headings†	8 pt-bold <i>italic</i> ‡	-
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	-
Leading	≥0.5 pt.	6 pt.
Hairlines	0.5 pt rule	-
Box and Border††	>0.5 pt rule	-
Characters Per Inch	No more than 38	≤ 38

† Headings are to be at least 2x the larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 †† FDA Recommends 2.5pt rule



200%



Actual Size 100%

FDA00272-2



Actual Size 100%

FDA00401-4

DRUG FACTS INFO	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8.1 pt.
Drug Facts Content	8 pt.	8 pt.
Headings*	8 pt bold call*	8 pt.
Subheadings	>6 pt.	6 pt.
Body Text	>6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	>0.5 pt.	1 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines	>0.5 pt. rule	2.5 pt.
Characters Per Inch	No more than 30	< 30

*Headings are to be at least 2x6 larger than Body Text.
 **Drug Facts 10k to be larger than all other text used in CDP.
 † FDA Recommendations 2.5pt rule



200%



Actual Size 100%

FDA00292-1




200%



Actual Size 100%

FDA00307-3



Actual Size 100%

FDA00183-1

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch	No more than 39	≤39

* Headings are to be at least 2 pts. larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DPP.
† FDA Recommends 2.5 pt. rule.

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - the common cold
 - temporarily reduces fever

Warnings

Allergic alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or blood drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take 8 or more alcohol drinks every day while using this product
- have a history of stomach ulcers or bleeding
- have a history of heart failure, kidney disease, or liver disease

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking sulfonamide
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Do not use this product if printed safety seal bearing 'Bayer HealthCare' under cap is torn or missing.

OPEN TO READ DRUG FACTS BEFORE USE

LIFT HERE

0 00000 00000 0

Drug Facts

Active Ingredient (in each caplet)

Naproxen sodium (naproxen 200 mg) (NSAID)Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 7 caplets in a 24-hour period

- ask a doctor

Children under 12 years



- ask a doctor

Other information

- each caplet contains sodium 20 mg
- store at 25°C (77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients: FDAC Plus 42 lake, hypromellose, microcrystalline cellulose, polyethylene glycol, polydioxane, talc, titanium dioxide

Questions or comments? 1-800-395-0689 (Mon-Fri 9AM - 5PM EST)



Actual Size 100%

FDA00298-3

DRUG FACTS INFO (updated 5/18/19)	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt. bold 10a ic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	6-8.5 pt.
Hairlines	0.5 pt. rule	.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 39	≤ 39

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than a 1 other type size used in DFP
† FDA Recommends 2.5pt rule




Actual Size 100%

FDA00297-2

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0 - 0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 30	≤30

* Headings are to be at least 2pts. larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in ODP.
 † FDA Recommends 2.5pt. rule

200%




Actual Size 100%


FDA00278-4



Panel 1 Top



Panel 2 Inside



Panel 3 Base

ICL updated 2/27/19		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt- <i>bold italic</i> *	4,5 pt
Subheadings	6 pt*	4,5 pt
Body Text	6 pt*	4,5 pt
Bullets	5 pt	6 pt
Hairlines	0,5 pt rule	-
Barlines [†]	> 0,5 pt rule	-
Characters per inch rule is not applicable to ICL. *ICL minimum font size is 4,5 pt. † 0,5 pt leading may be used as long as ascenders and descenders do not touch. † FDA recommends size 2,5 pt type		



Actual Size 100%

FDA00283-4

DRUG FACTS INFO <small>updated 5/19/18</small>		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0 pt. - 0.5 pt.
Hair lines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type sizes used in ODP
 † FDA recommends 2.5pt rule



200%



Actual Size 100%

FDA00405-2



Actual Size 100%

FDA00302-2

DRUG FACTS INFO Updated 5/18/19		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	6.5 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barlines*	>0.5 pt. rule	2.5 pt.
Characters Per Inch	No more than 30	≤ 30

* Headings are to be at least 6pt. larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommendation: 2.5pt. rule



200%



Actual Size 100%

FDA00271-1

See Carton for Full Information. Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction which may include hives, facial swelling, itching (hives) or dizziness. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding (red, brown, or black stools; bloody or black stools; have stomach pain that does not get better), have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body), slurred speech, leg swelling, or if pain or fever gets worse. If pregnant or breast-feeding, ask a health professional before use. Sometimes adults and children 12 years and older: take 1 tablet with water every 6 to 12 hours while symptoms last. For the first dose you may take 2 tablets within the first hour. Do not exceed 3 tablets in a 24-hour period. Children under 12 years: ask a doctor. Overdose or overdose? 1-800-455-4688 (Moo-Fit BAW - SPIN EST)

Dist. by: Bayer HealthCare LLC
Whispern, NJ 07981

100%

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt**	-
Drug Facts Continued	8 pt	-
Headings*	8 pt-bold italic*	-
Subheadings	≥6 pt	4.5 pt
Body Text	≥6 pt	4.5 pt
Bullets	5 pt	8 pt
Leading	≥0.5 pt	0 pt
Hairlines	0.5 pt rule	-
Box and Barlines†	>0.5 pt rule	-
Characters Per Inch:	No more than 39	≥39

* Headings are to be at least 8pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in ODP
 † FDA Recommends 2.5pt rule



Actual Size 100%

FDA00309-3

DRUG FACTS INFO	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	13 pt.
Drug Facts Content	8 pt.	13 pt.
Headings*	8 pt bold call*	13 pt.
Subheadings	>6 pt.	13 pt.
Body Text	>6 pt.	13 pt.
Bullets	5 pt.	12 pt.
Leading	>0.5 pt.	0.5 pt.
Headlines	0.5 pt rule	0.5 pt.
Box and Barlines	>0.5 pt rule	2.5 pt.
Characters Per Inch	No more than 30	<30

*Headings are to be at least 2x6 larger than Body Text.
 **Drug Facts Title to be larger than all other text used in CDP.
 †FDA Recommendations 2.5pt rule






200%



Actual Size 100%

FDA00277-3



Actual Size 100%

FDA00246-2

ICL updated 2/27/18		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold italic*	6 pt
Subheadings	6 pt*	6 pt
Body Text	6 pt*	6 pt
Bullets	5 pt	12 pt
Headlines	0.5 pt rule	-
Footnotes	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
*ICL minimum font size is 4.5 pt.
< 0.5 pt leading may be used so long as ascenders and
descenders do not touch.
† FDA recommends size 2.5 pt type



Actual Size 100%

FDA00400-2

DRUG FACTS INFO (Updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>5 pt.**	13 pt.
Drug Facts Content	8 pt.	15 pt.
Headings*	8 pt bold call*	15 pt.
Subheadings	>5 pt.	15 pt.
Body Text	>5 pt.	15 pt.
Bullets	5 pt.	12 pt.
Leading	>0.5 pt.	0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines	>0.5 pt. rule	2.5 pt.
Characters Per Inch	No more than 30	<30

*Headings are to be at least 2x5 larger than Body Text.
 **Drug Facts 10k to be larger than all other text used in CDP.
 †FDA Recommendations 2.5pt rule






200%



Actual Size 100%

FDA00299-1



DRUG FACTS INFO: (Modified Format – updated 5/18/18)

CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>7.0 pt	-
Drug Facts Continued	7 pt	-
Headings*	7 pt- bold italic *	6 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	6 pt	12 pt
Leading	≥0 pt	1 pt
Hairlines	0.5 pt rule	-
Box and Barlines†	>0.5 pt rule	-
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text
 ** Drug Facts Title to be larger than all other type size used in DPP
 † Box Bar line may be omitted and color contrast used to highlight Drug Facts title



Actual Size 100%

FDA00324-3

DRUG FACTS INFO Updated 5/19/19		
CATEGORY	REQUIRE	ACTUAL IN FILE
Drug Facts Title	>8 pt **	8 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	6 pt	12 pt
Leading	≥0.5 pt	0.5 pt
Headline	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	2.5 pt
Characters Per Inch:	No more than 36	≤ 36

* Headings are to be at least 2x6 larger than body text.
 ** Drug Facts file to be larger than all other type used in DFP
 † FDA recommends 2.5pt rule






200%



Actual Size 100%


FDA00275-5



DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	-
Drug Facts Continued	8 pt	-
Headings*	8 pt- bold italic *	6 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	9 pt
Leading	≥0.5 pt	.5 pt
Hairlines	0.5 pt rule	-
Box and Barlines ¹	>0.5 pt rule	-
Characters Per Inch:	No more than 39	< 39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 1 FDA recommends 2.5pt rule

200%



Actual Size 100%

FDA00288-3

