

Tagamet<sup>®</sup> HB 200

TAKE ANYTIME YOU NEED IT:  
before meal • during meal • after meal

Tagamet<sup>®</sup> HB 200  
Cimetidine Tablets 200 mg  
Acid Reducer

JUST ONE TABLET

RELIEVES and PREVENTS  
Heartburn and Acid Indigestion

Actual size



6  
TABLETS  
(6 DOSES)

Tagamet<sup>®</sup> HB 200

Tamper Evident: Each tablet individually sealed in a blister pack. Do not use if blister or foil is open or torn.

Dist. by Medtech Products, Inc.

Tarrytown, NY 10591

A Presidio Consumer Healthcare company

©2018 Trade dress is owned by

Medtech Products, Inc. All rights reserved.

Made in Spain. T815004101



LOT:  
EXP:

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

Tips for Managing Heartburn

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Drug Facts

Active ingredient (in each tablet)	Purpose
Cimetidine 200 mg	Acid reducer

**Uses** ■ relieves heartburn associated with acid indigestion and sour stomach  
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

**Allergy alert:** Do not use if you are allergic to cimetidine or other acid reducers

**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition.

See your doctor. ■ with other acid reducers

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

**Ask a doctor or pharmacist before use if you are taking** ■ a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ you need to take this product for more than 14 days  
■ stomach pain continues ■ your heartburn continues or worsens

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water
  - to prevent symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
  - do not take more than 2 tablets in 24 hours
  - children under 12 years: ask a doctor

**Other information** store at 15°-30°C (59°-86°F)

**Inactive ingredients** cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

**Questions?** Call toll-free 1-800-482-4394 weekdays tagamet.com

10629-01-17

Font size: 9 pt  
Leading: 6 pt  
Horizontal scale: 95%  
Alignment: Left

007<sup>HB</sup>  
**Tagamet**

**TAKE ANYTIME YOU NEED IT:**  
before meal • during meal • after meal

**Tagamet**  
<sup>HB</sup> **200**  
Cimetidine Tablets 200 mg  
Acid Reducer

**JUST ONE TABLET**  
**RELIEVES and PREVENTS**  
Heartburn and Acid Indigestion

Actual size



**30**  
**TABLETS**  
(30 DOSES)

**Tips for Managing Heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

### Drug Facts

**Active ingredient (in each tablet)** Cimetidine 200 mg.....**Purpose** Acid reducer

**Uses** ■ relieves heartburn associated with acid indigestion and sour stomach  
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

### Warnings

**Allergy alert:** Do not use if you are allergic to cimetidine or other acid reducers

**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition.

See your doctor. ■ with other acid reducers

#### Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

**Ask a doctor or pharmacist before use if you are taking** ■ a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ you need to take this product for more than 14 days  
■ stomach pain continues ■ your heartburn continues or worsens

**if pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Directions

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water
  - to prevent symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
  - do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information** store at 15°-30°C (59°-86°F)

**Inactive ingredients** cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

**Questions?** Call toll-free 1-800-482-4394 weekdays tagamet.com

**Tagamet**  
<sup>HB</sup> **200**

Dist. by Medtech Products Inc., Tarrytown, NY 10591  
A Prestige Consumer Healthcare company  
©2019 Trade dress is owned by Medtech Products Inc.  
All rights reserved. Made in Spain. T615S02102  
For more information, visit tagamet.com  
Tamper Evident: Each tablet individually sealed in a blister pack. Do not use if blister or foil is open or torn.



LOT:  
EXP:

0 42037 10363 7

10628-01-17

Quantity  
Font family: Myriad Pro  
Font size: 10.51517 pt

**Tagamet<sup>®</sup> HB 200**

**+20% MORE FREE**  
**36 TABLETS FOR THE PRICE OF 30**

TAKE ANYTIME YOU NEED IT:  
 before meal · during meal · after meal

**Tagamet<sup>®</sup> HB 200**  
 Cimetidine Tablets 200 mg  
 Acid Reducer

**JUST ONE TABLET**  
 RELIEVES and PREVENTS  
 Heartburn and Acid Indigestion

Actual size



**30 36**  
**TABLETS**  
 (30 36 DOSES)

**Tagamet<sup>®</sup> HB 200**

**Just ONE TABLET RELIEVES and PREVENTS Heartburn and Acid Indigestion**

Dist. by Mediatech Products Inc. Tarrytown, NY 10591  
 A Prestige Consumer Healthcare company  
 ©2019 Trade dress is owned by Mediatech Products Inc.  
 All rights reserved. Made in Spain. TGI5001102  
 For more information, visit tagamet.com

Temporarily sealed in a blister pack. Do not use if blister or foil is open or torn.

**Tips for Managing Heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Cimetidine 200 mg	Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to cimetidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

**Ask a doctor or pharmacist before use if you are taking**

- a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- you need to take this product for more than 14 days
- stomach pain continues
- your heartburn continues or worsens

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water
  - to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn
- do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information** store at 15°-30°C (59°-86°F)

**Inactive ingredients** cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

**Questions?** Call toll-free 1-800-482-4394 weekdays tagamet.com



LOT:  
 EXP:

**Tagamet**<sup>®</sup>  
HB 200

**BETTER VALUE**  
40 PILLS for the PRICE OF 30

TAKE ANYTIME YOU NEED IT:  
before meal • during meal • after meal

**Tagamet**<sup>®</sup>  
HB 200  
Cimetidine Tablets 200 mg  
Acid Reducer

**JUST ONE TABLET**  
RELIEVES and PREVENTS  
Heartburn and Acid Indigestion

Actual size



**40**  
TABLETS  
(40 DOSES)

**Tagamet**<sup>®</sup>  
HB 200

Just ONE TABLET RELIEVES  
and PREVENTS Heartburn  
and Acid Indigestion

Tamper Evident: Each tablet individually sealed in a blister pack. Do not use if blister or pills are open or torn.

Dist. by Mediatech Products Inc., Tarrytown, NY 10591  
A Prestige Consumer Healthcare company  
©2019 Trade dress is owned by Mediatech Products Inc.  
All rights reserved. Made in Spain. T01S007103  
For more information, visit tagamet.com



LOT:  
EXP:

**Tips for Managing Heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

13270-MM-YY

**Drug Facts**

**Active ingredient (in each tablet)** Cimetidine 200 mg  
**Purpose** Acid reducer

**Uses** ■ relieves heartburn associated with acid indigestion and sour stomach  
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to cimetidine or other acid reducers

**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition.  
See your doctor. ■ with other acid reducers

**Ask a doctor before use if you have**

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

**Ask a doctor or pharmacist before use if you are taking** ■ a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ you need to take this product for more than 14 days  
■ stomach pain continues ■ your heartburn continues or worsens

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water
  - to prevent symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
  - do not take more than 2 tablets in 24 hours
  - children under 12 years: ask a doctor

**Other information** store at 15°-30°C (59°-86°F)

**Inactive ingredients** cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

**Questions?** Call toll-free 1-800-482-4394 weekdays

tagamet.com

50 Tablets  
 USA  
 TGUS009102  
 J61602A  
 0-42037-10745-1  
 PRINTS 6/0

Legal  
 Regulatory

Tagamet<sup>®</sup> HB 200  
 Just ONE TABLET RELIEVES and PREVENTS Heartburn and Acid Indigestion

Tamper Evident: Each tablet individually sealed in a blister pack. Do not use if blister or foil is open or torn.

Dist. by Mediatech Products Inc., Tarrytown, NY 10591  
 A Prestige Consumer Healthcare company  
 ©2019 Trade dress is owned by Mediatech Products Inc.  
 All rights reserved. Made in Spain. TGUS009102

**Tagamet<sup>®</sup> HB 200**  
**200<sup>HB</sup>**

**TAKE ANYTIME YOU NEED IT:**  
*before meal • during meal • after meal*

**Tagamet<sup>®</sup> HB 200**  
**Cimetidine Tablets 200 mg Acid Reducer**

**JUST ONE TABLET**  
**RELIEVES and PREVENTS**  
**Heartburn and Acid Indigestion**

Actual size  
  
**50 TABLETS (50 DOSES)**

Drug facts  
 Font family: Prestige DF  
 Font size: 9 pt  
 Leading: 6 pt  
 Horizontal scale: 95%  
 Alignment: Left

**Tips for Managing Heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Cimetidine 200 mg.....	Acid reducer

**Uses** ■ relieves heartburn associated with acid indigestion and sour stomach  
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**  
**Allergy alert:** Do not use if you are allergic to cimetidine or other acid reducers  
**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition.  
 See your doctor. ■ with other acid reducers

**Ask a doctor before use if you have**  
 ■ frequent chest pain ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ heartburn with lightheadedness, sweating or dizziness  
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
 ■ kidney disease ■ liver disease

**Ask a doctor or pharmacist before use if you are taking** ■ a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ you need to take this product for more than 14 days  
 ■ stomach pain continues ■ your heartburn continues or worsens

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**  
 ■ adults and children 12 years and over:  
 ■ to relieve symptoms, swallow 1 tablet with a glass of water  
 ■ to prevent symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn  
 ■ do not take more than 2 tablets in 24 hours  
 ■ children under 12 years: ask a doctor

**Other information** store at 15°-30°C (59°-86°F)

**Inactive ingredients** cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

**Questions?** Call toll-free 1-800-482-4394 weekdays tagamet.com

0 42037 10745 1

LOT:  
 EXP:

quantity  
 Font family: Myriad Pro  
 Font size: 12.46756 pt  
 Leading: 11 pt  
 Horizontal scale: 100%

TIPS BODY  
 Font family: Helvetica  
 Font style: Bold Condensed  
 Font size: 7 pt  
 Leading: 8 pt  
 Vertical scale: 100%

TIPS HEADER  
 Font family: Helvetica  
 Font style: Bold Condensed  
 Font size: 9 pt  
 Leading: (10.8 pt)  
 Vertical scale: 100%