

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MOVIPREP safely and effectively. See full prescribing information for MOVIPREP.

**MOVIPREP (polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution)**

**Initial U.S. Approval: 2006**

### INDICATIONS AND USAGE

MoviPrep is an osmotic laxative indicated for cleansing of the colon as a preparation for colonoscopy in adults. (1)

### DOSAGE AND ADMINISTRATION

#### Preparation and Administration:

- Two doses of MoviPrep are required for a complete preparation for colonoscopy, using a “Two-Day” preferred method or “One-Day” alternative method dosing regimen. (2.1)
- MoviPrep must be reconstituted in water prior to ingestion. (2.1)
- Additional clear liquids must be consumed after each dose of MoviPrep in both dosing regimens. (2.1, 5.1)
- Do not take other laxatives while taking MoviPrep. (2.1, 5.5)
- Do not take oral medications within 1 hour of starting each dose. (2.1)

#### Dosing Regimen:

- Two-Day (Split-Dose) (Preferred Method)** Dose 1 the evening before the colonoscopy, and Dose 2 the morning of the colonoscopy (approximately 12 hours after the start of Dose 1, and at least 3 ½ hours prior to the colonoscopy). (2.2)
- One-Day (Evening Only) (Alternative Method):** Dose 1 at least 3 ½ hours prior to bedtime the evening before the colonoscopy and Dose 2 approximately 1 ½ hours after starting Dose 1 the evening before the colonoscopy. (2.3)
- For complete information on dosing, preparation and administration see full prescribing information (2.1, 2.2, 2.3)

### DOSAGE FORMS AND STRENGTHS

For Oral Solution: 2 pouches labeled Pouch A and 2 pouches labeled Pouch B. (3)

- Each Pouch A contains 100 grams of polyethylene glycol (PEG) 3350, NF, 7.5 grams of sodium sulfate, USP, 2.691 grams of sodium chloride, USP, and 1.015 grams of potassium chloride, USP. (3)
- Each Pouch B contains 4.7 grams of ascorbic acid, USP and 5.9 grams of sodium ascorbate, USP. (3)

### CONTRAINDICATIONS

- Gastrointestinal (GI) obstruction (4, 5, 6)
- Bowel perforation (4, 4, 6)
- Gastric retention (4)

- Ileus (4)
- Toxic colitis or toxic megacolon (4)
- Hypersensitivity to any ingredient in MoviPrep (4, 5, 10)

### WARNINGS AND PRECAUTIONS

- Risk of fluid and electrolyte abnormalities:** Encourage adequate hydration, assess concurrent medications, and consider laboratory assessments prior to and after use. (5.1, 7.1)
- Cardiac arrhythmias:** Consider pre-dose and post-colonoscopy ECGs in patients at increased risk. (5.2)
- Seizures:** Use caution in patients with a history of seizures and patients at increased risk of seizure, including medications that lower the seizure threshold. (5.3, 7.1)
- Patients with renal impairment or taking concomitant medications that affect renal function:** Use caution, ensure adequate hydration and consider laboratory testing. (5.4, 7.1, 8.6)
- Colonic mucosal ulcerations:** Consider potential for ulcerations when interpreting colonoscopy findings in patients with known or suspected inflammatory bowel disease. (5.5)
- Suspected GI obstruction or perforation:** Rule out the diagnosis before administration. (5.6)
- Patients at risk for aspiration:** Observe during administration. (5.7)
- Glucose-6-phosphate dehydrogenase deficiency (G-6-PD):** Use with caution. (5.8)
- Risks in patients with phenylketonuria:** Contains phenylalanine (5.9)
- Hypersensitivity reactions, including anaphylaxis:** Inform patients to seek immediate medical care if symptoms occur. (5.10)

### ADVERSE REACTIONS

Most common adverse reactions (≥ 5%) are:

- Two-Day (Split-Dose): malaise, nausea, abdominal pain, vomiting, and upper abdominal pain. (6.1)
- One-Day (Evening-Only): abdominal distension, anal discomfort, thirst, nausea, abdominal pain, sleep disorder, rigors, hunger, malaise, vomiting, and dizziness. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Salix Pharmaceuticals, a division of Bausch Health US, LLC, at 1-800-321-4576 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

### DRUG INTERACTIONS

Drugs that increase risk for fluid and electrolyte imbalance. (7.1)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 05/2019**

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

### 2 DOSAGE AND ADMINISTRATION

- 2.1 Important Preparation and Administration Instructions
- 2.2 Two-Day Split-Dosing Regimen (Preferred Method)
- 2.3 One-Day Evening Only Dosing Regimen (Alternative Method)

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

### 5 WARNINGS AND PRECAUTIONS

- 5.1 Serious Fluid and Electrolyte Abnormalities
- 5.2 Cardiac Arrhythmias
- 5.3 Seizures
- 5.4 Use in Patients with Renal Impairment
- 5.5 Colonic Mucosal Ulceration, Ischemic Colitis and Ulcerative Colitis
- 5.6 Use in Patients with Significant Gastrointestinal Disease
- 5.7 Aspiration
- 5.8 Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency
- 5.9 Risks in Patients with Phenylketonuria
- 5.10 Hypersensitivity Reactions

### 6 ADVERSE REACTIONS

- 6.1 Clinical Studies Experience

- 6.2 Postmarketing Experience

### 7 DRUG INTERACTIONS

- 7.1 Drugs That May Increase Risks due to Fluid and Electrolyte Abnormalities
- 7.2 Potential for Reduced Drug Absorption
- 7.3 Stimulant Laxatives

### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment

### 10 OVERDOSAGE

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action

### 14 CLINICAL STUDIES

### 16 HOW SUPPLIED/STORAGE AND HANDLING

### 17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

MoviPrep® is an osmotic laxative indicated for cleansing of the colon as a preparation for colonoscopy in adults.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Important Preparation and Administration Instructions

- Correct fluid and electrolyte abnormalities before treatment with MoviPrep [see *Warnings and Precautions (5.1)*].
- Two doses of MoviPrep are required for a complete preparation for colonoscopy. The time interval between the two doses depends on the regimen prescribed and the planned timing of the colonoscopy procedure. [see *Dosage and Administration (2.2, 2.3)*].
- The “Split-Dose” is the preferred method and consists of two separate doses: the first dose is taken the evening before the colonoscopy, and the second dose is taken the next day, the morning of the day of the colonoscopy [see *Dosage and Administration (2.2)*].
- The “Evening Only” is the alternative method and consists of two separate doses: both doses are taken in the evening before the day of the colonoscopy, with a minimum of 1.5 hours between the start of the first dose and the start of the second dose [see *Dosage and Administration (2.3)*].
- Both MoviPrep dosing regimens require administration of MoviPrep using the mixing container provided to reconstitute the contents of Pouch A and B with water to the Fill Line.
- Additional clear liquids (including water) must be consumed in both dosing regimens [see *Dosage and Administration (2.2, 2.3)*, *Warnings and Precautions (5.1)*].
- Consume only clear liquids (no solid food) from the start of MoviPrep treatment until after the colonoscopy.
- Do not eat or drink alcohol, milk, anything colored red or purple or any other foods containing pulp material.
- Do not take other laxatives while taking MoviPrep [see *Drug Interactions (7.3)*].
- Do not take oral medications within 1 hour before or after starting each dose of MoviPrep [see *Drug Interactions (7.2)*].
- Ensure completion of Dose 2, including all additional liquids, at least 2 hours before the colonoscopy.
- Storage: After reconstitution, store MoviPrep solution in an upright position and keep refrigerated. Use within 24 hours after it is mixed in water.

#### 2.2 Two-Day Split-Dosing Regimen (Preferred Method)

The Two-Day Split-Dosing regimen is the preferred dosing method.

Instruct adult patients that on the day before the clinical procedure, they can consume breakfast, followed by a light lunch (no solid foods), and clear soup and/or plain yogurt for dinner, which must be completed at least 1 hour prior to the start of the first MoviPrep dose.

Instruct adult patients to take two separate doses in conjunction with fluids as follows:

Dose 1 – In the evening before the colonoscopy, approximately 10 to 12 hours before Dose 2:

1. Empty the contents of 1 Pouch A and 1 Pouch B into the mixing container that comes with MoviPrep.
2. Add lukewarm water to the Fill Line on the mixing container (32 fluid ounces). Do not add other ingredients to the MoviPrep solution.
3. Thoroughly mix with a spoon or shake with lid on securely until the contents of Pouch A and B are completely dissolved.
4. Drink 8 ounces of the solution every 15 minutes. This should take about 1 hour. Be sure to drink all the solution.

5. Refill the mixing container halfway to the Fill Line (at least 16 ounces) with a clear liquid and drink all this liquid before going to bed.

Dose 2 – Take next morning, on the day of the colonoscopy, approximately 12 hours after the start of Dose 1 and at least 3 ½ hours prior to colonoscopy:

1. Empty the contents of 1 Pouch A and 1 Pouch B into the mixing container that comes with MoviPrep.
2. Add lukewarm water to the Fill Line on the mixing container (32 fluid ounces). Do not add other ingredients to the MoviPrep solution.
3. Thoroughly mix with a spoon or shake with lid on securely until the contents of Pouch A and B are completely dissolved.
4. Drink 8 ounces of the solution every 15 minutes. This should take about 1 hour. Be sure to drink all of the solution.
5. Refill the mixing container halfway to the Fill Line (at least 16 ounces) with a clear liquid and drink all this liquid at least 2 hours before the colonoscopy.
6. Consume additional water or clear liquids up to 2 hours before the colonoscopy or as prescribed by your healthcare provider. *Then stop drinking liquids until after the colonoscopy.*

Stop drinking MoviPrep temporarily or drink each portion at longer intervals if severe bloating, abdominal discomfort or distention occurs, until these symptoms resolve.

### **2.3 One-Day Evening Only Dosing Regimen (Alternative Method)**

The One-Day Evening Only regimen is the alternative dosing method for patients for whom the Split-Dosing regimen is inappropriate.

Instruct adult patients that on the day before the clinical procedure, they can consume breakfast, followed by a light lunch (no solid foods), and clear soup and/or plain yogurt for dinner, which must be completed at least 1 hour prior to the start of the first MoviPrep dose.

Instruct adult patients to take two separate doses in conjunction with fluids as follows:

Dose 1 – At least 3 ½ hours before bedtime the evening before the colonoscopy:

1. Empty the contents of 1 Pouch A and 1 Pouch B into the mixing container that comes with MoviPrep.
2. Add lukewarm water to the Fill Line on the mixing container (32 fluid ounces). Do not add other ingredients to the MoviPrep solution.
3. Thoroughly mix with a spoon or shake with lid on securely until the contents of Pouch A and B are completely dissolved.
4. Drink 8 ounces of the solution every 15 minutes. This should take about 1 hour. Be sure to drink all the solution.

Dose 2 – At least 1 ½ hours after starting Dose 1 on the evening before the colonoscopy:

1. Empty the contents of 1 Pouch A and 1 Pouch B into the mixing container that comes with MoviPrep.
2. Add lukewarm water to the Fill Line on the mixing container (32 fluid ounces). Do not add other ingredients to the MoviPrep solution.
3. Thoroughly mix with a spoon or shake with lid on securely until the contents of Pouch A and B are completely dissolved.
4. Drink 8 ounces of the solution every 15 minutes. This should take about 1 hour. Be sure to drink all of the solution.
5. Refill the mixing container to the Fill Line (32 fluid ounces) with a clear liquid and drink all this liquid before going to bed.

6. Consume additional water or clear liquids up to 2 hours before the colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after the colonoscopy.

Stop drinking MoviPrep temporarily or drink each portion at longer intervals if severe bloating, abdominal discomfort or distention occurs, until these symptoms resolve.

### 3 DOSAGE FORMS AND STRENGTHS

MoviPrep is supplied as a white to yellow powder for reconstitution and is available in a carton that contains 2 pouches labeled Pouch A and 2 pouches labeled Pouch B.

- Each Pouch A contains 100 grams of polyethylene glycol (PEG) 3350, NF, 7.5 grams of sodium sulfate, USP, 2.691 grams of sodium chloride, USP, and 1.015 grams of potassium chloride, USP.
- Each Pouch B contains 4.7 grams of ascorbic acid, USP and 5.9 grams of sodium ascorbate, USP.

### 4 CONTRAINDICATIONS

MoviPrep is contraindicated in the following conditions:

- Gastrointestinal (GI) obstruction [*see Warnings and Precautions (5.6)*]
- Bowel perforation [*see Warnings and Precautions (5.6)*]
- Gastric retention
- Ileus
- Toxic colitis or toxic megacolon
- Hypersensitivity to any ingredient in MoviPrep [*see Warnings and Precautions (5.10)*]

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Serious Fluid and Electrolyte Abnormalities

Advise patients to hydrate adequately before, during, and after the use of MoviPrep. If a patient develops significant vomiting or signs of dehydration after taking MoviPrep, consider performing post-colonoscopy lab tests (electrolytes, creatinine, and BUN).

Bowel preparations can cause fluid and electrolyte disturbances, which can lead to serious adverse reactions including cardiac arrhythmias, seizures, and renal impairment [*see Adverse Reactions (6.2)*]. Correct fluid and electrolyte abnormalities before treatment with MoviPrep. MoviPrep should be used with caution in patients using concomitant medications that increase the risk of electrolyte abnormalities [such as diuretics, angiotensin converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARBs)] or in patients with known or suspected hyponatremia. Consider performing pre-dose and post-colonoscopy laboratory tests (sodium, potassium, calcium, creatinine, and BUN) in these patients [*see Drug Interactions (7.1)*].

#### 5.2 Cardiac Arrhythmias

There have been rare reports of serious arrhythmias (including atrial fibrillation) associated with the use of ionic osmotic laxative products for bowel preparation. These occur predominantly in patients with underlying cardiac risk factors and electrolyte disturbances. Use caution when prescribing MoviPrep for patients at increased risk of arrhythmias (e.g., patients with a history of prolonged QT, uncontrolled arrhythmias, recent myocardial infarction, unstable angina, congestive heart failure, cardiomyopathy, or electrolyte imbalance). Consider pre-dose and post-colonoscopy ECGs in patients at increased risk of serious cardiac arrhythmias.

### **5.3 Seizures**

There have been rare reports of generalized tonic-clonic seizures and/or loss of consciousness associated with use of bowel preparation products in patients with no prior history of seizures. The seizure cases were associated with electrolyte abnormalities (e.g., hyponatremia, hypokalemia, hypocalcemia, and hypomagnesemia) and low serum osmolality. The neurologic abnormalities resolved with correction of fluid and electrolyte abnormalities.

Use caution when prescribing MoviPrep for patients with a history of seizures and in patients at increased risk of seizure, such as patients taking medications that lower the seizure threshold (e.g., tricyclic antidepressants), patients withdrawing from alcohol or benzodiazepines, or patients with known or suspected hyponatremia [*see Drug Interactions (7.1)*].

### **5.4 Use in Patients with Renal Impairment**

Use MoviPrep with caution in patients with renal impairment or patients taking concomitant medications that affect renal function (such as diuretics, ACE inhibitors, angiotensin receptor blockers, or nonsteroidal anti-inflammatory drugs) [*see Drug Interactions (7.1)*]. These patients may be at risk for renal injury. Advise these patients of the importance of adequate hydration before, during, and after use of MoviPrep, and consider performing pre-dose and post-colonoscopy laboratory tests (electrolytes, creatinine, and BUN) in these patients [*see Use in Specific Populations (8.6)*].

### **5.5 Colonic Mucosal Ulceration, Ischemic Colitis and Ulcerative Colitis**

Osmotic laxatives may produce colonic mucosal aphthous ulcerations, and there have been reports of more serious cases of ischemic colitis requiring hospitalization. Concurrent use of stimulant laxatives and MoviPrep may increase the risk of mucosal ulceration or ischemic colitis and is not recommended. Consider the potential for mucosal ulcerations resulting from the bowel preparation when interpreting colonoscopy findings in patients with known or suspected inflammatory bowel disease.

### **5.6 Use in Patients with Significant Gastrointestinal Disease**

If gastrointestinal obstruction or perforation is suspected, perform appropriate diagnostic studies to rule out these conditions before administering MoviPrep [*see Contraindications (4)*].

Use with caution in patients with severe ulcerative colitis.

### **5.7 Aspiration**

Patients with impaired gag reflex or other swallowing abnormalities are at risk for regurgitation or aspiration of MoviPrep. Observe these patients during the administration of MoviPrep. Use with caution in these patients.

### **5.8 Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency**

Since MoviPrep contains sodium ascorbate and ascorbic acid, MoviPrep should be used with caution in patients with glucose-6-phosphate dehydrogenase (G6PD) deficiency, especially G6PD deficiency patients with an active infection, with a history of hemolysis, or taking concomitant medications known to precipitate hemolytic reactions.

### **5.9 Risks in Patients with Phenylketonuria**

Phenylalanine can be harmful to patients with phenylketonuria (PKU). MoviPrep contains phenylalanine, a component of aspartame. Each MoviPrep treatment contains 131 mg of phenylalanine (after hydrolysis of the aspartame molecule *in vivo* to aspartic acid and phenylalanine). Before prescribing MoviPrep to a patient with PKU, consider the combined daily amount of phenylalanine from all sources, including MoviPrep.

## 5.10 Hypersensitivity Reactions

MoviPrep contains polyethylene glycol (PEG) and may cause serious hypersensitivity reactions including anaphylaxis, angioedema, rash, urticaria, and pruritus [see *Adverse Reactions (2)*]. Inform patients of the signs and symptoms of anaphylaxis, and instruct them to seek immediate medical care should signs and symptoms occur.

## 6 ADVERSE REACTIONS

The following serious or otherwise important adverse reactions for bowel preparations are described elsewhere in the labeling:

- Serious Fluid and Electrolyte Abnormalities [see *Warnings and Precautions (5.1)*]
- Cardiac Arrhythmias [see *Warnings and Precautions (5.2)*]
- Seizures [see *Warnings and Precautions (5.3)*]
- Patients with Renal Impairment [see *Warnings and Precautions (5.4)*]
- Colonic Mucosal Ulceration, Ischemic Colitis and Ulcerative Colitis [see *Warnings and Precautions (5.5)*]
- Patients with Significant Gastrointestinal Disease [see *Warnings and Precautions (5.6)*]
- Aspiration [see *Warnings and Precautions (5.7)*]
- Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency [see *Warnings and Precautions (5.8)*]
- Risks in Patients with Phenylketonuria [see *Warnings and Precautions (5.9)*]
- Hypersensitivity Reactions [see *Warnings and Precautions (5.10)*]

### 6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The safety of MoviPrep as a Two-Day Split-Dosing and One-Day Evening Only Dosing Regimen was evaluated in two randomized, active-controlled, multicenter, investigator-blinded clinical trials in adult patients scheduled to have an elective colonoscopy [see *Clinical Studies (14)*]. The safety analysis for Study 1 included 359 adult patients ranging in age from 18 to 88 years (mean age 59), with 52% female and 48% male patients. The safety analysis for Study 2 included 340 adult patients ranging in age from 21 to 76 years (mean age 53), with 53% male and 47% female patients.

Tables 1 and 2 display adverse reactions reported in at least 2% and 5% of patients in either treatment group in Study 1 and Study 2, respectively. Since diarrhea was considered as a part of the efficacy assessment, it was not defined as an adverse reaction in these trials.

**Table 1: Common Adverse Reactions<sup>1</sup> in Patients Undergoing Colonoscopy in Study 1**

	<b>MoviPrep Two-Day Split Dosing Regimen (N=180)</b>	<b>4 Liter PEG + Electrolytes Solution (N=179)</b>
Malaise	19%	18%
Nausea	14%	20%
Abdominal pain	13%	15%
Vomiting	8%	13%
Upper abdominal pain	6%	6%
Dyspepsia	3%	1%

<sup>1</sup> Reported in at least 2% of patients in either treatment group

**Table 2: Common Adverse Reactions<sup>1,2</sup> in Patients Undergoing Colonoscopy in Study 2**

	<b>MoviPrep One-Day Evening Only Dosing Regimen(N=169)</b>	<b>90 mL Oral Sodium Phosphate Solution (N=171)</b>
Abdominal distension	60%	41%
Anal discomfort	51%	52%
Thirst	47%	65%
Nausea	47%	47%
Abdominal pain	39%	32%
Sleep disorder	35%	29%
Rigors	34%	30%
Hunger	30%	71%
Malaise	27%	53%
Vomiting	7%	8%
Dizziness	7%	18%
Headache	2%	5%
Hypokalemia	0%	6%
Hyperphosphatemia	0%	6%

<sup>1</sup> Reported in at least 5% of patients in either treatment group

<sup>2</sup> Patients were specifically asked about the occurrence of the following symptoms: shivering, anal irritations, abdominal bloating or fullness, sleep loss, nausea, vomiting, weakness, hunger sensation, abdominal cramps or pain, thirst sensation, and dizziness.

## 6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of MoviPrep or other PEG-based products. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Cardiovascular:* Tachycardia, palpitations, hypertension, arrhythmia, atrial fibrillation, peripheral edema, asystole, acute pulmonary edema and syncope, and dehydration.

*Gastrointestinal:* upper gastrointestinal bleeding from a Mallory-Weiss tear, esophageal perforation [usually with gastroesophageal reflux disease (GERD)]

*Hypersensitivity reactions:* anaphylaxis (some of which were severe, including shock), rash, urticaria, pruritus, lip, tongue and facial swelling, dyspnea, chest tightness and throat tightness, rhinorrhea, dermatitis, fever, and chills.

*Nervous system:* tremor, seizure.

*Renal:* renal impairment and/or failure.

## 7 DRUG INTERACTIONS

### 7.1 Drugs That May Increase Risks due to Fluid and Electrolyte Abnormalities

Use caution when prescribing MoviPrep for patients with conditions and/or who are using medications that increase the risk for fluid and electrolyte disturbances or may increase the risk of renal impairment, seizures, arrhythmias, or QT prolongation in the setting of fluid and electrolyte abnormalities [see *Warnings and Precautions (5.1, 5.2, 5.3, 5.4)*]. Consider additional patient evaluations as appropriate.

### 7.2 Potential for Reduced Drug Absorption

MoviPrep can reduce the absorption of other co-administered drugs. Administer oral medications at least 1 hour before the start of administration of each dose of MoviPrep [see *Dosage and Administration (2.1)*].

### 7.3 Stimulant Laxatives

Concurrent use of stimulant laxatives and MoviPrep may increase the risk of mucosal ulceration or ischemic colitis. Avoid use of stimulant laxatives (e.g., bisacodyl, sodium picosulfate) while taking MoviPrep [see *Warnings and Precautions (5.5, 5.6)*].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

There are no available data on MoviPrep in pregnant women to inform a drug-associated risk for adverse developmental outcomes. Animal reproduction studies have not been conducted with MoviPrep.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

### 8.2 Lactation

#### Risk Summary

There are no data available on the presence of MoviPrep in human milk, the effects of the drug on the breastfed child, or the effects of the drug on milk production. The lack of clinical data during lactation precludes a clear determination of the risk of MoviPrep to a child during lactation; therefore, the developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for MoviPrep and any potential adverse effects on the breastfed child from MoviPrep or from the underlying maternal condition.

### 8.4 Pediatric Use

The safety and effectiveness of MoviPrep in pediatric patients have not been established.

### 8.5 Geriatric Use

Of the 413 patients in clinical trials receiving MoviPrep, 91 (22%) patients were aged 65 or older, while 25 (6%) patients were over 75 years of age. No overall differences in safety or effectiveness were observed between geriatric patients and younger patients, and other reported clinical experience has not identified differences in responses between geriatric patients and younger patients. However, elderly patients are more likely to have decreased hepatic, renal or cardiac

function and may be more susceptible to adverse reactions resulting from fluid and electrolyte abnormalities [*see Warnings and Precautions (5.1)*].

## 8.6 Renal Impairment

Use MoviPrep with caution in patients with renal impairment or patients taking concomitant medications that may affect renal function [*see Drug Interactions (7.1)*]. These patients may be at risk for renal injury. Advise these patients of the importance of adequate hydration before, during and after the use of MoviPrep, and consider performing baseline and post-colonoscopy laboratory tests (electrolytes, creatinine, and BUN) in these patients [*see Warnings and Precautions (5.4)*].

## 10 OVERDOSAGE

Overdosage of more than the recommended dose of MoviPrep may lead to severe electrolyte disturbances, including hyponatremia and/or hypokalemia, as well as dehydration and hypovolemia, with signs and symptoms of these disturbances. Certain severe electrolyte disturbances may lead to cardiac arrhythmias, seizures, and renal failure [*see Warnings and Precautions (5.1, 5.2, 5.3)*]. Monitor for fluid and electrolyte disturbances and treat symptomatically.

## 11 DESCRIPTION

MoviPrep (PEG-3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution) is an osmotic laxative consisting of 4 pouches (2 of Pouch A and 2 of Pouch B) containing white to yellow powder for reconstitution.

Each Pouch A contains 100 grams of polyethylene glycol (PEG) 3350, NF, 7.5 grams of sodium sulfate, USP, 2.691 grams of sodium chloride, USP, and 1.015 grams of potassium chloride, USP, plus the following excipients: aspartame, NF (sweetener), acesulfame potassium, NF (sweetener), and lemon flavoring. Pouch A contains 111.9 g of powder for oral solution.

Each Pouch B contains 4.7 grams of ascorbic acid, USP and 5.9 grams of sodium ascorbate, USP. Pouch B contains 10.6 g of powder for oral solution.

When 1 Pouch A and 1 Pouch B are dissolved together in water to a volume of 1 liter, MoviPrep (PEG-3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid) is an oral solution having a lemon taste.

The entire, reconstituted, 2-liter MoviPrep colon preparation contains 200 grams of PEG-3350, 15 grams of sodium sulfate, 5.38 grams of sodium chloride, 2.03 grams of potassium chloride, 9.4 grams of ascorbic acid, and 11.8 grams of sodium ascorbate plus the following excipients: aspartame (sweetener), acesulfame potassium (sweetener), and lemon flavoring.

A mixing container for reconstitution is enclosed.

Phenylketonurics: Contains Phenylalanine 131 mg per treatment.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The primary mode of action is osmotic action of polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid, which induce a laxative effect. The physiological consequence is increased water retention in the lumen of the colon, resulting in loose stools.

## 14 CLINICAL STUDIES

The colon cleansing efficacy and safety of MoviPrep was evaluated in two randomized, actively-controlled, multi-center, investigator-blinded trials in adult patients scheduled to have an elective colonoscopy.

In Study 1, patients were randomized to one of the following two colon preparation treatments: 1) 2 liters of MoviPrep with 1 additional liter of clear liquid split into two doses (during the evening before and the morning of the colonoscopy) and 2) 4 liters of polyethylene glycol plus electrolytes solution (4L PEG + E) split into two doses (during the evening before and the morning of the colonoscopy). Patients were allowed to have a morning breakfast, a light lunch, clear soup and/or plain yogurt for dinner. Dinner had to be completed at least one hour prior to initiation of the colon preparation administration.

The primary efficacy endpoint was the proportion of patients with effective colon cleansing as judged by blinded gastroenterologists on the basis of videotapes recorded during the colonoscopy.

The blinded gastroenterologists graded the colon cleansing twice (during introduction and withdrawal of the colonoscope) and the poorer of the two assessments was used in the primary efficacy analysis.

The efficacy analysis included 308 adult patients who had an elective colonoscopy. Patients ranged in age from 18 to 88 years old (mean age about 59 years old) with 52% female and 48% male patients. Table 3 displays the results.

**Table 3: Effectiveness of Overall Colon Cleansing of MoviPrep vs. 4 Liter Polyethylene Glycol plus Electrolytes Solution in Study 1**

	<b>Responders A<sup>2</sup> or B<sup>3</sup> (%)</b>	<b>C<sup>4</sup> (%)</b>	<b>D<sup>5</sup> (%)</b>
<b>MoviPrep (N=153)</b>	88.9	9.8	1.3
<b>4L PEG + E<sup>1</sup> (N=155)</b>	94.8	4.5	0.6

<sup>1</sup> 4L PEG + E is 4 Liter Polyethylene Glycol plus Electrolytes Solution.

<sup>2</sup> A: colon empty and clean or presence of clear liquid, but easily removed by suction

<sup>3</sup> B: brown liquid or semisolid remaining amounts of stool, fully removable by suction or displaceable, thus allowing a complete visualization of the gut mucosa

<sup>4</sup> C: semisolid amounts of stool, only partially removable with a risk of incomplete visualization of the gut mucosa

<sup>5</sup> D: semisolid or solid amounts of stool; consequently colonoscopy incomplete or needed to be terminated.

4L PEG+E's responder rate was not significantly higher than MoviPrep's responder rate.

In Study 2, patients were randomized to one of the following two colon preparation treatments: 1) 2 liters of MoviPrep with 1 additional liter of clear liquid in the evening prior to the colonoscopy and 2) 90 mL of oral sodium phosphate solution (90 mL OSPS) with at least 2 liters of additional clear liquid during the day and evening prior to the colonoscopy. Patients randomized to MoviPrep therapy were allowed to have a morning breakfast; a light lunch; and clear soup and/or plain yogurt for dinner. Dinner had to be completed at least one hour prior to initiation of the colon preparation administration.

The primary efficacy endpoint was the proportion of patients with effective colon cleansing as judged by the colonoscopist and one blinded gastroenterologist (on the basis of videotapes recorded during the colonoscopy). In case of a discrepancy between the colonoscopist and the blinded gastroenterologist, a second blinded gastroenterologist made the final efficacy determination.

The efficacy analysis included 280 adult patients who had an elective colonoscopy. Patients ranged in age from 21 to 76 years old (mean age about 53 years old) with 47% female and 53% male patients. Table 4 displays the results.

**Table 4: Effectiveness of Overall Colon Cleansing of MoviPrep Vs. 90 mL Oral Sodium Phosphate Solution in Study 2**

	<b>Responders A<sup>2</sup> or B<sup>3</sup> (%)</b>	<b>C<sup>4</sup> (%)</b>	<b>D<sup>5</sup> (%)</b>
<b>MoviPrep (N=137)</b>	73.0	23.4	3.6
<b>90 mL OSPS<sup>1</sup> (N=143)</b>	64.4	29.4	6.3

<sup>1</sup> OSPS is Oral Sodium Phosphate Solution.

<sup>2</sup> A: empty and clean or clear liquid (transparent, yellow, or green)

<sup>3</sup> B: brown liquid or semisolid remaining small amounts of stool, fully removable by suction or displaceable allowing a complete visualization of the underlying mucosa

<sup>4</sup> C: semisolid only partially removable/displaceable stools; risk of incomplete examination of the underlying mucosa

<sup>5</sup> D: heavy and hard stool making the segment examination uninterpretable and, consequently, the colonoscopy needed to be terminated.

MoviPrep's responder rate was not significantly higher than OSPS's responder rate.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

MoviPrep (polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution) is supplied as a white to yellow powder for reconstitution.

NDC 65649-201-75, MoviPrep, single-use outer carton:

- Each outer carton contains a disposable mixing container with lid for reconstitution of MoviPrep, prescribing information and patient information, and one inner carton.
- Each inner carton contains 2 pouches labeled Pouch A and 2 pouches labeled Pouch B.

### Storage

Store carton/container at room temperature, between 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F). When reconstituted, store upright and keep solution refrigerated. Use within 24 hours [see *Dosage and Administration (2.1)*].

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide and Instructions for Use).

Instruct patients:

- That two doses of MoviPrep are required for a complete preparation for colonoscopy either as a Split-Dose (2-Day), or Evening Only (1-Day) dosing regimen [see *Instructions for Use*].
- Not to take other laxatives while they are taking MoviPrep.
- That MoviPrep contains 131 mg of phenylalanine per treatment [see *Warnings and Precautions (5.9)*].
- That each pouch needs to be reconstituted in water before ingestion and that they should drink additional clear liquids. Examples of clear liquids can be found in the *Instructions for Use*.
- Not to take oral medications within one hour of starting each dose of MoviPrep.
- To follow the directions in the *Instructions for Use*, for either the Two-Day Split-Dosing or the One-Day Evening Only Dosing regimen, as prescribed.

- To consume additional clear liquids before, during, and after the use of MoviPrep to prevent dehydration [*see Warnings and Precautions (5.1)*].
- To contact their healthcare provider if they develop significant vomiting or signs of dehydration after taking MoviPrep or if they experience altered consciousness or seizures [*see Warnings and Precautions (5.1, 5.2, 5.3, 5.4)*]
- Not to eat or drink alcohol, milk, anything colored red or purple or any other foods containing pulp material.
- To stop drinking MoviPrep temporarily or drink each portion at longer intervals if they develop severe abdominal discomfort or distention until these symptoms diminish. If severe symptoms persist, tell patients to contact their healthcare provider.

**Distributed by:**

Salix Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

U.S. Patent Numbers: 7,169,381 and 7,658,914

MoviPrep is a trademark of Velinor AG used under license.

© 2019 Bausch Health Companies Inc. or its affiliates

## MEDICATION GUIDE

### MoviPrep® (moo-vee-prěp)

(polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution)

**Read this Medication Guide and Instructions for Use before your colonoscopy and again before you start taking MoviPrep.**

**What is the most important information I should know about MoviPrep?**

**MoviPrep and other bowel preparations can cause serious side effects, including:**

- **Serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood. These changes can cause:**
  - abnormal heartbeats that can cause death.
  - seizures. This can happen even if you have never had a seizure.
  - kidney problems.

**Your chance of having fluid loss and changes in body salts with MoviPrep is higher if you:**

- have heart problems.
- have kidney problems.
- take water pills (diuretics), high blood pressure medicine, or non-steroidal anti-inflammatory drugs (NSAIDs).

**Tell your healthcare provider right away if you have any of these symptoms of serious loss of body fluid (dehydration) while taking MoviPrep:**

- vomiting
- dizziness
- urinating less often than normal
- headache

**See “What are the possible side effects of MoviPrep?” for more information about side effects.**

**What is MoviPrep?**

MoviPrep is a prescription medicine used by adults to clean the colon before a colonoscopy. MoviPrep cleans your colon by causing you to have diarrhea (loose stools). Cleaning your colon helps your healthcare provider see the inside of your colon more clearly during your colonoscopy.

It is not known if MoviPrep is safe and effective in children.

**Who should not take MoviPrep?**

Do not take MoviPrep if your healthcare provider has told you that you have:

- a blockage in your intestine (bowel obstruction).
- an opening in the wall of your stomach or intestine (bowel perforation).
- problems with food and fluid emptying from your stomach (gastric retention).
- a problem with food moving too slowly through your intestines (ileus).
- a very dilated intestine (toxic megacolon).
- an allergy to any of the ingredients in MoviPrep. See the end of this leaflet for a complete list of ingredients in MoviPrep.

**What should I tell my healthcare provider before taking MoviPrep?**

Before taking MoviPrep, tell your healthcare provider **about all of your medical conditions, including** if you:

- have problems with serious loss of body fluid (dehydration) and changes in blood salts (electrolytes).
- have heart problems.
- have seizures or take medicines for seizures.
- have kidney problems or take medicines for kidney problems.
- have stomach or bowel problems, including ulcerative colitis.
- have problems with swallowing, gastric reflux, or if you inhale food or fluid into your lungs when eating or drinking (aspirate).
- have a condition called glucose-6-phosphate dehydrogenase (G6PD) deficiency that destroys red blood cells.
- are withdrawing from alcohol or benzodiazepines.
- have phenylketonuria (PKU). MoviPrep contains phenylalanine.
- are allergic to any of the ingredients in MoviPrep.

- are pregnant or plan to become pregnant. It is not known if MoviPrep will harm your unborn baby. Talk to your healthcare provider if you are pregnant.
- are breastfeeding or plan to breastfeed. It is not known if MoviPrep passes into your breast milk. You and your healthcare provider should decide if you will take MoviPrep while breastfeeding.

**Tell your healthcare provider about all the medicines you take**, including prescription and non-prescription medicines, vitamins, and herbal supplements.

**MoviPrep may affect how other medicines work.** Do not take medicines by mouth 1 hour before or after the start of MoviPrep.

Especially tell your healthcare provider if you take:

- medicines to treat a blood salt (electrolyte) imbalance.
- medicines for blood pressure or heart problems.
- medicines for seizures (antiepileptics).
- medicines for kidney problems.
- water pills (diuretics).
- non-steroidal anti-inflammatory drugs (NSAID).
- laxatives. Do not take other laxatives while taking MoviPrep.
- medicines for depression or other mental health problems.

Ask your healthcare provider or pharmacist for a list of these medicines if you are not sure if you are taking any of the medicines listed above.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

### **How should I take MoviPrep?**

**See the “Instructions for Use” for dosing instructions.** You must read, understand, and follow these instructions to take MoviPrep the right way.

- Take MoviPrep exactly as your healthcare provider tells you to take it. Your healthcare provider will tell you to take the Two-Day Split-Dosing regimen option or the One-Day Evening Only Dosing regimen option.
- On the day before the procedure, you can have breakfast, followed by a light lunch (no solid foods), and dinner of clear soup with or without plain yogurt, or plain yogurt only. You must finish dinner at least 1 hour before the start of the first MoviPrep dose.
- Drink only clear liquids before, during, and after you take MoviPrep, until 2 hours before your colonoscopy to help prevent fluid loss (dehydration).
- Do not eat solid food while taking MoviPrep until after your colonoscopy.
- Do not eat or drink alcohol, milk, anything colored red or purple or any other foods with pulp.
- It is important for you to drink the additional amount of clear liquids listed in the **Instructions for Use**.
- You may have stomach-area (abdomen) bloating after your first dose of MoviPrep.
  - If you have severe stomach-area (abdomen) discomfort or bloating, stop drinking MoviPrep for a short time or wait a longer time between each dose of MoviPrep until your stomach-area symptoms improve. If your stomach-area discomfort or bloating continues, tell your healthcare provider.
- If you take too much MoviPrep, call your healthcare provider.

### **What are the possible side effects of MoviPrep?**

**MoviPrep can cause serious side effects, including:**

- **Changes in certain blood tests.** Your healthcare provider may do blood tests after you take MoviPrep to check your blood for changes. Tell your healthcare provider if you have any symptoms of too much fluid loss, including:
  - vomiting
  - dizziness
  - feel faint, weak or lightheaded especially when you stand up (orthostatic hypotension)
  - heart problems
  - kidney problems
  - seizures
  - dry mouth
- Ulcers of the bowel or bowel problems (ischemic colitis): Tell your healthcare provider right away if you have severe stomach-area (abdomen) pain or rectal bleeding.
- Serious allergic reactions. Symptoms of a serious allergic reaction may include:

- skin rash
- swelling of the face, lips, tongue and throat
- raised red patches on your skin (hives)
- itching
- kidney problems

**The most common side effects of MoviPrep include:**

- anal discomfort
- thirst
- nausea
- stomach-area (abdomen pain or bloating)
- sleep problems
- chills
- hunger
- discomfort
- vomiting
- indigestion
- dizziness

These are not all the possible side effects of MoviPrep.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store MoviPrep?**

- Store MoviPrep that has not been mixed with water at room temperature, between 68° to 77°F (20° to 25°C).
- Store MoviPrep that has been mixed with water in an upright position in the refrigerator.
- MoviPrep should be taken within 24 hours after it has been mixed with water.

**Keep MoviPrep and all medicines out of the reach of children.**

**General information about the safe and effective use of MoviPrep.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use MoviPrep for a condition for which it was not prescribed. Do not give MoviPrep to other people, even if they are going to have the same procedure you are. It may harm them.

This Medication Guide summarizes the most important information about MoviPrep. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information that is written for healthcare professionals.

For more information, call 1-800-321-4576 or go to [www.MoviPrep.com](http://www.MoviPrep.com).

**What are the ingredients in MoviPrep?**

**Active ingredients:**

Pouch A: polyethylene glycol (PEG) 3350, sodium sulfate, sodium chloride, potassium chloride.

Pouch B: ascorbic acid and sodium ascorbate.

**Inactive ingredients:**

Pouch A: aspartame, acesulfame potassium, and lemon flavoring.

**Distributed by:**

Salix Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA  
U.S. Patent Numbers: 7,169,381 and 7,658,914

© 2019 Bausch Health Companies Inc. or its affiliates

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Issued: May 2019

## INSTRUCTIONS FOR USE

### MoviPrep® (polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution)

There are two different options for taking MoviPrep. Your healthcare provider will tell you to take the Two-Day Split-Dosing Regimen option **or** the One-Day Evening Only Dosing Regimen option.

#### Important Information on MoviPrep:

- You must drink all of the Dose 1 and Dose 2 of MoviPrep with either dosing regimen option. Make sure you finish Dose 2 at least 2 hours before your colonoscopy.
- After completing Dose 2, it is important that you drink additional clear liquids (including water), but you must stop drinking all liquids at least 2 hours before your colonoscopy.
- MoviPrep **must** be mixed with water. **Do not** add any other ingredients to MoviPrep.
- After you start taking MoviPrep, you can only drink clear liquids (no solid foods) until after your colonoscopy. Examples of clear liquids include:
  - water
  - clear fruit juices without pulp including apple, white grape, or white cranberry
  - strained limeade or lemonade
  - coffee or tea (**do not** use any dairy or non-dairy creamer)
  - clear broth
  - clear soda
  - gelatin (without added fruit or topping, no red or purple)
  - popsicles (without pieces of fruit or pulp, no red or purple)
- Drink plenty of clear liquids before, during, and after you take MoviPrep, up until 2 hours before your colonoscopy, to help prevent fluid loss (dehydration).
- **Do not** eat or drink anything within 2 hours before your colonoscopy.
- **Do not** eat or drink alcohol, milk, anything colored red or purple or containing pulp.
- **Do not** take other laxatives while taking MoviPrep.
- **Do not** take any medicines by mouth (oral) within 1 hour before or after starting each dose of MoviPrep.
- **Do not** eat any solid food while taking MoviPrep until after your colonoscopy.

To take each dose of MoviPrep, you will need:


- Mixing Container that comes with MoviPrep
- One Pouch A
- One Pouch B
- Lukewarm water
- For the Two-Day Split-Dosing Regimen:
  - On the day before the colonoscopy you can eat breakfast followed by a light lunch (no solid foods). For dinner you may have a clear soup with or without plain yogurt, or plain yogurt only.
  - You must finish eating at least 1 hour before you start taking MoviPrep.
  - You must take the first dose between 10 to 12 hours before the second dose. The second dose must be taken at least 3 ½ hours before the colonoscopy.
  - After you start taking MoviPrep you can only drink clear liquids.
- For the One-Day Evening Only Dosing Regimen:
  - On the day before the colonoscopy you can eat breakfast followed by a light lunch. For dinner you may have a clear soup with or without plain yogurt, or plain yogurt only.
  - You must finish eating at least 1 hour before you start taking MoviPrep.

- You must take the first dose 3 ½ hours before bedtime the evening before the colonoscopy.
- After you start taking MoviPrep you can only drink clear liquids.
- Take the second dose 1 ½ hours after starting Dose 1.


### **Two-Day Split-Dosing Regimen Dosing Instructions**

#### **Dose 1 – Take this dose the evening before your colonoscopy (10 to 12 hours before Dose 2):**


Step 1: Empty the contents of **one** Pouch A and **one** Pouch B into the Mixing Container that comes with MoviPrep.



Step 2: Add lukewarm water to the Fill Line on the Mixing Container. You will need at least 32 ounces.



Step 3: Mix to completely dissolve the medicine from Pouch A and Pouch B into the lukewarm water. To mix the solution, stir the medicine in the Mixing Container with a spoon, or close the lid and shake.




Step 4: Drink one 8 oz. (ounce) glass of the solution every 15 minutes. Be sure to drink all of the solution in the Mixing Container. It should take about 1 hour to drink all the liquid.

If you feel like you have severe stomach pain or discomfort you can stop taking MoviPrep for a short period of time and then continue taking it or you can take smaller sips of MoviPrep so that you space out your dose longer than 1 hour. If you still have severe stomach pain, call your healthcare provider.




Step 5: Refill the Mixing Container with 16 oz. (at least halfway to Fill Line and enough for two 8 oz. glasses) of a clear liquid and drink all of this liquid before you go to bed.



**Dose 2 – Take this dose the next morning on the day of your colonoscopy (start at least 3 ½ hours before your colonoscopy):**

Step 1: Repeat steps 1 through 4 from Dose 1 of the Split-Dose (2-Day) instructions.

Step 2: Fill the Mixing Container with 16 oz. (at least halfway to Fill Line and enough for two 8 oz. glasses) of a clear liquid and drink all of this liquid at least 2 hours before your colonoscopy.




Step 3: Drink only clear liquids up to 2 hours before your colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after your colonoscopy.

**One-Day Evening Only Dosing Regimen Instructions**

**Dose 1 (take at least 3 ½ hours before bedtime the evening before your colonoscopy):**


Step 1: Empty the contents of **one** Pouch A and **one** Pouch B into the Mixing Container that comes with MoviPrep.



Step 2: Add lukewarm water to the Fill Line on the Mixing Container. You will need at least 32 ounces.



Step 3: Mix to completely dissolve the medicine from Pouch A and Pouch B into the lukewarm water. To mix the solution, stir the medicine in the Mixing Container with a spoon, or close the lid and shake.



Step 4: Drink one 8 oz. (ounce) glass of the solution every 15 minutes. Be sure to drink all of the solution in the Mixing Container. It should take about 1 hour to drink all the liquid.




If you feel like you have severe stomach pain or discomfort you can stop taking MoviPrep for a short period of time and then continue taking it or you can take smaller sips of MoviPrep so that you space out your dose longer than 1 hour. If you still have severe stomach pain, call your healthcare provider.

**Dose 2 (take about 1 ½ hours after starting Dose 1):**

Step 1: Repeat Steps 1 through 4 from Dose 1 of the Evening Only (1-Day) instructions.

Step 2: After you complete steps 1 through 4, fill the Mixing Container again to the Fill Line with clear liquid and drink all of this liquid before you go to bed.



Step 3: Drink only clear liquids up to 2 hours before your colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after your colonoscopy.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

**Distributed by:**

Salix Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

U.S. Patent Numbers: 7,169,381 and 7,658,914  
© 2019 Bausch Health Companies Inc. or its affiliates

Issued: May 2019

675

**INSTRUCTIONS - See full Prescribing Information and Instructions For Use.**

**Important Information:**

- Take Moviprep as prescribed by your healthcare provider.
- The contents of each pouch of Moviprep **must** be mixed with water before taking.
- **Do not eat any solid food while taking Moviprep until after your colonoscopy.**
- Drink only clear liquids:
  - while taking Moviprep
  - after taking Moviprep and until 2 hours before your colonoscopy or as prescribed by your healthcare provider

**You may eat or drink any of the following clear liquids:**

- water
- clear fruit juices without pulp including apple, white grape, or white cranberry
- strained limeade or lemonade
- coffee or tea (**do not** use any dairy or non-dairy creamer.)
- clear broth
- clear soda
- gelatin (without added fruit or topping)
- popsicles (without pieces of fruit or fruit pulp)

**Do not eat or drink anything colored red or purple.**

**Do not eat or drink anything 2 hours before your colonoscopy or as prescribed by your healthcare provider.**

**Store at room temperature between 20° to 25°C (68° to 77°F).**

**Moviprep solution is taken by mouth in one of the following ways:**

**Two-Day Split-Dosing Instructions:**

**Dose 1 – Take this dose the evening before your colonoscopy (10 to 12 hours before Dose 2):**

1. Empty the contents of 1 Pouch A and 1 Pouch B into the container that comes with Moviprep.
2. Add lukewarm water to the Fill Line on the container.
3. Mix to completely dissolve the contents of Pouch A and B into the lukewarm water.
4. Drink one 8 oz. (ounce) glass (240 mL) every 15 minutes. Be sure to drink all of the solution. This should take about 1 hour.
5. Fill the container with 16 oz. (two 8 oz. glasses) of clear liquid and drink all of this liquid before you go to bed.

**Dose 2 – Take this dose the next morning on the day of your colonoscopy (start at least 3 ½ hours before your colonoscopy):**

1. Repeat steps 1 through 4 from Dose 1 of the Two-Day Split-Dosing instructions.
2. Fill the container with 16 oz. (two 8 oz. glasses) of clear liquid and drink all of this liquid at least 2 hours before your colonoscopy.
3. Drink only clear liquids up to 2 hours before your colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after your colonoscopy.

**Or**

**One-Day Evening Only Instructions:**

**Dose 1 – Take this dose at least 3 ½ hours before bedtime the evening before your colonoscopy:**

1. Empty the contents of 1 Pouch A and 1 Pouch B into the container that comes with Moviprep.
2. Add lukewarm water to the Fill Line on the container.
3. Mix to completely dissolve the contents of Pouch A and B into the lukewarm water.
4. Drink one 8 oz. (ounce) glass (240 mL) every 15 minutes. Be sure to drink all of the solution. This should take about 1 hour.

**Dose 2 – Take Dose 2 about 1 ½ hours after starting Dose 1 the evening before your colonoscopy:**

1. Repeat steps 1 through 4 from Dose 1 of the One-Day Evening Only instructions.
2. After you complete steps 1 through 4, fill the container again to the Fill Line with clear liquid and drink all of this liquid before you go to bed.
3. Drink only clear liquids up to 2 hours before your colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after your colonoscopy.

**Moviprep solution may be refrigerated before drinking.** Keep the mixed solution in an upright position and store in the refrigerator. Moviprep should be taken within 24 hours after it is mixed in water. Do not add other ingredients to the Moviprep solution.



Moviprep is a trademark of Velinor AG used under license.

**Rx only**

**Moviprep**

(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride, Sodium Ascorbate and Ascorbic Acid for Oral Solution)

100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g

Lot:  
Exp:

9675400 10054110



PEG-3350	100 g
Sodium sulfate	7.5 g
Sodium chloride	2.691 g
Potassium chloride	1.015 g
Chloride	59.8 mmol/L
Sulfate	52.8 mmol/L
Potassium	14.2 mmol/L
Ascorbate	29.8 mmol/L

Each Pouch A contains:	100 g PEG-3350
	7.5 g Sodium sulfate
	2.691 g Sodium chloride
	1.015 g Potassium chloride
	59.8 mmol/L Chloride
	52.8 mmol/L Sulfate
	14.2 mmol/L Potassium
	29.8 mmol/L Ascorbate

Each Pouch B contains:	5.9 g Sodium ascorbate
	4.7 g Ascorbic acid

also contains aspartame	1.015 g Potassium chloride
	2.691 g Sodium chloride
	7.5 g Sodium sulfate
	100 g PEG-3350

This carton contains: 2 Pouches labeled Pouch A, and 2 Pouches labeled Pouch B

Keep out of reach of children.

Phenylketonurics: contains phenylalanine – a maximum of 131 mg per treatment.

**Moviprep**  
(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride, Sodium Ascorbate and Ascorbic Acid for Oral Solution)  
100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g

**Rx only**

NDC 65649-201-75

**Distributed by:**  
Salix Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

See full Prescribing Information and patient information provided prior to use.  
Keep out of reach of children.

**Moviprep**  
(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride, Sodium Ascorbate and Ascorbic Acid for Oral Solution)  
100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g

205 mm

13 mm

145 mm

40 mm

145 mm

39 mm

NDC 65649-201-75

Rx only

# MoviPrep®

(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride,  
Sodium Ascorbate and Ascorbic Acid for Oral Solution)

100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g

**On mixing in 1 liter of water, one Pouch A  
and one Pouch B provide:**

PEG-3350	29.6 mmol/L
Sodium	181.6 mmol/L
Chloride	59.8 mmol/L
Sulfate	52.8 mmol/L
Potassium	14.2 mmol/L
Ascorbate	29.8 mmol/L

## Container for mixing of MoviPrep®

### Directions for use:

Dissolve the contents of one Pouch **A** and one Pouch **B** in 1 liter of water and drink one 8 oz. glass of the solution (approximately 240 mL) every 15 minutes until gone, followed by 16 oz. of clear fluid (approximately 500 mL). Repeat as directed.

**Important:** See package insert for full prescribing information. When mixed, store in an upright position in the refrigerator. Use within 24 hours.

**Keep out of reach  
of children.**

Phenylketonurics: Contains  
Phenylalanine – a maximum  
of 131 mg per treatment.

**Salix** 

Distributed by:  
Salix Pharmaceuticals, a division of  
Bausch Health US, LLC  
BrIDGEWATER, NJ 08807 USA

MoviPrep is a trademark of Veltrop AG used  
under license  
© 2019 Bausch Health Companies Inc. or its affiliates

965001

Salix Pharmaceuticals  
Printed Side  
T122618A


No Coating Area   
No Print Area 

IC Bars

70014286

Area for  
Serialization, Lot/Exp

**MoviPrep**  
(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride,  
Sodium Ascorbate and Ascorbic Acid for Oral Solution)  
100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g



Rx only

NDC 65649-201-75

Rx only

NDC 65649-201-75

**MoviPrep**

(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride,  
Sodium Ascorbate and Ascorbic Acid for Oral Solution)

100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g

Phenylephrine:  
contains phenylephrine – a maximum of 131 mg per treatment.

PHARMACIST: Dispense the enclosed  
Medication Guide to each patient.

Keep out of reach of children.

This carton contains: One container to mix MoviPrep®

2 Pouches Labeled Pouch A:	2 Pouches Labeled Pouch B:	On mixing in 1 liter of water, one Pouch A and one Pouch B provide:
PEG-3350 100 g	Sodium ascorbate 5.9 g	PEG-3350 29.6 mmol/L
Sodium sulfate 7.5 g	Ascorbic acid 4.7 g	Sodium 181.6 mmol/L
Sodium chloride 2.691 g		Chloride 59.8 mmol/L
Potassium chloride 1.015 g		Sulfate 52.8 mmol/L
also contains aspartame		Potassium 14.2 mmol/L
		Ascorbate 29.8 mmol/L

Salix  
PHARMACEUTICALS

INSTRUCTIONS - See inside for full Prescribing Information and Instructions For Use.

**MoviPrep (moo-vee-prép)**  
(PEG-3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate and ascorbic acid for oral solution, 100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g)

**Important Information:**

- Take MoviPrep as prescribed by your healthcare provider.
- The contents of each pouch of MoviPrep must be mixed with water before taking.
- Do not eat any solid food while taking MoviPrep until after your colonoscopy.**
- Drink only clear liquids:
  - while taking MoviPrep
  - after taking MoviPrep and until 2 hours before your colonoscopy or as prescribed by your healthcare provider

**You may eat or drink any of the following clear liquids:**

- water
- clear fruit juices without pulp including apple, white grape, or white cranberry
- strained limeade or lemonade
- coffee or tea (do not use any dairy or non-dairy creamer.)
- clear broth
- clear soda
- gelatin (without added fruit or topping)
- popicles (without pieces of fruit or fruit pulp)

Do not eat or drink anything colored red or purple.

Do not eat or drink anything 2 hours before your colonoscopy or as prescribed by your healthcare provider.

Store at room temperature between 20° to 25°C (68° to 77°F).

Manufactured by:  
Salix Pharmaceuticals, a division of  
Bauch Health US, LLC  
Bridgewater, NJ 08807 USA  
Product of United Kingdom  
US Patent Numbers: 7,189,381 and 7,868,014  
MoviPrep is a trademark of Veltrop AG used under license.  
© 2019 Bauch Health Companies Inc. or its affiliates  
9656201 70014286



INSTRUCTIONS (continued from back panel)

MoviPrep solution is taken by mouth in one of the following ways:

**Two-Day Split-Dosing Instructions:**

**Dose 1 – Take this dose the evening before your colonoscopy (10 to 12 hours before Dose 2):**

- Empty the contents of 1 Pouch A and 1 Pouch B into the container that comes with MoviPrep.
- Add lukewarm water to the Fill Line on the container.
- Mix to completely dissolve the contents of Pouch A and B into the lukewarm water.
- Drink one 8 oz. (ounce) glass (240 mL) every 15 minutes. Be sure to drink all of the solution. This should take about 1 hour.
- Fill the container with 16 oz. (two 8 oz. glasses) of clear liquid and drink all of this liquid before you go to bed.

**Dose 2 – Take this dose the next morning on the day of your colonoscopy (start at least 3 ½ hours before your colonoscopy):**

- Repeat steps 1 through 4 from Dose 1 of the Two-Day Split-Dosing instructions.
- Fill the container with 16 oz. (two 8 oz. glasses) of clear liquid and drink all of this liquid at least 2 hours before your colonoscopy.
- Drink only clear liquids up to 2 hours before your colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after your colonoscopy.

Or

**One-Day Evening Only Instructions:**

**Dose 1 – Take this dose at least 3 ½ hours before bedtime the evening before your colonoscopy:**

- Empty the contents of 1 Pouch A and 1 Pouch B into the container that comes with MoviPrep.
- Add lukewarm water to the Fill Line on the container.
- Mix to completely dissolve the contents of Pouch A and B into the lukewarm water.
- Drink one 8 oz. (ounce) glass (240 mL) every 15 minutes. Be sure to drink all of the solution. This should take about 1 hour.

**Dose 2 – Take Dose 2 about 1 ½ hours after starting Dose 1 the evening before your colonoscopy:**

- Repeat steps 1 through 4 from Dose 1 of the One-Day Evening Only instructions.
- After you complete steps 1 through 4, fill the container again to the Fill Line with clear liquid and drink all of this liquid before you go to bed.
- Drink only clear liquids up to 2 hours before your colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after your colonoscopy.

MoviPrep solution may be refrigerated before drinking. Keep the mixed solution in an upright position and store in the refrigerator. MoviPrep should be taken within 24 hours after it is mixed in water. Do not add other ingredients to the MoviPrep solution.

8 5/16

5 1/16

5/8


Glue

Glue

Glue

Glue

Glue



360 mm

180 mm

180 mm

80 mm

130 mm

**MoviPrep<sup>®</sup>** Rx only **Pouch A**

(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride, Sodium Ascorbate and Ascorbic Acid for Oral Solution)

100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g

**Instructions for use:**  
Dissolve the contents of one Pouch A and one Pouch B in 1 liter of water and drink one 8 oz. glass of the solution (approximately 240 mL) every 15 minutes until gone.

**This one Pouch A must be mixed with one Pouch B.**

Distributed by:  
Salix Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA  
9656401 42130004

**Important:** See package insert for full prescribing information.  
Store at room temperature between 20° to 25°C (68° to 77°F).

**Keep out of reach of children.**  
Phenylketonurics: Contains Phenylalanine – a maximum of 131 mg per treatment.

MoviPrep is a trademark of Vellnor AG used under license.  
© 2019 Bausch Health Companies Inc. or its affiliates

NDC 65649-201-75

**MoviPrep<sup>®</sup>**

(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride, Sodium Ascorbate and Ascorbic Acid for Oral Solution)

100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g

Pouch A	
Each Pouch A contains:	
PEG-3350	100 g
Sodium sulfate	7.5 g
Sodium chloride	2.691 g
Potassium chloride	1.015 g
also contains aspartame	

This pouch contains 111.9 g of powder for oral solution.

**Rx only**

See enclosed Instructions for Use.

**Salix**  
PHARMACEUTICALS

LOT  
EXP  
Pre-print

1041

10 mm

14 mm

18 mm

**NOTE:**

If only two lines of print is required use line position 18 and 14mm. Character size is 2mm high.

Unwind Direction



**NOTE:**  
 If only two lines of print is required use line position 18 & 14mm. Character size is 2mm high.



David  
Lewis

Digitally signed by David Lewis  
Date: 11/05/2019 07:53:42AM  
GUID: 508da72000029f287fa31e664741b577