



2 Count  
Packets

# Advil

**SINUS  
CONGESTION & PAIN**

Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)  
Phenylephrine HCl 10 mg Nasal Decongestant



# Advil

**SINUS  
CONGESTION & PAIN**

2  
Packets of 1  
Coated Tablet

**NON-DROWSY**

# Advil

**SINUS  
CONGESTION & PAIN**

Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)  
Phenylephrine HCl 10 mg Nasal Decongestant

- ✓ Nasal Congestion
- ✓ Sinus Pressure
- ✓ Nasal Swelling
- ✓ Headache
- ✓ Fever
- ✓ Body Aches



**ONE  
TABLET  
DOSE!**

2  
Packets of 1  
Coated Tablet



Do Not Use if pouch  
is broken or torn.

WHITE HOLDOUT BELOW  
THIS LINE FOR LOT CODE  
& EXPIRATION DATE  
(NO VARNISH)

# Advil

**SINUS  
CONGESTION & PAIN**

For most recent product information,  
visit [www.advil.com](http://www.advil.com)

Questions or comments?  
Call weekdays from 9 AM to 5 PM EST  
toll free at 1-800-88-ADVIL

Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc.

PAA135323.FDA01

090177e192d30715FinalFinal On: 12-Feb-2020 14:48 (GMT)

**PUT  
HERE**

PA4140916 FDA01

<b>READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION</b>	
<b>Drug Facts</b>	<b>Active ingredients (in each tablet)</b>
Pseudoephedrine HCl 10 mg	Oxymetazoline HCl 0.05 mg
<b>Uses</b>	
Relieves these symptoms associated with the common cold or flu:	
• sneezing • runny nose • nasal congestion • watery, itchy eyes • minor body aches and pains • reduces swelling of the nasal passages • temporarily reduces nasal breathing through the nose	

### PANEL 1

<b>Drug Facts (continued)</b>	<b>Warnings</b> Allergic reaction: Sulfonamide antibiotics may cause a severe allergic reaction, especially in children. Symptoms may include: • hives • rash or skin irritation • difficulty breathing • swelling of the face, lips, tongue, or throat • severe dizziness • fainting
<b>Other information</b>	This product contains an NSAID, which may cause severe stomach bleeding in people taking aspirin, ibuprofen, or other NSAIDs. Long-term use of this product may increase the risk of stomach bleeding.
<b>Do not use</b>	• if you are taking aspirin, ibuprofen, or other NSAIDs • if you have had stomach ulcers or bleeding problems • if you are pregnant, trying to get pregnant, or breastfeeding

### PANEL 2

<b>Drug Facts (continued)</b>	<b>Do not use</b>
• if you are taking aspirin, ibuprofen, or other NSAIDs • if you have had stomach ulcers or bleeding problems • if you are pregnant, trying to get pregnant, or breastfeeding	• if you are taking aspirin, ibuprofen, or other NSAIDs • if you have had stomach ulcers or bleeding problems • if you are pregnant, trying to get pregnant, or breastfeeding

### PANEL 3

<b>Drug Facts (continued)</b>	<b>Warnings</b> Allergic reaction: Sulfonamide antibiotics may cause a severe allergic reaction, especially in children. Symptoms may include: • hives • rash or skin irritation • difficulty breathing • swelling of the face, lips, tongue, or throat • severe dizziness • fainting
<b>Other information</b>	This product contains an NSAID, which may cause severe stomach bleeding in people taking aspirin, ibuprofen, or other NSAIDs. Long-term use of this product may increase the risk of stomach bleeding.
<b>Do not use</b>	• if you are taking aspirin, ibuprofen, or other NSAIDs • if you have had stomach ulcers or bleeding problems • if you are pregnant, trying to get pregnant, or breastfeeding

### PANEL 4

<b>Drug Facts (continued)</b>	<b>Warnings</b> Allergic reaction: Sulfonamide antibiotics may cause a severe allergic reaction, especially in children. Symptoms may include: • hives • rash or skin irritation • difficulty breathing • swelling of the face, lips, tongue, or throat • severe dizziness • fainting
<b>Other information</b>	This product contains an NSAID, which may cause severe stomach bleeding in people taking aspirin, ibuprofen, or other NSAIDs. Long-term use of this product may increase the risk of stomach bleeding.
<b>Do not use</b>	• if you are taking aspirin, ibuprofen, or other NSAIDs • if you have had stomach ulcers or bleeding problems • if you are pregnant, trying to get pregnant, or breastfeeding

### BASE PANEL

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	N/A
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

PA4140916 FDA01

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
07/30/2020 04:08:00 PM