


1.14.1.1 Draft Carton and Container Labels

PROPOSED SINGLE DOSE UNIT FOIL POUCH



1.14.1.1 Draft Carton and Container Labels

PROPOSED SINGLE DOSE UNIT CARTON

LF60015802-01
All dimensions in mm
M. Olczak 21.02.2019

9890907
AB63720

Drug Facts (continued)
boric acid, calcium chloride dihydrate, glycerin, potassium chloride, sodium borate decahydrate, sodium chloride, water for injection. Hydrochloric acid and/or sodium hydroxide may be used to adjust pH.

Drug Facts (continued)
Questions or comments? (Call: 1-800-563-5340)
LUMIFY is a trademark of Bausch & Lomb Incorporated or its affiliates.
© 2019 Bausch & Lomb Incorporated or its affiliates Product of Spain

BAUSCH+LOMB
LUMIFY®

BRIMONIDINE TARTRATE
OPHTHALMIC SOLUTION 0.025%
REDNESS RELIEVER EYE DROPS

NDC 24208-537-01

- Works in 1 minute
- Lasts up to 8 hours

Sterile

1 SINGLE-USE VIAL
SAMPLE - NOT FOR SALE
0.013 FL OZ (0.4 mL)

LOT
EXP
Varnish free
35 x 14

LUMIFY® eye drops, from the eye care experts at Bausch + Lomb. Just one drop works within a minute and lasts up to 8 hours to reduce redness.

DO NOT USE IF POUCH IS TORN OR OPEN

Distributed by: Bausch + Lomb, a division of Bausch Health US, LLC, Bridgewater, NJ 08807 USA
US Patent Number: 8,293,742

Drug Facts	
Active ingredient Brimonidine tartrate (0.025%)	Purpose Redness reliever
Use ■ relieves redness of the eye due to minor eye irritations	
Warnings For external use only Do not use ■ if solution changes color or becomes cloudy ■ if single-use vial is not intact Stop use and ask a doctor if ■ you experience	

Drug Facts (continued)
eye pain, changes in vision, continued redness or irritation of the eye ■ condition worsens or persists for more than 3 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)
■ remove contact lenses before use ■ wait at least 10 minutes before re-inserting contact lenses after use
■ if using other ophthalmic products while using this product, wait at least 5 minutes between each product
■ to avoid contamination, do not touch tip of container to any surface ■ children under 5 years of age: consult a doctor

Directions ■ adults and children 5 years of age and over: ■ instill 1 drop in the affected eye(s)
■ discard vial immediately after use; do not re-use
■ do not use more than 4 times daily (every 6–8 hours)

Other information ■ store at 15°-25°C (59°-77°F)

Inactive ingredients benzalkonium chloride

Varnish free

1.14.1.1 Draft Carton and Container Labels


PROPOSED SINGLE DOSE UNIT VIAL

(image scaled to 200%)



1.14.1.1 Draft Carton and Container Labels

PROPOSED TRADE SINGLE DOSE UNIT FOIL POUCH



1.14.1.1 Draft Carton and Container Labels

PROPOSED TRADE SINGLE DOSE UNIT CARTON

LF60015802-01
All dimensions in mm
M. Olczak 21.02.2019

Drug Facts (continued)
boric acid, calcium chloride dihydrate, glycerin, potassium chloride, sodium borate decahydrate, sodium chloride, water for injection. Hydrochloric acid and/or sodium hydroxide may be used to adjust pH.

Drug Facts (continued)
Questions or comments? (Call: 1-800-553-6340)
LUMIFY is a trademark of Bausch & Lomb Incorporated or its affiliates.
© 2019 Bausch & Lomb Incorporated or its affiliates Product of Spain

BAUSCH + LOMB
LUMIFY[®]
BRIMONIDINE TARTRATE
OPHTHALMIC SOLUTION 0.025%
REDNESS RELIEVER EYE DROPS

Works in 1 minute
Lasts up to 8 hours

1 SINGLE-USE VIAL
Sterile 0.013 FL OZ (0.4 mL)

NDC 24208-537-10

LUMIFY[®] eye drops, from the eye care experts at Bausch + Lomb. Just one drop works within a minute and lasts up to 8 hours to reduce redness.

DO NOT USE IF POUCH IS TORN OR OPEN

Distributed by: Bausch + Lomb, a division of Bausch Health US, LLC, Bridgewater, NJ 08807 USA
US Patent Number 8,293,742

Drug Facts

Active ingredient	Purpose
Brimonidine tartrate (0.025%)	Redness reliever

Use ■ relieves redness of the eye due to minor eye irritations

Warnings
For external use only

Do not use ■ if solution changes color or becomes cloudy ■ if single-use vial is not intact

Stop use and ask a doctor if ■ you experience

Drug Facts (continued)

eye pain, changes in vision, continued redness or irritation of the eye ■ condition worsens or persists for more than 3 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ adults and children 5 years of age and over: ■ instill 1 drop in the affected eye(s) ■ discard vial immediately after use; do not re-use ■ do not use more than 4 times daily (every 6-8 hours)

Drug Facts (continued)

■ remove contact lenses before use ■ wait at least 10 minutes before re-inserting contact lenses after use ■ if using other ophthalmic products while using this product, wait at least 5 minutes between each product ■ to avoid contamination, do not touch tip of container to any surface ■ children under 5 years of age: consult a doctor

Other information ■ store at 15°-25°C (59°-77°F)

Inactive ingredients benzalkonium chloride,

LOT
EXP

Varnish free
35 x 14

Varnish free

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
03/05/2020 04:47:52 PM