
\*Each tablet contains 20 mg of cabozantinib (equivalent to 25 mg of cabozantinib (S)-malate).

Take once each day on an empty stomach. Cabometyx should not be taken with food. Take Cabometyx at least 1 hour before or at least 2 hours after eating. Swallow Cabometyx tablets whole. Do not crush Cabometyx tablets.

Store Cabometyx at 20°C to 25°C (68°F to 77°F); excursions are permitted from 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

Keep out of the reach of children.



\*Each tablet contains 40 mg of cabozantinib (equivalent to 51 mg of cabozantinib (S)-malate).

Take once each day on an empty stomach. Cabometyx should not be taken with food. Take Cabometyx at least 1 hour before or at least 2 hours after eating. Swallow Cabometyx tablets whole. Do not crush Cabometyx tablets.

Store Cabometyx at 20°C to 25°C (68°F to 77°F); excursions are permitted from 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

Keep out of the reach of children.





\*Each tablet contains 60 mg of cabozantinib (equivalent to 76 mg of cabozantinib (S)-malate).

Take once each day on an empty stomach. Cabometyx should not be taken with food. Take Cabometyx at least 1 hour before or at least 2 hours after eating. Swallow Cabometyx tablets whole. Do not crush Cabometyx tablets.

Store Cabometyx at 20°C to 25°C (68°F to 77°F); excursions are permitted from 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

Keep out of the reach of children.





Digitally signed by Ramesh Raghavachari

Date: 11/05/2020 02:57:00PM

GUID: 502d0913000029f375128b0de8c50020

,