

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt	Horizontal Scale:	75%
Headers:	4.50 pt	Leading:	4.50 pt	Barlines:	NA pt		
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt	Average Tracking:	-10

00000000

LOT:

EXP:

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*

Pain reliever/fever reducer

nonsteroidal anti-inflammatory drug

Uses

■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache
■ backache ■ the common cold ■ menstrual cramps
■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ ulcers. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you

NDC 50580-486-05

Motrin[®] IB

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

100 Coated Caplets ^{Caplet-Shaped Tablets}

than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).
Directions ■ **do not take more than directed** ■ **the smallest effective dose should be used.** Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist. ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used. ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor
Other information ■ store between 20-25°C (68-77°F) ■ **do not use if tell inner seal imprinted "Sealed For Your Safety" is broken or missing**
Inactive ingredients carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide
Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect) (collect)
Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©J&JCI 20XX **www.motrin.com**

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt	Horizontal Scale:	75%
Headers:	4.50 pt	Leading:	4.50 pt	Barlines:	NA pt		
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt	Average Tracking:	-10

Reference ID: 4889366

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: NA pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%	
Headers: 4.50 pt	Leading: 4.50 pt	Barlines: NA pt		
Subheaders: 4.50 pt	Drug Facts (continued): NA pt	Hairlines: NA pt	Average Tracking: -16	

00000000



LOT:
EXP:

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer

Purposes
*nonsteroidal anti-inflammatory drug

Uses
■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis
■ toothache ■ backache ■ the common cold ■ menstrual cramps
■ temporarily reduces fever

Warnings:
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product** ■ take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better. ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area. ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions** ■ **do not take more than directed** ■ **the smallest effective dose should be used.** **Adults and children 12 years and older:** ■ take 1 caplet every 4 to 6 hours while symptoms persist. ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used. ■ **do not exceed 6 caplets in 24 hours, unless directed by a doctor.** **Children under 12 years:** ■ ask a doctor. **Other information** ■ store between 20-25°C (68-77°F) ■ **do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing.** **Inactive ingredients** camauaba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©J&JCI 20XX www.motrin.com

VALUE SIZE!

NDC 50580-486-06

Motrin[®] IB

Ibuprofen Tablets USP 200 mg

Pain Reliever / Fever Reducer (NSAID)


actual size

225 Coated Caplets*

Capsule-Shaped Tablets

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt	Horizontal Scale:	83%
Headers:	4.50 pt	Leading:	5.00 pt	Barlines:	NA pt		
Subheaders:	4.50 pt	Drug Facts (continued):	NA pt	Hairlines:	NA pt	Average Tracking:	-10

00000000

EXP:

LOT:

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)

nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - toothache
 - backache
- minor pain of arthritis
- the common cold
- menstrual cramps

Ibuprofen reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include: ■ hives ■ stomach aches ■ asthma (wheezing) ■ shock ■ skin reddening

■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach**

bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug ■ take other drug containing prescription or

nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have a 3 or more alcoholic drinks every

day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke**

warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal.

The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you

have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or

after heart surgery. **Ask a doctor before use if** ■ you have problems or serious side effects from taking

pain relievers or fever reducers ■ have problems using aspirin or you ■ you have a history of

stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis,

kidney disease, asthma, or had a stroke ■ you are taking a diuretic. **Ask a doctor or pharmacist**

before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this

Purposes

Pain reliever/fever reducer

NDC 50580-486-07

Motrin[®] IB

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

Motrin[®] IB

300

Coated Caplets[®]

*Capsule-Shaped Tablets

benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. **When**

using this product ■ take with food or milk ■ stomach upset occurs. **Stop use and ask a doctor if**

■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have

bloody or black stools ■ have stomach pain that does not get better. ■ If you have symptoms of heart

problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body

■ slurred speech ■ leg swelling. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or

lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms

appear. **If pregnant or breast-feeding,** ask a health professional before use. It is especially

important not to use ibuprofen 10 weeks or later in pregnancy because it can be so by

a doctor because it may cause problems in the unborn child or complications during delivery. **Keep**

out of reach of children. In case of overdose, get medical help or contact a Poison Control Center

right away. (1-800-222-1222)

Directions ■ **do not take more than directed** ■ **the smallest effective dose should be**

used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while

symptoms persist ■ **do not take more than 12 caplets in 24 hours** ■ **do not take more than 3**

caplets per day ■ **do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years:** ■ **ask a doctor**

Other information: ■ store between 20-25°C (68-77°F) ■ **do not use if foil inner seal imprinted**

"Sealed For Your Safety" is broken or missing

Inactive ingredients: carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6, aluminum lake,

hydrophobic, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol,

pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-877-895-3665 (toll-free) or 1-215-275-8755 (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**

McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA

6/2011 20XX

www.motrin.com

Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: NA pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 90%	
Headers: 4.50 pt	Leading: 5.00 pt	Barlines: NA pt		
Subheaders: 4.50 pt	Drug Facts (continued): NA pt	Hairlines: NA pt	Average Tracking: -10	

00000000



LOT:

EXP:

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis
■ toothache ■ backache ■ the common cold ■ menstrual cramps
■ temporarily reduces fever

Warnings
Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product. ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product**

For Hospital and Government Use Only



NDC 50580-486-02

Motrin[®] IB
Ibuprofen Tablets usp, 200 mg
Pain Reliever / Fever Reducer (NSAID)

500
Coated Caplets*


*Capsule-Shaped Tablets

■ take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better. ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)


Directions
■ **do not take more than directed** ■ **the smallest effective dose should be used**
Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. **Children under 12 years:** ■ ask a doctor

Other information
■ store between 20-25°C (68-77°F) ■ **do not use if foil inner seal imprinted "Sealed For Your Safety" is broken or missing**

Inactive ingredients
carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6 aluminum lake, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
©J&J 20XX **www.motrin.com**



Labeling Format Information					
Font: HelveticaCond Bold, Cond Black, Cond Black Italic					
Drug Facts:	NA pt	Body Text:	4.50 pt	Bullets:	3.50 pt
Warnings:	4.50 pt	Warnings:	4.50 pt	Warnings:	NA pt
Subheaders:	4.50 pt	Drug Facts continued:	NA pt	Warnings:	NA pt
Average Tracking: -25					




Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 65%	
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt		
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0	



Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 65%	
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt		
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0	



Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 65%	
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt		
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0	





Labeling Format Information: Fonts: Helvetica®-Medium, Bold, Bold Italic					
Drug Facts:	9.00 pt	Body Text:	6.00 pt	Bullets:	5.00 pt
Warnings:	8.00 pt	Leading:	6.50 pt	Barlines:	2.50 pt
Subheaders:	8.00 pt	Drug Facts (continued):	8.00 pt	Headlines:	12.00 pt
				Average Tracking:	-15



Labeling Format Information: Fonts: Helvetica®-Medium, Bold, Bold Italic					
Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 85%		
Headings: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt			
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Barlines: 0.50 pt	Average Tracking: 0		



Labeling Format Information: Fonts: HelveticaDF-Cond Medium, Cond Bold, Cond Bold Italic				
Drug Facts: 8.50 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 95%	
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt		
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Tracking: 0	

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
11/16/2021 03:01:48 PM