XXXXXX

LOT/EXP PRINTED OR DEBOSSED ONLINE FORMAT BELOW:



12789-MM-YY

Dist. by Medtech Products Inc. Tarrytown, NY 10591 A Prestige Consumer Healthcare company ©2020 Trade dress is owned by Medtech Products Inc. All rights reserved. Made in Spain. TGUS013101

Drug Facts

Active ingredient (in each tablet) Cimetidine 200 mg Purpose Anid reducer

Uses relieves heartburn associated with acid indigestion and sour stomach prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers.

Do not use I if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. I with other acid reducers

Ask a doctor before use if you have # frequent chest pain # frequent wheezing, particularly with heartburn # unexplained weight loss # nausea or vomiting # stomach pain # had heartburn over 3 months. This may be a sign of a more serious condition. # heartburn with lightheadedness, sweating or dizziness * chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness # kidney disease # liver disease.

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if you need to take this product for more than 14 days stomach pain continues your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions Adults and children 12 years and over: ** to relieve symptoms, swallow 1 tablet with a glass of water ** to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn ** do not take more than 2 tablets in 24 hours. Children under 12 years: ask a dootor.

Other information store at 15°-30°C (59°-86°F)

Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions?

Call toll-free 1-800-482-4394 weekdays.

Keep outer carton for full information.

NEW VALUE SIZE

TAKE ANYTIME YOU NEED IT: before meal · during meal · after meal

Tagamet T≈200

> Cimetidine Tablets 200 mg Acid Reducer

JUST ONE TABLET
RELIEVES and PREVENTS

Heartburn and Acid Indigestion



TABLETS (70 DOSES)

TIPS FOR MANAGING HEARTBURN:

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- · If you smoke, quit smoking
- · Raise the head of your bed
- Wear loose fitting clothing around your stomach

Tamper Evident: Do Not Use if seal under cap is broken or missing.

NO COATING







Drug Facts

Active ingredient Purpose (in each tablet) Cimetidine 200 mg

Acid reducer

Uses

relieves heartburn associated with acid indigestion and sour stomach prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert Do not use if you are allergic to cimetidine or other acid reducers.

Do not use

if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers

Tamper Evident: Do Not Use if eal under cap is broken or missing. ©2020 Dist, by Medtech Products Inc., Terrytown, NY 10591.

A Prestige Consumer Healthcare company. Made in Spain. TGUS013201

BACK

Drug Facts (continued)

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders: or lightheadedness

■ kidney disease

■ liver disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if you need to take this product for more than 14 days ■ stomach pain continues ■ your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

BASE

Drug Facts (continued) Directions

in 24 hours.

- Adults and children 12 years and over: ■ to relieve symptoms, swallow
- 1 tablet with a glass of water ■ to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn do not take more than 2 tablets

Other information store at 15°-30°C (59°-86°F)

Children under 12 years: ask a doctor

Inactive ingredients cellulose, corn starch, hypromellose,

magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions? 1-800-482-4394

This is a representation of an electronic record that was signed
electronically. Following this are manifestations of any and all
electronic signatures for this electronic record.

/s/

NUSHIN F TODD 03/15/2021 01:37:30 PM