



### PLANNING YOUR SUCCESS.

- The key to accomplishing anything important is commitment. When it comes to quitting smoking, that is especially true. Nicorette Lozenges can help if you really want to quit. Nicorette Lozenges help reduce withdrawal symptoms including nicotine craving associated with quitting smoking.
- Your chances of staying off cigarettes are much better if you start with at least 9 Nicorette Lozenges daily. For best results, use the lozenges on a regular schedule (as outlined in this User's Guide).
- Start using Nicorette Lozenges on your quit date.

- 4) This User's Guide outlines a 12 week plan for Nicorette Lozenges. Even though you may feel confident about your non-smoking status after a few weeks, it's important to stick with the plan to help you remain smoke-free. Even a single cigarette can put you right back to square one.
- Nicorette Lozenges work best when used together with a support plan.
   See insert between pages 12 and 15 for instructions on enrollment in the MyQuit Individualized Stop Smoking Program.
- 6) After the first six weeks, start using fewer Nicorette Lozenges, as directed in the instructions, gradually reducing your use over the next six weeks. If you feel the need to use

the lozenges for a longer period to keep from smoking, talk to your health care provider.

 If you have questions about using Nicorette Lozenges, call toll-free 1-888-569-1743, or talk to your pharmacist or family doctor.

### YES! YOU WANT TO QUIT.

Wonderful. You've made the most important decision of all, to stop smoking. And by choosing **Nicorette** Lozenges to help you, you're starting on the right path. Now remember, using **Nicorette** Lozenges doesn't just mean taking a **Nicorette** Lozenge. It means setting and following a program like the one we suggest in this User's Guide.

Your own success depends on your effort, your level of addiction to tobacco, and your commitment to

### following your program.

### LET'S FACE IT.

Quitting smoking isn't easy! You or someone you know may have tried unsuccessfully. That's okay. It's hard to stop smoking the first time you try. The important part is to learn from your previous attempts, consider what went wrong and keep trying to quit until you succeed. Look to this User's Guide for support as you undergo this terrific task. The guide includes important information on how to use **Nicorette** Lozenges and also gives you tips to help you stop smoking. Refer back to it often for advice, answers, and encouragement to help you stay on track.

## **GET MOTIVATED. STAY MOTIVATED.** Everyone has a reason for quitting — whether you're concerned about your

health, your appearance, family or peer pressure, or the effect of secondhand smoke on your loved ones - all of the above, or something else entirely. Whatever your reasons, write them down There's a WALLET CARD inside the back cover of this User's Guide. Write your reasons on the card and carry it with you. When you have an urge to smoke or experience a difficult moment it can help you focus on your reasons for quitting. Lots of people quit with a co-worker, spouse or friend and use them as a quitting buddy. You can help each other out by providing extra encouragement in tough moments.

There may be support groups in your area for people trying to quit. Call your local chapter of the American Lung Association, American Cancer Society or American Heart Association for further information. Toll-free phone numbers are printed on the WALLET CARD on the back cover of this User's Guide.

# UNDERSTANDING THE DOUBLE-EDGED SWORD.

Smoking has two addictive components, a physical and a mental need for the nicotine in tobacco. You need to conquer both to succeed. **Nicorette** Lozenges can ease your physical nicotine addiction. But your readiness and resolve are necessary to help overcome the mental side of your cigarette dependence. So once you're ready, it's time to begin. But first, read and consider the following important warnings. IMPORTANT WARNINGS.

This product is only for those who want to stop smoking.

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

### Do not use

 if you are allergic to soya (mint flavor only)

### Ask a doctor before use if you have

- · a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.

- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

# Ask a doctor or pharmacist before use if you are

- · using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

### Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat

 you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

### YOU'RE READY TO START.

Okay, you're ready. To become a non-smoker, start today. Now before you do anything else, you have a bit of planning to do. Read this User's Guide all the way through. You want to make sure you bought the right dose to start. If you typically smoke your first cigarette within 30 minutes of waking up, use the 4 mg Nicorette Lozenges. If you smoke your first cigarette more than 30 minutes after waking up, use the 2 mg Nicorette Lozenges. Next, plan your quitting schedule. Get a calendar to follow your progress and mark the following four important dates (see the reminders on page 26 of this booklet).

### THE PROGRAM.

STEP 1. (Weeks 1-6) Starting on your quit date it's best to use at least 9 Nicorette Lozenges each day, one every 1-2 hours.

First choose the day you plan to quit (make it soon). Place the Step 1 reminder on this date. That's the day you will start using **Nicorette** Lozenges to calm your cravings for nicotine and help you stay smoke-free. Prior to the quit date, get rid of all your cigarettes to remove temptations and make it more difficult to start smoking again.

Use a Nicorette Lozenge every 1 to 2 hours and at least 9 lozenges each day for the first 6 weeks to help prevent unexpected cravings and improve your chances of quitting. These aren't ordinary lozenges. Place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 - 30 minutes). Minimize swallowing. Do not chew or swallow the lozenge. You may feel a warm or tingling sensation. Occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 to 30 minutes). Remember to read the instructions on page 11 before you take your first Nicorette Lozenge.

#### STEP 2. (The next three weeks, that is weeks 7-9). At the beginning of week 7 start using fewer Nicorette Lozenges, one every 2-4 hours.

After six weeks, you should wait a little longer between lozenges, one lozenge every two to four hours. This will help you gradually use fewer Nicorette Lozenges. Put the Step 2 reminder on the first day of week 7 to help remind you when to start reducing the number of Nicorette Lozenges you take.

STEP 3. (The last three weeks, that is weeks 10-12). At the beginning of week 10, reduce Nicorette Lozenge use even further, one every 4-8 hours. At the beginning of week 10 further decrease the number of Nicorette Lozenges you use each day to reduce the amount of nicotine you get. You should do this by using one lozenge every 4 to 8 hours. Put the Step 3 reminder on the first day of week 10 so you know when you should be starting this last step to becoming smoke and nicotine-free.

END. At the end of week 12 you'll complete Nicorette Lozenge therapy. Put the "EX-SMOKER" reminder on your calendar on the date 12 weeks after the day you stopped smoking and started using Nicorette Lozenges.

#### **BE PREPARED.**

Since smoking is an addiction, it is hard to quit. Even after you stop, there will be times when you WANT a cigarette, sometimes strongly. (See also section on "CHALLENGES TO WATCH FOR"). The best defense is to be prepared. Plan now for handling tough times so

you don't give in. Think about situations where you usually get an urge to smoke or where you might experience strong urges to smoke (for example, spending time with smokers or drinking alcohol — try to avoid these situations if those things tempt you to smoke).

Change your habits. For example, take your coffee break somewhere else. Take a walk. In other words, break the association between your usual habits and cigarettes.

If you do encounter a situation where you feel a strong urge to smoke, fight it! Take a break from the situation; keep yourself busy or distracted with other activities. Remind yourself why you want to quit, and above all, remind yourself that having "just one" really will hurt your goal of quitting! Assemble a "survival package" — items that can keep you distracted in case you get an urge to smoke. For example, cinnamon gum or hard candy, relaxing music, things to keep your hands busy like a smooth stone or paper clips, etc.

Track your quit progress. Use a journal or the MyQuit app on your mobile phone to note if and when you get an urge to smoke. Note how many pieces of **Nicorette** Lozenges you use each day. If you slip and have a cigarette, don't give up. Stop smoking again and get back on your program with **Nicorette** Lozenges.

Establish your support network. Keep friends' and family members' phone numbers ready to get the moral support you need. Before quitting, ask friends and family to support and encourage you. Think of specific ways they can help.

Reward yourself. Set aside little gifts to encourage yourself which you can earn by overcoming difficult hurdles.

HOW Nicorette LOZENGES WORK. Nicotne Replacement Therapy. They deliver nicotine to your body, temporarily relieving craving and nicotine withdrawal symptoms when you quit smoking. But unlike cigarettes, Nicorette Lozenges deliver a lower, steady level of nicotine to your blood. When used as directed, Nicorette Lozenges help you regulate, control, and gradually reduce your body's craving for nicotine.

The good news is that **Nicorette** Lozenges contain no tar or carbon monoxide, and therefore don't present the same medical risks as cigarettes.

However, the lozenges still deliver nicotine, the addictive ingredient in cigarettes. And for some people the nicotine in **Nicorette** Lozenges can occasionally cause mouth or throat irritation, headaches, nausea, hiccups, upset stomach or dizziness.

#### USING Nicorette LOZENGES PROPERLY.

Remember, Nicorette Lozenges aren't like ordinary lozenges such as cough drops. This lozenge is designed to deliver nicotine into your system through the lining of your mouth, not in your stomach like most other medicines. It is important to minimize swallowing the dissolved medicine in these lozenges so that it can be properly absorbed in your mouth.

Do not use more than one lozenge at a time, or many lozenges one after another since this can cause hiccups, heartburn, nausea or other side effects.

Read all the following instructions before using **Nicorette** Lozenges. Refer to them often to make sure you're using **Nicorette** Lozenges correctly. IMPORTANT: Don't worry or give up if you do not like the taste of the lozenge at first. Nicorette Lozenges are a medication, not a candy. Most people get used to the taste after a day or two. Remember, staying with the plan will help you quit. Begin using Nicorette Lozenges on your quit date.

- Remove the Nicorette Lozenge from the immediate container. Place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 - 30 minutes). Minimize swallowing. Do not chew or swallow the lozenge. You may feel a warm or tingling sensation.
- Occasionally move the lozenge from one side of your mouth to the other side until completely dissolved (about 20 - 30 minutes).

To reduce cravings or urges to smoke and other withdrawal symptoms, use **Nicorette** Lozenges according to the following dosage schedule:

Weeks 1	Weeks 7	Weeks 10		
through 6	through 9	through 12		
1 lozenge	1 lozenge	1 lozenge		
every	every	every		
1 to 2 hours	2 to 4 hours	4 to 8 hours		

Do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day. At the end of 12 weeks (3 months) you will have completed treatment.

#### FOR THE BEST CHANCE OF QUITTING, use Nicorette Lozenges on a regular schedule, using at least

MyQU Program brought to you by	it
Nicorette nicotine polacrilex lozenge Lozenge and GSK	To learn more visit www.quit.com/join-myquit or call 1-800-770-0708
Trademarks are owned by or licens	sed to the GSK group of companies. 13

# Nicorette has created the

# MyQuit program

to provide you with support wherever and whenever you need it. The program provides a network of tools to help you get through those tough moments.

- To Your Inbox: Stay motivated with our personalized email support program that sends encouragement and tips straight to your inbox.
- On The Web: Get helpful advice, tips and inspiration from ex-smokers on Quit.com.
- On The Go: Log cravings and track your personal progress on your mobile phone with the MyQuit App.

14

9 lozenges a day during the first 6 weeks. That will help your body better adjust to the lack of cigarettes and better help prevent cravings. Some people may need more lozenges to reduce their cravings. Do not exceed the recommended maximum daily dosage of 20 lozenges per day. Do not continuously use one lozenge after another, since this may cause you hiccups, heartburn, nausea or other side effects.

Do not eat or drink 15 minutes before using or while the lozenge is in your mouth.

# CUTTING BACK ON YOUR Nicorette LOZENGE USAGE.

The whole reason for using **Nicorette** Lozenges is to decrease and slowly eliminate your need for nicotine, while you control cravings. So, as the schedule on Page 12 indicates, you should gradually reduce the amount of Nicorette Lozenges you take per day. Some people find it easier to reduce by substituting ordinary sweets or sugar-free candy for some of the Nicorette Lozenges they would normally use. As time goes on, you can increase the number of pieces of candy as you further reduce your use of Nicorette Lozenges. It is important to complete treatment. If you still feel the need to use Nicorette Lozenges to keep from smoking after week 12, talk with your health care provider.

# MAKE QUITTING EASIER ON YOURSELF.

Soon after your quit date, parties, bars, celebrations, and socializing may all tempt you to smoke. Please remember 15

these tips to help you resist those urges and stay smoke-free.

### The Day You Quit Smoking:

- Look to your family and friends for support. Let them know what to do or avoid doing to help you quit.
- Throw away ALL cigarettes, ashtrays, matches, lighters. You don't need them. You don't want them and you want to make it difficult to go back.
- Keep yourself occupied. Take a walk. See a movie. See friends. Do anything to keep your mind off cigarettes.
- Calculaté all the money you'll save by not buying cigarettes. Probably well over \$1,000 a year! \$1,000 a year? Think of what you can spend it on!
- Know what situations are going to make you want to smoke. Plan now
   16

how you'll avoid them or deal with them so you don't smoke.

- Keep Nicorette Lozenges next to your bed so you're prepared when you get up. A lot of people get cravings first thing in the morning.
- Make an appointment to see your dentist and get the tobacco stains cleaned off. While you're getting rid of the evidence of cigarettes in the house, do the same for your teeth. Have clothes or drapes that smell of smoking cleaned.
- Now that your house is smoke-free, try to spend most of your time in smoke-free environments.
- If you usually smoked with coffee or alcohol, try to keep away from them for now. Remember you are also trying to break a habit.

- Smoking is a "hands-on" habit. So use something else to occupy your hands: a rubber band or a pen.
- Now's a good time to get active.
   Find activities to take your mind off cigarettes and relax. Take up jogging, swimming, or walking.
- Don't stress out about gaining weight. Dieting now may weaken your efforts to quit smoking. Eat sensibly and exercise daily; drink large quantities of water and fruit juices; this can help your chances of staying smoke-free.
- Laugh. Watch a sitcom. Read a comic book. It really helps.
- REMEMBER: Urges to smoke are temporary. They'll pass, even if you don't smoke.

### WHAT YOU CAN EXPECT.

As you are successful at staying smokefree, initially you will probably notice a few of the following typical withdrawal symptoms, so don't be surprised. Use of **Nicorette** Lozenges reduces these symptoms, but may not eliminate them entirely. They will go away with time. Stay focused on your goal of becoming an ex-smoker. Research shows that if you manage to avoid all smoking in the first week (that means not having a single puff), your chances of success increase dramatically.

The First Few Days. You may feel nervous or irritable or have difficulty concentrating during the first few days after you quit smoking. Your body needs time to regain balance. Initially, you might feel a little out of sorts, get headaches, feel light-headed, or have trouble sleeping. Your smoker's cough may get worse before it improves. But fear not, it's a positive sign. Coughing helps clean your lungs of the tar residue you got from smoking.

After a Couple of Weeks. Your confidence and ability to cope with urges to smoke should be getting stronger. But don't be over-confident and think you can smoke just one cigarette. Even now, having even a single puff can lead to a return to smoking cigarettes regularly. Be prepared, and remember why you wanted to stop smoking. Have you noticed that your sense of taste and smell has improved? You are probably coughing less and finding it easier to breathe. You've also probably noticed your withdrawal symptoms are \_\_\_\_\_\_ subsiding (though don't worry if they're still there: they last longer for some people). These are all positive signs that your body is getting used to your success at stopping smoking.

By The End of The First Month. You are less likely to have cravings for cigarettes as often. However sudden cravings may still happen, and when they do, be on your guard, as they can be strong and seem to come out of the blue. Be prepared for these challenging times. The key is do what you can so these unexpected cravings can't beat you. Keep focused on the ways non-smokers are more attractive than smokers. Their breath smells better. Their clothes and hair are fresher. Their teeth are cleaner and brighter. Their skin is less likely to wrinkle.

Not smoking around children and your friends is also healthier for them too.

WHAT IF YOU DO SLIP AND SMOKE?

"What if I relapse?" One cigarette is a slip-up, but it's not the end of the quit effort. Everybody slips at something. The key is this: forgive yourself and stop at that one cigarette. Don't let this slip ruin your good intentions, keep at your quit attempt. So, throw out your cigarettes and continue with your quit attempt, keeping in mind what went wrong and led to the slip.

If you do go back to smoking, certainly don't throw out your **Nicorette** Lozenges. Keep them for the next time you're ready to quit. In fact research says that even if you are back to smoking regularly the best thing you can do is learn and try again. Try to understand the reason you had those cigarettes that made you slip. That's important, because now you can plan better to deal with these moments next time. It's true you stumbled, but don't think of yourself as having failed. Encourage yourself by treating the last attempt as a learning experience, even a "trial run" for the real thing.

Take a look at the usage instructions and check that you used the **Nicorette** Lozenges correctly and for the full 12 weeks of the program. When you try again make sure you use enough and the right way. That way you'll be best equipped to deal with the unexpected cravings.

Don't forget; quitting isn't easy and it takes practice to do anything. Stopping smoking is no different.

### YOU'VE MADE IT.

Once your twelve week quitting program is over, you've taken your last **Nicorette** Lozenge. Now you are both cigarette and nicotine-free. Get up and give yourself a standing ovation. We mean it. Do you realize that you have just done a really difficult thing?

Now's a good time to think back on the process. Think of all your reasons for quitting smoking. Think of your goals. Think of how they're going to be a reality now.

Think of what you're going to do with your newly liberated cigarette money. The places you can now go smokefree. Think of the extra time you may have added to your life and what you can do with it. And although you may still experience the occasional temptation, 20 and cigarettes still want you back, think positively. Think forward. And consider yourself a proud non-smoker.

FREQUENTLY ASKED QUESTIONS. 1. When I stop smoking and start using Nicorette Lozenges how will I feel? Nicorette Lozenges help reduce cravings, but be prepared for some nicotine withdrawal symptoms. After you stop smoking they can begin almost at once and are normally at their strongest during the first three or four days. For some people, any of the following may occur:

- unexpected craving or urges for cigarettes
- anxiety, irritability, restlessness, mood changes, nervousness
- drowsiness
- trouble concentrating

increased appetite and weight gain

 headaches, muscular pain, constipation, fatigue

Nicorette Lozenges are designed to reduce the craving for nicotine you used to satisfy with cigarettes. Nicorette Lozenges can also help provide relief from other withdrawal symptoms such as irritability and nervousness.

#### 2. Are Nicorette Lozenges just swapping one type of nicotine addiction for another?

Nicorette Lozenges do contain nicotine, however there is probably less nicotine in your daily dose of lozenges than in your cigarettes. Nicorette Lozenges give you enough nicotine to help you combat the physical withdrawal symptoms so you can cope with the mental side of stopping smoking. Also, since the nicotine from the lozenges goes into your blood stream more slowly, it produces less of the effects of nicotine that people find rewarding. In fact, when used as directed in the 12 week program, **Nicorette** Lozenges gradually wean you off your dependence for both nicotine and cigarettes.

3. Can Nicorette Lozenges do any harm? Some people with conditions like heart disease or people taking prescription medicine for asthma or depression should not use this product without talking to their doctor — check the "IMPORTANT WARN NGS" on pages 6 to 7. You may also experience side effects such as hiccups, mouth or throat irritation, heartburn or other stomach problems such as nausea especially if Nicorette Lozenges are chewed or swallowed. In any case, Nicorette Lozenges do not contain the tar, carbon monoxide, and other toxins present in cigarette smoke.

### 4. Will I put on weight?

In the first couple of months after quitting smoking, some people do put on a few pounds. But think of it this way. Overall, you'll be healthier and look better. You can always tackle your weight by changing your diet and increasing the amount you exercise once you have gotten through the difficult part of stopping smoking.

# 5. Does taking Nicorette Lozenges cost more than smoking?

If you normally smoke a pack and a half a day, your total cost of using

Nicorette Lozenges during the 12 week period is about the same as smoking. But guess what? After you've finished the Nicorette Lozenge program all that money you used to spend on cigarettes is now savings. And think of the health issues you'll hopefully be able to avoid.

# 6. What if I have a cigarette and start smoking?

Don't panic. First, don't think badly of yourself. Throw away your cigarettes and forgive yourself. Then think about what went wrong and get back on track. In fact, people who have already tried to stop smoking are more likely to be successful the next time.

Reference ID: 4886336

### CHALLENGES TO WATCH FOR.

Once you quit smoking, you are likely to experience periodic, and sometimes intense, temptations to smoke. Certain situations present special challenges. Some common ones include:

#### Stress and upset.

When you are feeling stressed or upset, you may think a cigarette will make everything better. It won't. Find other ways to relax and unwind.

#### The blues.

You may be especially vulnerable when you feel bored or blue. Remember that having a cigarette will just make you feel worse.

#### Smoking cues.

Seeing cigarettes or watching other people smoke can trigger temptation.

Remember that you choose not to smoke anymore.

### Alcohol.

Drinking and smoking seem to go together, and alcoholic beverages may weaken your resolve, making drinking dangerous to your quit effort. Avoid drinking early in your quit effort, and try to drink with non-smokers.

### Automatic slips.

Sometimes you may find yourself preparing to smoke without even realizing it. Watch out for those moments when your hand seems to "automatically" reach for a cigarette.

Watch out for these situations: they can trigger a relapse. You probably know which one(s) are most dangerous for you; plan ahead to deal with the

situation effectively. Always remember that you're trying to break a habit, and the most important thing is to do something to combat the urge in these situations.

### COPING AFTER QUITTING.

The key to staying smoke-free is to prepare for and cope with challenges as they occur. If you find yourself tempted to smoke, do something!

Here are some things to consider:


- Escape. Leave the situation, even for a few minutes. Most temptations don't last long.
- Distract yourself. Get your mind off smoking. Think of something else or get busy with something.

- Relax. Don't let stress get to you. Think of pleasant, relaxing things; breathe slowly and regularly. Let the stress drain out of you.
- Talk yourself out of it. What you say to yourself matters. So, remind yourself how important it is for you to quit; remind yourself you can't have just one; or just command yourself to STOP.


# For more information please visit www.nicorette.com














	$\bigcirc$			$\bigcirc$			
	$\bigcirc$						
American Lung Association: 1-800-586-487 American Cancer Society: 1-800-227-2345 American Heart Association: 1-800-242-872 Quitting Buddy or Friend who has Quit:	WAL	ו נוניביבי בונת פתטע כי	ni 9011 901 16 dmozo1 printi 9002				or walking.
herican Lung Association: 1-800-586-48 merican Cancer Society: 1-800-227-234 herican Heart Association: 1-800-242-87 Quitting Buddy or Friend who has Quitt	ER I	ely. A ways remember that you're ob of si ping the do	to deal with the situation effectiv trying to break a habit, and the n	ant yo u back, think positively.	temp tation, and cigarettes stillw Think forward. An d consider you	ve. Find activities to take your Take up jogging, swimming,	Now's a good time to get act
Bud		ey can trigg er a re lapse. You ey can trigg er a re lapse. You	ht:sno itis ut for the se sit uations: the probably known which one (s) are	ed to your life and what you can	l voy seo sig e AT. yenom ettes cigarette vous seo so		occnb\table your hands: a rubber Smoking is a "hands-on" habi break a habit.
Ass cer s t Ass ldy o		<del>iomen ts wh en your h and se ems to</del> tte.	r ealizing it. watch out for a cigare "au tomaticall y" reach for a cigare		Think of what you're going to do	nember you are also trying to	
sociation: 1-800-586-4872 Society: 1-800-227-2345 sociation: 1-800-242-8721 or Friend who has Quit:	WALLET CARD	r eve arin g to smoke withou t eve n			They regoin g to be a reality work ing to be a reality work.	on ments.	your time in smoke-free envi
ety:		ir quiteffort, and try tod rink with	ыо - с и улога о циклид еа пу пу уло епо с с у улога о циклид еа пу пу уло	the process Think of all your	thin g? Now's a go od time to t hink back of		evidence of cigarettes in the
: 1-8 : 1-80 : 1-80	ÖR	p tog ether, and alcoholic be verages g drinking da ngerous to your q uit	may weaken your resolve, makin		nicotin e-free. Get up an d give you mean it Do you realize that you h	your dentist and get the your dentist and get the	Ma ke an ap poin tment to see tobacco stains cleaned off. W
o ha	₽ <u> </u> 0		temptation. Remember that you	vyouare both cigarette and	Once your twelve week quitting your last <b>Nko rette</b> Lozenge. Nov	lot of people g et cravings first	
	ē i i		Seeing cues Seeing cigarettes or wa tching ot			em or deal with them so you	Prian now now youniavoid tn brian ow now youniavoid tn brian keep Nicorette Lozenges n⊜
487 345 872	.? i		You may be especially vulnerable Remember that having a cigaret		s vsea finsi priittiup (taproft) nod	ng to make you want to smoke.	
			zəuldərT		righ t way. Th at way you'll be best une xpected cra vings.	save by n ot buying cigarettes.	Probably well over \$1,000 a y
		u pset, you may think a cigarette will Find other wa ysto relax a nd unwind.		adt to skeeks of the	Nico rette Lozen ges correctly a na Nico rette Lozen ges correctly a na program. When you tryagain ma		ب لاوو p your self occupied. Ta ke frien ds. Do any thing to k eep
			ommoo əmo 2.2 apri əlisi ara	off host nov te dt Voedo hae oan	run*for the real thing. Take a look at the usage instructi	htrays, matches, ligh ters. You ant them and you want to	make it difficult to go back. make it difficult to go back.
		kely to experience periodic, and so moke. Certain situations present	Once you du it smoking, you are	ailed. Encourage yourself by	teal is a state attent of the self as having the self of the se		what to do or a void doing to
			CHALLENGES TO WATCH	ot now you can plain better to	you slip. That's important, becaus you slip. That's important, becaus deal with these moments next th		priyom 2 u v voking
		nen think a bout w hat went wrong pple who have a lready tried to sto p	and get back on track. In fact, pe		nes uoy for the b est thing you can you Try to understand the reason you		t uoy tom at the yem prisileisos
		ly of yo urself. Th row away your	6. What if I have a clgarette and Don't panic. First, don't think bad	en if you a re backtosmoking brithe next time you're ready to	Nico rette Lozen ges. Keep them quit. In fact, research says that ev		<b>МА КЕ QUIT TING EASIE</b> l soon after your quit d ate, part
			to avoid.		If you do go backto smoking te	g after week 12, talk with your	realth care provider.
		you used to spend on dgarettes is you used to spend on dgarettes is alth issues you'll hopefull y be able		with your quit attempt, keeping	ruin your good intentions, keep a out your dg arettes and contin ue in mind what went wrong a nd le		ed the still feel the
		2 week period is about the same as	Nico rette Lozen ges dunn u the 1	dils sid t fet tet bon't let t his slip	of the q uit effort E verybod y slips forgive yo urself a nd stop a t that	of candy as you further reduce	ncrease the number of pieces
		Sea costmore than smoking?	5. Does taking Nicorette Lozen		NAPLIF IF VOU DOSLIPAHW Mhat if I relapse?"One cigarette	sier to red uce by substituting and y for some of the <b>Nicorette</b>	se ti briñ e lqo eq e mo 2. vabier
		g the amount you exercise once you			smoking around children and yo the m too.	for nicotine, while you control dule indicates, you should f <b>Nicorette</b> Lozenges you ta ke	eriavin gs. S o, as th e ab ove sche
			In the nist couple of months aire put on a few pounds. But think o healthier and look better You car	lesslikely to wrinkle. Not	r bris a cloth of ter. The ir cloth es and h cleaner a nd brighte r Their skin is	orette Lozeng es is to de crea se	
			4. Will I puton welght?	ou. Keep focused on the ways than smokers. Their breath	non-s mo kers a re more a ttra ctive une xpected cravings can't beat y	RNicorette LOZENGE	CUTTING BACK ONYOU
		carbon mon œi de, and ot her t oxins		he blue. Be prepared for these	stronging times. The key is do your of the strong of the s	before using or while the	o n ot ea t or drin k 15 minutes oze nge is in your mouth.
		pr swallowed. In any case, <b>Nico rette</b>	heart burn or other stoma dr prob	Vern a daive to nabb us tevewo	By The End of The Hrst Month. H. noff of settes as often .H d, o by off n of the n the y d o, b		via o ne lozeng e after anot her, Diccups, hea it burn, na usea o r
		a ctor — طه ودلا the <b>"MP OR TA NT</b> U ser's Guide. You may also ccups, mouth or throat infitation,	vitt to thort a the front of this	Pricessa at stopping smoking.	your body is getting used to you		sagnazol 0 2 to age sob y list
		or de pre ssion sh ould not use th is e heart disease or people ta king	So me people with condition slike prescription medicine for ast hms	don't won'y if they 're still there:	and of the state of the second state of the se	ed more lozen ges to reduce and better help prevent	adj ust to the lack of dg are ttes adj ust to the lack of dg are ttes
nies or its licensor. 84L0058J2	©2021 GSK group of compa	yyarmi ?	3. Can Nico rette Lozenges do a	g nibriñ brie szel g nidguoz vid so	to stop smoking. Ha ve you not ica smol ha s improved? You ve are p ro it easier t o breathe. You've also p	e s seprezol e te ste el te prisu, e	asy during the first 6 weeks. Th Lose nge s on a regular schedul day during the first 6 weeks. Th
		d reward ing. In fact, whe n used as <b>Nicorette</b> Lozenges gra dually or both nicotine and ciga rettes.	effects of nicorine that pendence f directed in the 12 week program wean you off your de pendence f	and remember why you wanted n lead to a retum to smoking	cigarettes regularly. Be prepared		gron avail hiw nov (arthrom E F <b>OR THE BEST CHANCE OF Q</b>
			goes into your blood stream mor	ettin g strong er. But don't b e smoke just on e cigarette. Even	with urges to smoke should be g	wy. At the end of 12 weeks	biscol činschitero miserio a con o C biscol segnesco lo Sinschitero miserio a arresteria di linu usu (schroom Si
		ns so yo u can cop é with the mental bugh nicotine to h ép you combat	<b>Nico rette</b> Lozen ges give you en o		your lungs of the tar residue your After a Couple of Weeks. Your c	vours 4 to 8 hours	1 to 2 hours 2 to 4
		icotine, however there is probably lozeng es than in your cigarettes.	Nico rette Lozen ges d o con tain n		it improves. But fear no t, it's a po		Weeks1 to 6 Weeks 1 lozengeevery 1 lozeng
		ə u pi o ayu yo əd x əu o buld dawı	ג. A ne Nicorrette Lozenges just: מלמו כלוס חלטר אסל אפו?	o reg ain bal ance. Initia IIy, you bead aches, feel lig ht-he aded, or	smoking. Your bod y needs time t might feel a little out of sorts, ge	ອບາ ດາ ມົນແກ່ງ ຄະຫຼຸ	o llowin d dosa de sche dule: Mubrours, use <b>Nkorette</b> Loze
tisiv se se please visit corette.com		viild strif as in the blift			əəi yamı voy <b>.200 Wə Traf Taha</b> Hitoriny concentrating during th		io reduce cravings or urgesto
ti sin oscola aoitea	ano fai o so ta so d	to reduce the craving for nicotine s. <b>Nico rette</b> Lozen ges can a lso hel p	you used to satisfy with cigarette	k (that means not having a sin gle		l completely dissolved (about	
one; or just command yourself			• he adaches, mus cu lar pain, cons	h shows t hat if you manag e to fo cu sed on yo ur goal of	The y will go a way with time. Stay be coming an exismoker. Researc	n or tingl ing sen sation.	<b>on od</b> .priwollsws 9 siminiM 1 sw s f 9 y s m v f ee l a wai 2) Occasionally move the lozer
you sayto yourselfmatters. So, syou sayto yourselfmatters. So, ndir; reminal	<ul> <li>Talk yourself out of it. What</li> </ul>	uiec	<ul> <li>trou ble concentrating</li> <li>increased app etite and weight</li> </ul>	Use of <b>Nico rette</b> Lozenges	probably notice a few of the follo symptoms, so don't be surprised reduces these symptoms, but ma		e) əvlossi b yhvols oʻt əpnəzol
o yo <b>u</b> . Think of pleasant, relaxing reg <b>u</b> lariy. Let th e stress drain out			<ul> <li>anxiety, it rita bility, rest les sness,</li> <li>anxiety, it rita bility, rest les sness,</li> </ul>		Plot and the stew of the following the following state of the second sec		/ourquitdate. 1) Remove the Nicorette Loze
v with something.	<ul> <li>Distract yo unself. Get your 1 somet hing else or g et b usy</li> <li>1 foot a stress det to b</li> </ul>		<ul> <li>up sected craving or urges fo</li> </ul>		an don't smoke.	nalq shirt dika the plan	asteafter a day or two. Rem vill help you quit Begin usin
	temp tations don't last long	סטט פון אין געטט אין			P DEAD IN WILLING STORY OF SAUGHERS	lcorette Lo zenges are a	asteo fithe of a construction of the state of the source o
o ufind yourself tempted to see some things to consider:	challenges as they occur. If ye	cravings, but be prepared for some	Will i feel? Ni co rette Lozen ges h elp re duce	fr ee.	drink large quan titles of water a your chances of stayin g smoke • Laug h. Watch a sitcom. Read a c	ərbiə xiliton o binov îl quəvbi	<b>Vico rette</b> Lozen ges correctly. M PORTA UT: D on't wo rry or
ritiw a qoo briand for and cop a vita		rtusing Nicorette Lozen geshow. SNOITS:			your efforts to quit smoking. Ea		Loze nges. Refer to them often Loze nges. Refer to them often

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD 11/09/2021 11:14:13 PM