




BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	6.5pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	7pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	




BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	6.5pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	7pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Vertical PDP Graphics Orientation (Recommended)


GLUE ASSISTS FOR POSITION ONLY. PATTERN MAY VARY BY MANUFACTURING PROCESS

MFG Glue Assist No Varnish


Reads this way
Horizontal PDP Graphics Orientation (Mandatory)



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8.1pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	8pt		
Horizontal Scale: 100%		Average Kerning: 0%	




BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	




BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	




BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	




BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8.1pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	8pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Do not use if printed safety seal under cap is torn or missing.

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

NDC 69452-179-78

Strength to last 12 hours

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

80 CAPSULES



Actual Size



Lot No.:
Exp. Date:

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:	<ul style="list-style-type: none"> take 1 capsule every 8 to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8- to 12-hour period do not exceed 3 capsules in a 24-hour period
children under 12 years:	<ul style="list-style-type: none"> ask a doctor

Drug Facts (continued)

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Do not use if printed safety seal under cap is torn or missing.

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

NDC 69452-179-78

Strength to last 12 hours

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Compare to the active ingredient of ALEVE® Liquid Gels ***

80 CAPSULES



Actual Size



Lot No.:
Exp. Date:

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:	<ul style="list-style-type: none"> take 1 capsule every 8 to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8- to 12-hour period do not exceed 3 capsules in a 24-hour period
children under 12 years:	<ul style="list-style-type: none"> ask a doctor

Drug Facts (continued)

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

***This product is not manufactured or distributed by the owners of ALEVE® Liquid Gels.

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

NO COATING

Naproxen Sodium Capsules, 220 mg (NSAID) Pain Reliever/Fever Reducer

Do not use if printed safety seal under cap is torn or missing.

EASY OPEN ARTHRITIS CAP
Strength to last 12 hours
Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

NDC 69452-179-78

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

80 CAPSULES

NO COATING

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients
FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

EASY OPEN ARTHRITIS CAP

80 CAPSULES

Strength to last 12 hours

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Manufactured for: Bionpharma Inc. 600 Alexander Road, Princeton, NJ 08540



Lot No:
Exp. Date:

NO COATING

Naproxen Sodium Capsules, 220 mg (NSAID) Pain Reliever/Fever Reducer

Do not use if printed safety seal under cap is torn or missing.

EASY OPEN ARTHRITIS CAP
Strength to last 12 hours

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

NDC 69452-179-78

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Compare to the active ingredient of ALEVE® Liquid Gels***

80 CAPSULES

NO COATING

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients
FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

***This product is not manufactured or distributed by the owners of ALEVE® Liquid Gels.

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer


EASY OPEN ARTHRITIS CAP

80 CAPSULES

Strength to last 12 hours


BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Manufactured for:
Bionpharma Inc.
600 Alexander Road,
Princeton, NJ 08540




3 69452 17978 9

Lot No:
Exp. Date:



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8.1pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	8pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8.1pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Bullines:	2.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	8pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Do not use if printed safety seal under cap is torn or missing.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:	<ul style="list-style-type: none"> take 1 capsule every 8 to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8- to 12-hour period do not exceed 3 capsules in a 24-hour period
children under 12 years:	<ul style="list-style-type: none"> ask a doctor

Drug Facts (continued)

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured by:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Strength to last 12 hours

100 CAPSULES

Actual Size



NDC 69452-179-20

Drug Facts

Active ingredient (in each capsule)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers



Lot No.:
Exp. Date:

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Do not use if printed safety seal under cap is torn or missing.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:	<ul style="list-style-type: none"> take 1 capsule every 8 to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8- to 12-hour period do not exceed 3 capsules in a 24-hour period
children under 12 years:	<ul style="list-style-type: none"> ask a doctor

Drug Facts (continued)

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

***This product is not manufactured or distributed by the owners of ALEVE® Liquid Gels.

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Strength to last 12 hours

100 CAPSULES

Actual Size

Compare to the active ingredient of ALEVE® Liquid Gels ***

NDC 69452-179-20

Drug Facts

Active ingredient (in each capsule)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers



Lot No.:
Exp. Date:

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Do not use if printed safety seal under cap is torn or missing.

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

NDC 69452-179-22

NDC 69452-179-22

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

120 CAPSULES

120 CAPSULES



Lot No.:
Exp. Date:

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*. Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes
Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients
FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		

Horizontal Scale: 100% Average Kerning: 0%

Do not use if printed safety seal under cap is torn or missing.

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

Strength to last 12 hours

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

NDC 69452-179-22

Strength to last 12 hours

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

NDC 69452-179-22

Compare to the active ingredient of ALEVE® Liquid Gels ***

120 CAPSULES



Actual Size

Compare to the active ingredient of ALEVE® Liquid Gels ***

120 CAPSULES



Actual Size



Lot No.:
Exp. Date:

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*. Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes
Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients
FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

***This product is not manufactured or distributed by the owners of ALEVE® Liquid Gels.

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521


BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		

Horizontal Scale: 100% Average Kerning: 0%



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8.1pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	8pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Drug Facts (continued)

Active ingredient (in each capsule) Naproxen sodium 220 mg (NSAID). Pain reliever/fever reducer.

Purposes Naproxen sodium 220 mg (NSAID). Pain reliever/fever reducer.

Uses Temporarily relieves minor aches and pains due to:

- the common cold
- headache
- backache
- muscle aches
- menstrual cramps
- toothache

Warnings Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day when using this product

 Take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- If you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease the benefit of aspirin

● taking any other drug

Drug Facts

Directions

- Take 1 capsule every 8 to 12 hours while symptoms last.
- For the first dose you may take 2 capsules within the first hour and other children and adults.
- Take 1 capsule every 8 to 12 hours while symptoms last.
- Drink a full glass of water with each dose.
- Taken with food, this product may take longer to work.

Other information

- Each capsule contains sodium 20 mg.
- Store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- Protect from light.
- Read all directions and warnings before use. Keep cap on.

Questions or comments?

- Proprietary glycol, purified water, sorbitan, sorbitol, acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol.

1-888-235-2466 (Mon-Fri 9AM - 5PM CST)

Naproxen Sodium Capsules, 220 mg (NSAID)
 Pain Reliever/Fever Reducer

Strength to last 12 hours

2 PACK 2 x 80 CT

Naproxen Sodium Capsules, 220 mg (NSAID)
 Pain Reliever/Fever Reducer

2 PACK 2 x 80 CT

Naproxen Sodium Capsules, 220 mg (NSAID)
 Pain Reliever/Fever Reducer

2 PACK 2 x 80 CT

Naproxen Sodium Capsules, 220 mg (NSAID)
 Pain Reliever/Fever Reducer

2 PACK 2 x 80 CT

Do not use if printed safety seal under cap is torn or missing.

Lot No.:
Exp. Date:

LOT & EXP

Naproxen Sodium Capsules, 220 mg (NSAID)
 Pain Reliever/Fever Reducer

Strength to last 12 hours

160 CAPSULES

Naproxen Sodium Capsules, 220 mg (NSAID)
 Pain Reliever/Fever Reducer

Strength to last 12 hours

160 CAPSULES

2 Bottles x 80 Capsules Each

Naproxen Sodium Capsules, 220 mg (NSAID)
 Pain Reliever/Fever Reducer

Strength to last 12 hours

160 CAPSULES

Manufactured for:
Bionpharma Inc.
 600 Alexander Road,
 Princeton, NJ 08540

00000000000000

L0000000
R0521

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	10pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	9pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Do not use if printed safety seal under cap is torn or missing.

Drug Facts

Active ingredient (in each capsule)
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Purposes
 Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients
 FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
 600 Alexander Road,
 Princeton, NJ 08540

L0000000
 R0521

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Strength to last 12 hours

160 CAPSULES

Actual Size

NDC 69452-179-79

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Strength to last 12 hours

160 CAPSULES

Actual Size

NDC 69452-179-79



Lot No.:
 Exp. Date:

BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		

Horizontal Scale: 100% Average Kerning: 0%

Do not use if printed safety seal under cap is torn or missing.

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

NDC 69452-179-79

160 CAPSULES

Compare to the active ingredient of ALEVE® Liquid Gels ***

Compare to the active ingredient of ALEVE® Liquid Gels ***

NDC 69452-179-79

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Strength to last 12 hours

160 CAPSULES



Actual Size



Actual Size



Lot No.:
Exp. Date:

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*. Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk; if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients
FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

***This product is not manufactured or distributed by the owners of ALEVE® Liquid Gels.

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

BIONPHARMA


Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique


Drug Facts:	8pt	Leading:	7.2pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	.6pt
Body Text:	6pt	Hairlines:	.2pt
Drug Facts (continued):	7pt		

Horizontal Scale: 100% Average Kerning: 0%

Front




Back




BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	6.5pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	7pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Front



Back



BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	8pt	Leading:	6.5pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	7pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	7pt		

Horizontal Scale: 100% Average Kerning: 0%

1.75" 4.048" 2.827"

Do not use if printed safety seal under cap is torn or missing.

Manufactured for:
Bionpharma Inc.
600 Alexander Road,
Princeton, NJ 08540 R0521

LOT
EXP

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

STRENGTH TO LAST 12 HOURS

160 CAPSULES
(L QUID FILLED CAPSULES)

ALEVE

STRENGTH TO LAST 12 HOURS

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

STRENGTH TO LAST 12 HOURS

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes
Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - menstrual cramps
 - temporarily reduces fever
- muscular aches
- backache
- headache
- toothache
- the common cold

Internal die slit on face stock only - allows Top ply to break free while maintaining lot/exp area to base label.

1.25"

LOT / EXP NO PRINT AREA 5/8" Coating Area

TOP

0.3125" 1.7275" 6.585"

100% Proprietary Coating 2 - Mirrored Plates

BEHIND TOP

3.25"

Drug Facts (continued)

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Drug Facts (continued)

- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - sturred speech
 - weakness in one part or side of body
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

 adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

 children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
1-888-235-2466 (Mon-Fri 9AM-5PM EST)


BASE

100% & 60% Proprietary Coating 1


1.7275" 6.585" 0.3125" 8.625"

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	14pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	8pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Front




Back




BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	6.5pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	7pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Front




Back




BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	6.5pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	7pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Front



Back



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	8pt	Leading:	6.5pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	7pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer ● right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

under 12 years	● ask a doctor
adults and children 12 years and older	<ul style="list-style-type: none"> ● take 1 capsule every 8 hours to 12 hours while symptoms last ● for the first dose you may take 2 capsules within the first hour ● do not exceed 2 capsules in any 8-hour to 12-hour period ● do not exceed 3 capsules in a 24-hour period

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F). ● protect from light
- read a 1 directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
call toll-free 1-888-235-2466 (Mon-Fri 9 AM-5 PM EST)

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Actual Size
NP1
20 capsules**
liquid-filled capsule

12 hour menstrual pain relief

temporarily relieves minor cramps, backache, and headache

NDC 691452-270-11

Drug Facts

Active ingredient (in each capsule) Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to: ● menstrual cramps ● muscular aches ● headache ● backache ● minor pain of arthritis ● toothache ● the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ● hives ● facial swelling ● asthma (wheezing) ● shock ● skin reddening ● rash ● blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

L0000000
R0521

Manufactured for:
BION PHARMA
600 Alexander Road, Princeton, NJ 08540

3 69452 27011 0

Lot No.:
Exp. Date:

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



naproxen sodium capsules, 220 mg (NSAID)
 pain reliever/fever reducer

Do not use if printed safety seal under cap is torn or missing.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer ● right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

under 12 years	● ask a doctor
adults and children 12 years and older	<ul style="list-style-type: none"> ● take 1 capsule every 8 hours to 12 hours while symptoms last ● for the first dose you may take 2 capsules within the first hour ● do not exceed 2 capsules in any 8-hour to 12-hour period ● do not exceed 3 capsules in a 24-hour period

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F). ● protect from light
- read a 1 directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
 call toll-free 1-888-235-2466 (Mon-Fri 9 AM-5 PM EST)

Drug Facts

Active ingredient (in each capsule) Purposes
 Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to: ● menstrual cramps ● muscular aches ● headache ● backache ● minor pain of arthritis ● toothache ● the common cold ● temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ● hives ● facial swelling ● asthma (wheezing) ● shock ● skin reddening ● rash ● blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

naproxen sodium capsules, 220 mg (NSAID)
 pain reliever/fever reducer

***compare to the active ingredient in Menstrido!™

Actual Size
 NP1
 20 capsules***
 liquid-filled capsule

12 hour menstrual pain relief

temporarily relieves minor cramps, backache, and headache

NDC 69452-270-11

L0000000
 R0521

***This product is not manufactured or distributed by the owners of Menstrido!™.

Manufactured for:
BION PHARMA
 600 Alexander Road, Princeton, NJ 08540

Lot No.:
 Exp. Date:

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Do not use if printed safety seal under cap is torn or missing.

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
 - the smallest effective dose should be used
 - drink a full glass of water with each dose
 - if taken with food, this product may take longer to work
- | | |
|--|--|
| under 12 years | ask a doctor |
| adults and children 12 years and older | take 1 capsule every 8 hours to 12 hours while symptoms last |
| | for the first dose you may take 2 capsules within the first hour |
| | do not exceed 2 capsules in any 8-hour to 12-hour period |
| | do not exceed 3 capsules in a 24-hour period |

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light.
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?

call toll-free 1-888 235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

Drug Facts

Active ingredient (in each capsule) Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)
Purposes Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - muscular aches
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

 take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

40 capsules**

**liquid-filled capsule

NP1

Actual Size

12 hour menstrual pain relief


temporarily relieves minor cramps, backache, and headache

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

NDC 69482-270-15



Lot No.:
Exp. Date:



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Do not use if printed safety seal under cap is torn or missing.

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

under 12 years	ask a doctor
adults and children 12 years and older	<ul style="list-style-type: none"> take 1 capsule every 8 hours to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8-hour to 12-hour period do not exceed 3 capsules in a 24-hour period

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
call toll-free 1-888 235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

Drug Facts

Active ingredient (in each capsule)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - muscular aches
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

 take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Actual Size
NP1

40 capsules**
*liquid-filled capsule

12 hour menstrual pain relief

temporarily relieves minor cramps, backache, and headache

***compare to the active ingredient in Menstridol™

NDC 69482-270-15

***This product is not manufactured or distributed by the owners of Menstridol™.

3 69482 27015 8

Lot No.:
Exp. Date:

Do not use if printed safety seal under cap is torn or missing.

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

under 12 years

- ask a doctor

adults and children 12 years and older

- take 1 capsule every 8 hours to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8-hour to 12-hour period
- do not exceed 3 capsules in a 24-hour period

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F). ● protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?

call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

**Liquid-filled capsule

80 capsules**



Actual Size

NP1

12 hour menstrual pain relief

temporarily relieves minor cramps,
backache, and headache

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

NDC 69452-270-78

Drug Facts

Active ingredient (in each capsule)

Naproxen sodium 220 mg
(naproxen 200 mg) (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - headache
 - minor pain of arthritis
 - the common cold
- temporarily reduces fever
- muscular aches
- backache
- toothache

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.



Lot No.:
Exp. Date:

BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia

Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		

Horizontal Scale: 100%

Average Kerning: 0%

Do not use if printed safety seal under cap is torn or missing.

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

under 12 years

- ask a doctor

adults and children 12 years and older

- take 1 capsule every 8 hours to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8-hour to 12-hour period
- do not exceed 3 capsules in a 24-hour period

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F). ● protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?

call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

**liquid-filled capsule

80 capsules**



Actual Size

NP1

12 hour menstrual pain relief

temporarily relieves minor cramps,
backache, and headache

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

NDC 69452-270-78

***compare to the active ingredient in Menstridol™

Drug Facts

Active ingredient (in each capsule)

Naproxen sodium 220 mg
(naproxen 200 mg) (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - headache
 - minor pain of arthritis
 - the common cold
 - temporarily reduces fever
 - muscular aches
 - backache
 - toothache

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

***This product is not manufactured or distributed by the owners of Menstridol™.



Lot No.:
Exp. Date:

BIONPHARMA


Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia

Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		

Horizontal Scale: 100%

Average Kerning: 0%



Do not use if printed safety seal under cap is torn or missing.

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

Adults and children 12 years and older	<ul style="list-style-type: none"> Take 1 capsule every 8 hours to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8-hour to 12-hour period do not exceed 3 capsules in a 24-hour period
Children under 12 years	<ul style="list-style-type: none"> ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients
FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Actual Size
NP1
12 hour back & muscle pain relief

50 capsules**
**hydro-filled capsule

for temporary relief of minor back and muscle aches and pains

NDC 60452-260-16

Drug Facts

Active ingredient (in each capsule) Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*	Purposes Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug
---	--

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - muscular aches
 - minor pain of arthritis
 - menstrual cramps
 - headache
 - toothache
 - the common cold
 - temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery



Lot No.:
Exp. Date:

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

Adults and children 12 years and older	<ul style="list-style-type: none"> take 1 capsule every 8 hours to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8-hour to 12-hour period do not exceed 3 capsules in a 24-hour period
Children under 12 years	<ul style="list-style-type: none"> ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients
FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
call toll-free 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540

L0000000
R0521

Drug Facts

Active ingredient (in each capsule)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - muscular aches
 - minor pain of arthritis
 - menstrual cramps
 - headache
 - toothache
 - the common cold
 - temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

Do not use if printed safety seal under cap is torn or missing.

50 capsules**

Actual Size

NP1

12 hour back & muscle pain relief


for temporary relief of minor back and muscle aches and pains

***Compare to the active ingredient in ALEVE® BACK & MUSCLE PAIN

NDC 60452-269-16

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

***This product is not manufactured or distributed by the owners of ALEVE® Back & Muscle pain.



3 69452 26916 9

Lot No.:
Exp. Date:

naproxen sodium capsules, 220 mg (NSAID)
pain reliever/fever reducer

BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia			
Drug Facts:	8pt	Leading:	7pt
Header:	7pt	Bullets:	5pt
Subheader:	6pt	Barlines:	0.6pt
Body Text:	6pt	Hairlines:	0.2pt
Drug Facts (continued):	7pt		
Horizontal Scale: 100%		Average Kerning: 0%	

Front

.125" corner radius

2.375"

NDC 69452-269-20

naproxen sodium capsules, 220 mg (NSAID)

pain reliever/fever reducer

12 hour back & muscle pain relief

Actual Size

NP1

for temporary relief of minor back and muscle aches and pains

100 capsules**

**liquid-filled capsule

Drug Facts

Active ingredient (in each capsule) Naproxen sodium 220 mg (naproxen 200 mg) (NSA D) *.....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Purposes

Uses ● temporarily relieves minor aches and pains due to:
 ● backache ● muscular aches ● minor pain of arthritis ● menstrual cramps ● headache
 ● toothache ● the common cold
 ● temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ● hives ● facial swelling ● asthma (wheezing) ● shock
 ● skin reddening ● rash ● blisters
Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed
Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ● if you have ever had an allergic reaction to any other pain reliever/fever reducer ● right before or after heart surgery

6 9 4 5 2 2 6 9 2 0 6
L000000 RB21
3

Non Varnish Area

6.00"

Back

2.375"

Drug Facts (continued)

Ask a doctor before use if: ● the stomach bleeding warning applies to you ● you have a history of stomach problems, such as heartburn ● you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ● you are taking a diuretic ● you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are: ● under a doctor's care for any serious condition ● taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ● taking any other drug

When using this product: ● take with food or milk if stomach upset occurs

Stop use and ask a doctor if:
 ● you experience any of the following signs of stomach bleeding:
 ● heartburn ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ● you have symptoms of heart problems or stroke:
 ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling
 ● pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● you have difficulty swallowing ● it feels like the capsule is stuck in your throat ● redness or swelling is present in the painful area ● any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ● do not take more than directed ● the smallest effective dose should be used ● drink a full glass of water with each dose ● if taken with food, this product may take longer to work

Adults and children 12 years and older:
 ● take 1 capsule every 8 hours to 12 hours while symptoms last ● for the first dose you may take 2 capsules within the first hour

Children under 12 years:
 ● do not exceed 2 capsules in any 8-hour to 12-hour period ● do not exceed 3 capsules in a 24-hour period ● ask a doctor

Other information ● each capsule contains: sodium 20 mg ● store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F). ● protect from light ● read all directions and warnings before use.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol

Questions or comments?
 call toll-free 1-888-235-2466 (Mon - Fri 9AM - 6PM EST)

6.00"

BIONPHARMA	
Labeling Format Information:	
<p>Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia</p>	
Drug Facts:	8pt
Header:	7pt
Subheader:	6pt
Body Text:	6pt
Drug Facts (continued):	7pt
Leading:	5pt
Bullets:	5pt
Barlines:	0.6pt
Hairlines:	0.2pt
Horizontal Scale: 100% Average Kerning: 0%	

Reference ID: 4903813

Front

.125" corner radius

2.375"

***Compare to the active ingredient in
ALEVE® BACK & MUSCLE PAIN

NDC 69452-269-20

naproxen sodium
capsules, 220 mg (NSAID)

pain reliever/fever reducer

12 hour back & muscle pain relief

Actual Size

for temporary relief of
minor back and muscle
aches and pains

100 capsules**

**liquid-filled capsule

Do not use if printed safety seal
under cap is torn or missing.

6.00"

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D) *.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses ● temporarily relieves minor aches and pains due to:
● backache ● muscular aches ● minor pain of arthritis ● menstrual cramps ● headache
● toothache ● the common cold
● temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ● hives ● facial swelling ● asthma (wheezing) ● shock
● skin reddening ● rash ● blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSA Ds (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed
Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ● if you have ever had an allergic reaction to any other pain reliever/fever reducer
● right before or after heart surgery

Non Varnish Area

PEEL HERE →

***This product is not manufactured or distributed by the owners of ALEVE® Back & Muscle Pain.
Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540
Lot#EYEP
L0000000 RB521
6 9 4 5 2 2 6 9 2 0 6
3

Back

2.375"

Drug Facts (continued)

Ask a doctor before use if ● the stomach bleeding warning applies to you ● you have a history of stomach problems, such as heartburn ● you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ● you are taking a diuretic ● you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are ● under a doctor's care for any serious condition ● taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ● taking any other drug

When using this product ● take with food or milk if stomach upset occurs

Stop use and ask a doctor if
● you experience any of the following signs of stomach bleeding:
● heartburn ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better
● you have symptoms of heart problems or stroke:
● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling
● pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● you have difficulty swallowing ● it feels like the capsule is stuck in your throat ● redness or swelling is present in the painful area ● any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ● do not take more than directed ● the smallest effective dose should be used ● drink a full glass of water with each dose ● if taken with food, this product may take longer to work

Adults and children 12 years and older
● take 1 capsule every 8 hours to 12 hours while symptoms last
● for the first dose you may take 2 capsules within the first hour
● do not exceed 2 capsules in any 8-hour to 12-hour period
● do not exceed 3 capsules in a 24-hour period

Children under 12 years
● ask a doctor

Other information ● each capsule contains: sodium 20 mg ● store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F). ● protect from light ● read all directions and warnings before use.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
call toll-free 1-888-235-2466 (Mon - Fri 9AM - 6PM EST)

6.00"

BIONPHARMA Labeling Format Information:	
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique Georgia	
Drug Facts:	8pt
Header:	7pt
Subheader:	6pt
Body Text:	6pt
Drug Facts (continued):	7pt
Leading:	5pt
Bullets:	5pt
Barlines:	0.6pt
Hairlines:	0.2pt
Horizontal Scale: 100%	Average Kerning: 0%

Reference ID: 4903813

Drug Facts (continued)

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:	<ul style="list-style-type: none"> ● take 1 capsule every 8 to 12 hours while symptoms last ● for the first dose you may take 2 capsules within the first hour ● do not exceed 2 capsules in any 8- to 12-hour period ● do not exceed 3 capsules in a 24-hour period
children under 12 years:	<ul style="list-style-type: none"> ● ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- read all directions and warnings before use. Keep carton.

Inactive ingredients
FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

***compare to the active ingredient in ALEVE® Liquid Gels

A+health

naproxen sodium capsules, 220 mg (NSAID)

pain reliever/fever reducer

80 capsules**

**liquid-filled capsule

strength to last 12 hours

A+health

naproxen sodium capsules, 220 mg (NSAID)

Do not use if printed safety seal under cap is torn or missing.

A+health

naproxen sodium capsules, 220 mg (NSAID)

pain reliever/fever reducer

80 capsules**

**liquid-filled capsule

strength to last 12 hours

**compare to the active ingredient in ALEVE® Liquid Gels

Drug Facts

Active ingredient (in each capsule)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSA D)*	Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - menstrual cramps
 - the common cold
 - temporarily reduces fever
 - muscular aches
 - headache
 - backache
 - toothache

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- rash
- asthma (wheezing)
- shock
- skin reddening
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

R0521
L0000000

**This product is not manufactured or distributed by the owners of ALEVE® Liquid Gels.

Manufactured for:
BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540



Lot: LOT & EXP
Exp. Date:

BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	10pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2.431pt
Body Text:	6pt	Hairlines:	.486pt
Drug Facts (continued):	9pt		

Horizontal Scale: 94% Average Kerning: 0%

.125" corner radius


Front

***compare to the active ingredient in ALEVE® Liquid Gels

a+health

naproxen sodium capsules, 220 mg (NSAID)

pain reliever/fever reducer



120 capsules** strength to last 12 hours

**liquid-filled capsule

Do not use if printed safety seal under cap is torn or missing.

<p>Drug Facts</p> <p>Active ingredient (in each capsule) Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses</p> <ul style="list-style-type: none"> ● temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ● minor pain of arthritis ● headache ● temporarily reduces fever ● muscular aches ● toothache ● backache ● the common cold ● menstrual cramps <p>Warnings</p> <p>Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ● hives ● facial swelling ● asthma (wheezing) ● shock</p> <p>● skin reddening ● rash ● blisters.</p> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed</p> <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use ● if you have ever had an allergic reaction to any other pain reliever/fever reducer ● right before or after surgery</p>	<p>Purposes</p> <p>69452125922</p> <p>LOTEXP</p> <p>1</p> <p style="text-align: center;">Non Varnish Area</p> <p style="text-align: center;">PEEL HERE</p>
--	---

***This product is not manufactured or distributed by the owners of ALEVE® Liquid Gels.
Manufactured for: BIONPHARMA
800 Alexander Road, Princeton, NJ 08540
L0000000
R0621

6.25"

Back

2.375"

Drug Facts (continued)

Ask a doctor before use if: ● stomach bleeding warning applies to you ● you have a history of stomach problems, such as heartburn ● you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ● you are taking a diuretic ● you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are: ● under a doctor's care for any serious condition ● taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ● taking any other drug

When using this product: ● take with food or milk if stomach upset occurs

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● redness or swelling is present in the painful area
- any new symptoms appear ● you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ● do not take more than directed ● the smallest effective dose should be used ● drink a full glass of water with each dose ● if taken with food, this product may take longer to work

adults and children 12 years and older:	take 1 capsule every 8 to 12 hours while symptoms last ● for the first dose you may take 2 capsules within the first hour	● do not exceed 2 capsules in any 8- to 12-hour period
children under 12 years:	do not exceed 3 capsules in a 24-hour period	ask a doctor

Other information ● each capsule contains sodium 20 mg ● store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F). ● protect from light ● read all directions and warnings before use.

Inactive ingredients: FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)

6.25"

BIONPHARMA	
Labeling Format Information:	
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique	
Drug Facts:	9pt
Header:	8pt
Subheader:	6pt
Body Text:	6pt
Drug Facts (continued):	8pt
Leading:	6.5pt
Bullets:	5pt
Barlines:	2pt
Hairlines:	.5pt
Horizontal Scale: 85%	
Average Kerning: 0%	

Reference ID: 4903813

.125" corner radius


Front

***compare to the active ingredient
in ALEVE® Liquid Gels

a+health

**naproxen sodium
capsules, 220 mg
(NSAID)**

pain reliever/fever reducer




180 capsules** strength to last 12 hours

**liquid-filled capsule

Do not use if printed safety seal under cap is torn or missing.

<p>Drug Facts</p> <p>Active ingredient (in each capsule) Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses</p> <ul style="list-style-type: none"> ● temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ● minor pain of arthritis ● headache ● temporarily reduces fever ● muscular aches ● toothache ● backache ● the common cold ● menstrual cramps <p>Warnings</p> <p>Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ● hives ● facial swelling ● asthma (wheezing) ● shock</p> <p>● skin reddening ● rash ● blisters.</p> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems</p> <p>● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed</p> <p>Heart attack and stroke warning: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use ● if you have ever had an allergic reaction to any other pain reliever/fever reducer</p> <p>Do not use ● right before or after surgery</p>	<p>Purposes</p> <p>Pain reliever/fever reducer</p>
--	---



69452125925 2

LotEXP

L0000000

ROB21

PEEL
HERE

Non Varnish Area

2.375"

6.25"

Back

2.375"

6.25"

Drug Facts (continued)	
<p>Ask a doctor before use if ● stomach bleeding warning applies to you ● you have a history of stomach problems, such as heartburn ● you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ● you are taking a diuretic ● you have problems or serious side effects from taking pain relievers or fever reducers</p> <p>Ask a doctor or pharmacist before use if you are ● under a doctor's care for any serious condition ● taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ● taking any other drug</p> <p>When using this product ● take with food or milk if stomach upset occurs</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ● you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ● you have symptoms of heart problems or stroke: <ul style="list-style-type: none"> ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling ● pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● redness or swellings present in the genital area ● any new symptoms appear ● you have difficulty swallowing ● it feels like the capsule is stuck in your throat <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Directions ● do not take more than directed ● the smallest effective dose should be used ● drink a full glass of water with each dose ● if taken with food, this product may take longer to work</p> <p>adults and children 12 years and older: ● take 1 capsule every 8 to 12 hours while symptoms last ● for the first dose you may take 2 capsules within the first hour ● do not exceed 2 capsules in any 8- to 12-hour period ● do not exceed 3 capsules in a 24-hour period</p> <p>children under 12 years: ● ask a doctor</p> <p>Other information ● each capsule contains sodium 20 mg ● store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F). ● protect from light ● read all directions and warnings before use.</p> <p>Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol</p> <p>Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)</p>

BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	9pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	8pt		

Horizontal Scale: 85% Average Kerning: 0%

BIONPHARMA

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	10pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2.431pt
Body Text:	6pt	Hairlines:	.486pt
Drug Facts (continued):	9pt		

Horizontal Scale: 94%

Average Kerning: 0%

Be Health

Be Health

NDC 69452-179-78

Naproxen Sodium Capsules, 220 mg (NSAID)

Do not use if printed safety seal under cap is torn or missing.



80 Capsules**

**Liquid-filled capsule

†Compare to the active ingredient in ALEVE® Liquid Gels

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain reliever/fever reducer

Strength to last 12 hours

***This product is not manufactured or distributed by the owners of ALEVE® Liquid Gels.

Distributed by:

BIONPHARMA
600 Alexander Road,
Princeton, NJ 08540



N 3 69452 17978 9

List No.:
Exp. Date:

NDC 69452-179-78

Be Health

†Compare to the active ingredient in ALEVE® Liquid Gels

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain reliever/fever reducer

Strength to last 12 hours

80 Capsules**

**Liquid-filled capsule

L0000473
R0621

Drug Facts (continued)

When using this product:

- take with food or milk, if stomach upset occurs

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

● do not take more than directed

- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

● take 1 capsule every 8 to 12 hours while symptoms last

- for the first dose, you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

children under 12 years: ● ask a doctor

Other Information

● each capsule contains: sodium 20 mg

- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- protect from light
- avoid alcohol and warfarin before use.

Inactive Ingredients

FD&C Blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?

1-888-235-2465 (Mon-Fri 9AM-5PM EST)

Drug Facts

Active ingredient (in each capsule)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*, a nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- menstrual cramps
- the common cold
- temporarily reduces fever

- backache
- muscular aches
- headache
- toothache

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

● if you have ever had an allergic reaction to any other pain reliever/fever reducer

- right before or after heart surgery

Ask a doctor before use if

● stomach bleeding warning applies to you


- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are


● under a doctor's care for any serious condition

- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

Front



Back



BIONPHARMA	
Labeling Format Information:	
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique	
Drug Facts:	9pt
Header:	8pt
Subheader:	6pt
Body Text:	6pt
Drug Facts (continued):	N/A
Leading:	6.12 - 6.5pt
Bullets:	5pt
Barlines:	2pt
Hairlines:	.5pt
Horizontal Scale: 85%	
Average Kerning: 0%	

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
12/13/2021 03:53:13 PM