UNPRINTED AREA FOR
LOT & EXP CODING
(UNVARNISHED)

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Product inside sealed in plastic blister with foil backing. Do Not Use if plastic blister or foil barrier is broken.

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
For most recent product information, visit www.advil.com
Trademarks owned or licensed by GSK ©2021 GSK or licensor

81% For Position Only

3 0573 0199 22 3

Drug Facts

Active ingredients (in each tablet) **Purposes**

Ibuprofen 200 mg (NSAID)* ..... Pain reliever/fever reducer
Phenylephrine HCl 10 mg ..... Nasal decongestant
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms associated with the common cold or flu:
  - headache
  - fever
  - sinus pressure
  - nasal congestion
  - minor body aches and pains

Drug Facts (continued)

- reduces swelling of the nasal passages
- temporarily restores freer breathing through the nose

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age because this product contains too much medication for children under this age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if

- stomach bleeding warning app ies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn

Drug Facts (continued)

- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other product that contains phenylephrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because buprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the fo lowing signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days

000065893

Drug Facts (continued)

Other information

- store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
- read all warnings and directions before use. Keep carton.

Inactive ingredients acesulfame potassium, artificial flavor, carnauba wax, colloidal si icon dioxide, corn starch, croscarmellose sodium, glycerin, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, sodium lauryl sulfate, stearic acid, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

Questions or comments? Call weekdays from **9 AM to 5 PM EST** toll free at **1-800-88-ADVIL**

Drug Facts (continued)

- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- **do not take more than directed**
- adults and children 12 years of age and over:
  - take 1 tablet every 4 hours while symptoms persist.
  - do not use more than 6 tablets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use because this product contains too much medication for children under this age

NON-DROWSY

Advil

50% MORE
10 FREE Tablets

SINUS CONGESTION & PAIN

Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg Nasal Decongestant

- ✓ Nasal Congestion
- ✓ Sinus Pressure
- ✓ Nasal Swelling
- ✓ Headache
- ✓ Fever
- ✓ Body Aches

I200-P10

ONE TABLET DOSE!

30 ~~20~~ Coated Tablets



Reference ID: 4891800



UNVARNISHED AREA
FOR LOT # AND EXP. DATE

Drug Facts

Active Ingredients (in each tablet)	Purposes
Ibuprofen 200 mg (NSAID)*	Pain reliever/fever reducer
Phenylephrine HCl 10 mg	Nasal decongestant

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms associated with the common cold or flu:
  - headache ■ fever
  - sinus pressure ■ nasal congestion ■ minor body aches and pains
- reduces swelling of the nasal passages
- temporarily restores freer breathing through the nose

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives ■ facial swelling ■ asthma (wheezing) ■ shock
- skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older ■ have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age because this product contains too much medication for children under this age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other product that contains phenylephrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- adults and children 12 years of age and over:
  - take 1 tablet every 4 hours while symptoms persist.
  - do not use more than 6 tablets in any 24-hour period
- children under 12 years of age: do not use because this product contains too much medication for children under this age

Other information

- store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
- read all warnings and directions before use

Inactive ingredients

acesulfame potassium, artificial flavor, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, sodium lauryl sulfate, stearic acid, sucralose, synthetic iron oxide, talc, titanium dioxide, tragacanth, xanthan gum

Questions or comments?

Call weekdays from **9 AM to 5 PM EST** toll free at **1-800-88-ADVIL**

Do Not Use if pouch is broken or torn

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
For most recent product information, visit www.advil.com
Trademarks owned or licensed by GSK ©2021 GSK or licensor

Drug Facts (continued)

When using this product
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - weakness in one part or side of body
  - trouble breathing
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely

NON-DROWSY

Advil

SINUS CONGESTION & PAIN

Ibuprofen 200 mg Pain Reliever/Fever Reducer **(NSAID)**
Phenylephrine HCl 10 mg Nasal Decongestant

- ✓ Nasal Congestion
- ✓ Sinus Pressure
- ✓ Nasal Swelling
- ✓ Headache
- ✓ Fever
- ✓ Body Aches

I200-P10

ONE TABLET DOSE!

50 Packets of 1 Coated Tablet

NON-DROWSY
Advil
SINUS CONGESTION & PAIN
Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg Nasal Decongestant
✓ Nasal Congestion ✓ Headache
✓ Sinus Pressure ✓ Fever
✓ Nasal Swelling ✓ Body Aches
I200-P10
ONE TABLET DOSE!
50 Packets of 1 Coated Tablet

NON-DROWSY
Advil
SINUS CONGESTION & PAIN
Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg Nasal Decongestant
✓ Nasal Congestion
✓ Sinus Pressure
✓ Nasal Swelling
✓ Headache
✓ Fever
✓ Body Aches
I200-P10
ONE TABLET DOSE!
50 Packets of 1 Coated Tablet

NON-DROWSY
Advil
SINUS CONGESTION & PAIN
Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg Nasal Decongestant
✓ Nasal Congestion
✓ Sinus Pressure
✓ Nasal Swelling
✓ Headache
✓ Fever
✓ Body Aches
I200-P10
ONE TABLET DOSE!
50 Packets of 1 Coated Tablet

000065888

FPO
3 0573 0199 01 8

Reference ID: 4891800

Advil SINUS CONGESTION & PAIN

Ibuprofen, 200 mg/Phenylephrine HCl, 10 mg
Pain Reliever/Fever Reducer (NSAID)*/Nasal Decongestant
1 Tablet Non-Drowsy ***nonsteroidal anti-inflammatory drug**

Uses temporarily relieves these symptoms associated with the common cold or flu: headache, fever, sinus pressure, nasal congestion, minor body aches and pains; reduces swe ling of nasal passages; temporarily restores freer brea hing through the nose.
Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSA D, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others], have 3 or more alcoho ic drinks every day while using this product, take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use in children under 12 years of age because this product contains too much medication for ch ldren under this age, if you have ever had an a lergic reaction to any other pain re lever/fever reducer, right before or after heart surgery, f you are now taking a prescription monoamine oxidase inhibitor (MAOI), (certain drugs for depression, psychiatric, or emotional cond tions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. **Ask a doctor before use if** stomach bleeding warning applies to you; you have

TO OPEN: FOLD BACK ON DOTTED LINE. TEAR ON ARROW OR USE SCISSORS. Do Not Use if pouch is broken or torn.

FRONT

problems or serious side effects from taking pain relievers or fever reducers; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke; you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition, taking any other product that contains phenylephrine or any other nasal decongestant, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin or taking any other drug. **When using this product** take w th food or m lk f stomach upset occurs. **Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools or have stomach pain that does not get better; you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swe ling; pain gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; nasal congestion lasts for more than 7 days; symptoms continue or get worse; redness or swe ling is present in the painful area; you get nervous, dizzy, or sleepless; any new symptoms appear.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless defin tely directed to do so by a doctor because t may cause problems in the unborn child or comp ications during delivery. **Keep out of reach of ch ldren.** In case of overdose, get medical help or contact a Poison Control Center right away. ***Directions*** **Do not take more than directed.** Adults and ch ldren 12 years of age and over: take 1 tablet every 4 hours while symptoms persist. Do not use more than 6 tablets in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use because his product contains too much medication for children under this age. ***Other information*** Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). Read all warnings and directions before use. ***Inactive ingredients*** acesulfame potassium, artificial flavor, carnauba wax, colloidal si icon dioxide, corn starch, croscarmellose sodium, glycerin, hyprome lose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, sodium lauryl sulfate, stearic acid, sucralose, synthetic iron oxide, talc, itanium dioxide, triacetin, xanthan gum
☎ toll free at **1-800-88-ADVIL.** For most recent product information, visit www.advil.com
Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor 000065880

BACK

LASER ENCODING AREA FOR LOT AND EXPIRATION INFORMATION. PRINTED IN ONE SOLID COLOR, BLACK OR DARK BLUE.

problems or serious side effects from taking pain relievers or fever reducers; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke; you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition, taking any other product that contains phenylephrine or any other nasal decongestant, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin or taking any other drug. **When using this product** take w th food or m lk f stomach upset occurs. **Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools or have stomach pain that does not get better; you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swe ling; pain gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; nasal congestion lasts for more than 7 days; symptoms continue or get worse; redness or swe ling is present in the painful area; you get nervous, dizzy, or sleepless; any new symptoms appear.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless defin tely directed to do so by a doctor because t may cause problems in the unborn child or comp ications during delivery. **Keep out of reach of ch ldren.** In case of overdose, get medical help or contact a Poison Control Center right away. ***Directions*** **Do not take more than directed.** Adults and ch ldren 12 years of age and over: take 1 tablet every 4 hours while symptoms persist. Do not use more than 6 tablets in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use because his product contains too much medication for children under this age. ***Other information*** Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). Read all warnings and directions before use. ***Inactive ingredients*** acesulfame potassium, artificial flavor, carnauba wax, colloidal si icon dioxide, corn starch, croscarmellose sodium, glycerin, hyprome lose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, sodium lauryl sulfate, stearic acid, sucralose, synthetic iron oxide, talc, itanium dioxide, triacetin, xanthan gum
☎ toll free at **1-800-88-ADVIL.** For most recent product information, visit www.advil.com
Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor 000065880

BACK

[redacted]

Reference ID: 4891800

AREA FOR LOT AND EXP
TO BE PRINTED IN DARK COLOR

you have problems or serious side effects from taking pain re ievers or fever reducers; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke; you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition, taking any other product that contains phenylephrine or any other nasal decongestant, taking aspirin for heart attack or stroke, because ibuprofen may decrease his benefit of aspirin or taking any other drug. **When using this product** take w th food or milk if stomach upset occurs. **Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding: feel faint, vom t blood, have bloody or black stools or have stomach pain that does not get better; you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swe ling; pain gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; nasal congestion lasts for more than 7 days; symptoms continue or get worse; redness or swel ing is present in the painful area; you get nervous, dizzy, or sleepless; any new symptoms appear. **If pregnant or breast-feeding,** ask a hea th professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ***Directions*** **Do not take more than directed.** Adults and children 12 years of age and over: take 1 tablet every 4 hours while symptoms persist. Do not use more than 6 tablets in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use because this product contains too much medication for children under this age. ***Other information*** Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). Read a l warnings and directions before use. ***Inactive ingredients*** acesulfame potassium, artificial flavor, carnauba wax, colloidal silicon dioxide, corn starch, croscarme lose sodium, glycerin, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregela inized starch, propyl gallate, sodium lauryl sulfate, stearic acid, sucralose, synthetic iron oxide, talc, titanium dioxide, triace in, xanthan gum

☎ toll free at **1-800-88-ADVIL** For most recent product information, visit www.advil.com GSK CH, Warren, NJ 07059

00006589 1

BACK

PANEL 1

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

LIFT HERE
For More Drug Facts

Drug Facts

Active ingredients (in each tablet) — **Purposes**

Ibuprofen 200 mg (NSAID)* ..... Pain reliever/fever reducer
Phenylephrine HCl 10 mg ..... Nasal decongestant
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms associated with the common cold or flu:
  - headache
  - fever
  - sinus pressure
  - nasal congestion
  - minor body aches and pains
- reduces swelling of the nasal passages
- temporarily restores freer breathing through the nose

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Drug Facts (continued)

Symptoms may include:
- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.
The chance is higher if you:
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

000065890

PANEL 2

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
- in children under 12 years of age because this product contains too much medication for children under this age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are
- under a doctor's care for any serious condition
- taking any other product that contains phenylephrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse

BASE PANEL

Drug Facts (continued)

- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- adults and children 12 years of age and over:
  - take 1 tablet every 4 hours while symptoms persist.
  - do not use more than 6 tablets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use because this product contains too much medication for children under this age

Other information

- store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
- read all warnings and directions before use. Keep carton.

Drug Facts (continued)

Inactive ingredients acesulfame potassium, artificial flavor, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, sodium lauryl sulfate, stearic acid, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

Questions or comments? Call weekdays from **9 AM to 5 PM EST** toll free at **1-800-88-ADVIL**

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

LIFT HERE

Drug Facts

Active ingredients (in each tablet) **Purposes**

Ibuprofen 200 mg (NSAID)* ..... Pain reliever/fever reducer

Phenylephrine HCl 10 mg ..... Nasal decongestant

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms associated with the common cold or flu:
  - headache
  - fever
  - sinus pressure
  - nasal congestion
  - minor body aches and pains
- reduces swelling of the nasal passages
- temporarily restores freer breathing through the nose

00006587 9

PANEL 1

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age because this product contains too much medication for children under this age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

PANEL 2

Drug Facts (continued)

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating

Drug Facts (continued)

due to an enlarged prostate gland, or had a stroke

- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other product that contains phenylephrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood

PANEL 3

Drug Facts (continued)

- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- adults and children 12 years of age and over:
  - take 1 tablet every 4 hours while symptoms persist.
  - do not use more than 6 tablets in any 24-hour period unless directed by a doctor

PANEL 4

Drug Facts (continued)

- children under 12 years of age: do not use because this product contains too much medication for children under this age

Other information

- store at 20 -25ºC (68 -77ºF). Avoid excessive heat above 40ºC (104ºF).
- read all warnings and directions before use. Keep carton.

Inactive ingredients

acesulfame potassium, artificial flavor, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, sodium lauryl

Drug Facts (continued)

sulfate, stearic acid, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

Questions or comments?

Call weekdays from **9 AM to 5 PM EST** toll free at **1-800-88-ADVIL**

BASE PANEL

TO SHOW CONSTRUCTION ONLY



[redacted]

Reference ID: 4891800

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
11/19/2021 03:13:09 PM