



Do not use if pouch is torn or damaged.

Active ingredient (in each tablet)

Famotidine 20 mg..... Purpose
Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach

• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

To Open: While Folded on Line, Tear At Slit



Warnings: Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. **Do not use** • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers. **Ask a doctor before use if you have** • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting stomach pain • kidney disease. **Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of each of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions** • adults and children 12 years and over: • to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew. • to **prevent** symptoms swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor. **Other information** • read the directions and warnings before use • keep the carton. It contains important information. • store at 20°-25°C (68°-77°F) • protect from moisture. **Inactive ingredients** carnauba wax, hydroxypropyl cellulose, hypromellose magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide **Questions or comments? 1-800-755-4008** (toll-free) or **215-273-8755** (collect)

Famotidine (active ingredient) made in Hungary

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**

McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA

©McNeil Consumer Pharmaceuticals Co. 2021

Pepcid® and Pepcid AC® are registered trademarks of McNeil CPC

30050385



Heartburn Relief & Prevention

on-the GO

Individually wrapped tablets perfect for:



NDC 16837-855-02

MAXIMUM STRENGTH

Pepcid[®] AC

Famotidine Tablets 20 mg Acid Reducer

Just One Tablet!
Prevents & Relieves Heartburn Due to Acid Indigestion



actual size

2 Tablets



MAXIMUM STRENGTH

Pepcid[®] AC

Famotidine (active ingredient) made in Hungary

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 2021
Pepcid[®] and Pepcid AC[®] are registered trademarks of McNeil CPC

30050389

Important: Read all product information before using. Keep this box for important information.

Drug Facts (continued)

Stop use and ask a doctor if:
• Your heartburn continues or worsens
• You need to take this product for more than 14 days

Drug Facts

Active ingredient
Famotidine 20 mg Acid reducer

Purpose
(In each tablet)

Uses
• Relieves heartburn associated with acid indigestion and sour stomach
• Prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers
• If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• With other acid reducers
• Ask a doctor before use if you have had heartburn over 3 months. This may be a sign of a more serious condition.
• Heartburn with **lightheadedness, sweating, or dizziness**
• Chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
• Frequent wheezing, particularly with heartburn
• Explained weight loss
• Nausea or vomiting
• Stomach pain
• Kidney disease

Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• With other acid reducers
• Ask a doctor before use if you have had heartburn over 3 months. This may be a sign of a more serious condition.
• Heartburn with **lightheadedness, sweating, or dizziness**
• Chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
• Frequent wheezing, particularly with heartburn
• Explained weight loss
• Nausea or vomiting
• Stomach pain
• Kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug, Acid reducers may interact with certain prescription drugs.

Questions or comments?
1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, fatty, spicy, or acidic foods, citrus fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



30050389

30050389

EXP

50 Pouches of 20 mg each
Just One Tablet!
 Prevents & Relieves Heartburn
 Due to Acid Indigestion
Famotidine Tablets 20 mg
Acid Reducer
Pepcid[®] AC
MAXIMUM STRENGTH

MAXIMUM STRENGTH
Pepcid[®] AC

Drug Facts	Purpose
Active ingredient (in each tablet) Famotidine 20 mg	Acid reducer
Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages	
Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness Frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease	
Drug Facts (continued) Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)	
Directions • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor	
Other information • read the directions and warnings before use • keep the carton. It contains important information. • store at 20°-25°C (68°-77°F) • protect from moisture • do not use if pouch is torn or damaged	
Inactive ingredients carnauba wax, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide	
Questions or comments? 1-800-755-4008 (toll-free) or 215-273-8755 (collect)	

Famotidine (active ingredient) made in Hungary
 Distributed by: JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©McNeil Consumer Pharmaceuticals Co. 2021
 Pepcid[®] and Pepcid AC[®] are registered trademarks of McNeil CPC.

7 16837 85572 9

MAXIMUM STRENGTH
Pepcid[®] AC
Famotidine Tablets 20 mg
Acid Reducer
Just One Tablet!
 Prevents & Relieves Heartburn
 Due to Acid Indigestion

50 Pouches of 1 Tablet per Pouch
20 mg each
(Individual Pouches)

NDC 16837-855-52
MAXIMUM STRENGTH
Pepcid[®] AC
Famotidine Tablets 20 mg
Acid Reducer
Just One Tablet!
 Prevents & Relieves Heartburn
 Due to Acid Indigestion

50 Pouches of 1 Tablet per Pouch
20 mg each
(Individual Pouches)

MAXIMUM STRENGTH
Pepcid[®] AC
Famotidine Tablets 20 mg
Acid Reducer
Just One Tablet!
 Prevents & Relieves Heartburn
 Due to Acid Indigestion

50 Pouches of 1 Tablet per Pouch
20 mg each
(Individual Pouches)

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
02/18/2022 12:46:41 PM