



This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfd>

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.  
*Active ingredient (in each tablet)*

Famotidine 20 mg. . . . . *Purpose*  
Acid reducers

*Uses* • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages. **Warnings:** A large alert: Do not use if you are a leptic to form, lime or other acid

reducers. Do not use

• If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • Tell other acid reducers. Ask a doctor before use if you have • had heartburn over 3 months.

30047631

**MAXIMUM STRENGTH**

NDC 16837-855-75

**Pepcid<sup>®</sup>**  
**AC**

Famotidine Tablets 20 mg Acid Reducer

**Just One Tablet!**

Prevents & Relieves Heartburn  
Due to Acid Indigestion

**75 Tablets**

This may be a sign of a more serious condition. • heartburn with light-headedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulder; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease. **Ask a doctor or pharmacist before use if you are taking a prescription drug.**

Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet the glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor

Famotidine (active ingredient) made in Hungary  
Distributed by: JOHNSON & JOHNSON CONSUMER PHARMACEUTICALS, A Division of  
Fort Washington, PA 19101 USA ©McNeil Consumer Pharmaceuticals Co. 2020  
PEPCID<sup>®</sup> and PEPCID AC<sup>®</sup> are registered trademarks of McNeil-PPC

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

**125 TABLETS (75ct + 50ct)**

**MAXIMUM STRENGTH**

NDC 16837-855-00

**Pepcid<sup>®</sup>**  
**AC**

**Famotidine Tablets 20 mg  
Acid Reducer**

**Just One Tablet!**

*Prevents & Relieves Heartburn  
Due to Acid Indigestion*



actual size

**75 Tablets**



**MAXIMUM STRENGTH**

**Pepcid<sup>®</sup>**  
**AC**

**Famotidine Tablets 20 mg  
Acid Reducer**

**Just One Tablet!**

*Prevents & Relieves Heartburn  
Due to Acid Indigestion*



actual size


**50 Tablets**



© J&JCI 2021

FPO

MCN-TBD-XXXX



7

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
12/20/2022 12:42:38 PM