

Location for lot number and expiration date

*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.

Active ingredient (in each tablet).....**Purpose**

Use ■ Treats frequent heartburn (occurs 2 or more days a week)
■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
■ Allergic reaction occurs, stop use and seek medical help right away.
■ Do not use if you have:
■ heartburn with swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
■ frequent chest pain
■ signs of a serious condition. See your doctor.

These may be signs of a serious condition. See your doctor.
■ liver disease
■ heartburn over or you have
■ heartburn with certain prescription drugs.
■ Stop use and ask doctor if:
■ you need to take this product for more than 14 days
■ you develop a rash or joint pain
■ you get diarrhea

If pregnant or breast-feeding, ask a health care professional before use.
■ You need to breast-feed your child. In days of overuse, stop medical help or contact a Poison Control Center right away. (1-800-222-2222.)

**KEEP THE CARTON. IT CONTAINS
IMPORTANT INFORMATION.**



Age Group	Percentage
18-24	10%
25-34	15%
35-44	20%
45-54	25%
55-64	30%
65-74	35%
75-84	40%
85-94	45%
95-104	50%

Dissolve Tabs



Strawberry
Flavor

One 14-day course of treatment
May take 1 to 4 days for full effect

Actual Size

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.

- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight. ■ If you smoke, quit smoking.

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Reference ID: 5070463




FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the “Drug Facts” box

**Location for
lot number and
expiration date**

Reference ID: 5070463

Reference ID: 5070463


Reference ID: 5070463



Reference ID: 5070463

Reference ID: 5070463

Reference ID: 5070463




FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the “Drug Facts” box



8U155 FA C8-B 100422



Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Do not eat late at night or just before bedtime.
- Do not eat slowly and do not eat big meals.

© 2007 Takeda Pharmaceuticals U.S.A., Inc.

This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc. The registered trademark Prevacid®.

Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Made in Israel

Drug Facts

Active ingredient (in each tablet)

Use

■ Treats frequent heartburn (occurs **2 or more** days a week)

■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- Do not use if you are allergic to lansoprazole
- Skin rash
- Stomach pain
- Diarrhea
- Rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- Trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- Chest pain or shoulder pain with shortness of breath, sweating, pain
- Spreading to arms, neck or shoulders, or lightheadedness

Frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask your doctor before use if you have

- Liver disease
- Heartburn over 3 months. This may be a sign of a more serious condition.
- Frequent wheezing, particularly with heartburn
- Unexplained weight loss
- Nausea or vomiting
- Stomach pain

Ask your doctor or pharmacist before use if you are taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- Your heartburn continues or worsens
- You need to take this product for more than 14 days
- You need to take more than 1 course of treatment every 4 months
- You get diarrhea
- You develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Purpose

Acid Reducer

Drug Facts (continued)

Directions • adults 18 years of age and older
This product is to be used once a day (every 24 hours), every day for 14 days
• I may take 1 to 3 days for full effect (although some people get complete relief of symptoms within 24 hours)

14-Day Course of Treatment

- Take 1 tablet before eating in the morning
- Do not eat crust or chew tablets
- Place the tablet on tongue; tablet disintegrates, with or without water.
- Do not take more than 1 tablet a day
- Take every day for 14 days
- Do not take more than 1 tablet a day
- Do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed)

- You may repeat a 14-day course every 4 months
- Do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Children under 18 years of age: ask a doctor before use. Headturn in children may sometimes be caused by a serious condition.

Other information

- Read the directions and warnings before use
- Keep the carton. It contains important information.
- Store at 20-25°C (68-77°F) • Keep product out of high heat and humidity
- Take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscapellone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, sodium chloride, sodium stearoyl fumarate, polyoxylated 80 polyethylene glycol, polyoxylated 40 polyethylene glycol, polyoxylated 20 polyethylene glycol, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg

Lansoprazole

**Delayed Release
Orally Disintegrating Tablets 15 mg**

Safety Feature – Do not use if printed tablet blister unit is open or torn.

**KEEP THE CARTON. IT CONTAINS
IMPORTANT INFORMATION.**



*Compare to the active ingredient
in Prevacid® 24 HR**

Value 3 Pack



Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg
Acid Reducer

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



**Strawberry
Flavor**

Actual
size

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the “Drug Facts” box



FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



8U155 FA C8-N 100422



Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Do not eat late at night or just before bedtime.
- Do not eat slowly and do not eat big meals.

© 2007 Takeda Pharmaceuticals U.S.A., Inc.

This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc. The registered trademark Prevacid®.

Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Made in Israel

Drug Facts

Active ingredient (in each tablet)

Use

■ Treats frequent heartburn (occurs **2 or more** days a week)

■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- Do not use if you are allergic to lansoprazole
- Skin rash
- Stomach pain
- Diarrhea
- Rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- Trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- Chest pain or shoulder pain with shortness of breath, sweating, pain
- Spreading to arms, neck or shoulders, or lightheadedness

Frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask your doctor before use if you have

- Liver disease
- Heartburn over 3 months. This may be a sign of a more serious condition.
- Frequent wheezing, particularly with heartburn
- Unexplained weight loss
- Nausea or vomiting
- Stomach pain

Ask your doctor or pharmacist before use if you are taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- Your heartburn continues or worsens
- You need to take this product for more than 14 days
- You need to take more than 1 course of treatment every 4 months
- You get diarrhea
- You develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Purpose

Acid Reducer

Drug Facts (continued)

Directions • adults 18 years of age and older
This product is to be used once a day (every 24 hours), every day for 14 days
• I may take 1 to 3 days for full effect (though some people get complete relief of symptoms within 24 hours)

14-Day Course of Treatment

• Take 1 tablet before eating in the morning
• Do not eat crust or chew tablets
• Place the tablet on tongue; tablet disintegrates, with or without water.
• Do not take more than 1 tablet a day
• Take every day for 14 days
• Do not take more than 1 tablet a day
• Do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed)

• You may repeat a 14-day course every 4 months
• Do not take for more than 14 days or more often than every 4 months
• Unless directed by a doctor

• Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

• Read the directions and warnings before use
• Keep the carton. It contains important information.
• Store at 20-25°C (68-77°F) • Keep product out of high heat and humidity
• Protect product from moisture
• Take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscapellone, flavor, hypromellose, hypromellose phthalate, maize polyphosphate, glycol, silicon dioxide, sodium stearoyl fumarate, polydextrose, mannitol, polyethylene glycol, polyethylene glycol succinate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg



Lansoprazole

**Delayed Release
Orally Disintegrating Tablets 15 mg**

Safety Feature – Do not use if printed tablet blister unit is open or torn.

**KEEP THE CARTON. IT CONTAINS
IMPORTANT INFORMATION.**



NDC 64239-181-03

Treats Frequent Heartburn



Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg
Acid Reducer

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water




Strawberry Flavor



FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the “Drug Facts” box



8U155 FA C8-P 100422



Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Do not eat late at night or just before bedtime.
- Do not eat slowly and do not eat big meals.

© 2007 Takeda Pharmaceuticals U.S.A., Inc.

This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc. The registered trademark Prevacid®.

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Made in Israel

[illegible]

Drug Facts (continued)

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- do not eat or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water.
- the tablet can also be swallowed whole with water.
- take every day for 14 days ■ do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton, it contains important information.
- store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
- protect product from moisture
- take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscapellon, flavor, hypromellose, hypromellose phthalate, maize polyborate, glycol, polyethylene glycol, silicon dioxide, sodium stearoyl fumarate, polysorbate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg

Lansoprazole

**Delayed Release
Orally Disintegrating Tablets 15 mg**

Safety Feature – Do not use if printed tablet blister unit is open or torn.

**KEEP THE CARTON. IT CONTAINS
IMPORTANT INFORMATION.**



*Compare to the active ingredient
in Prevacid® 24 HR**

Bonus Size! 14 Free



Lansoprazole

**Delayed Release
Orally Disintegrating Tablets 15 mg
Acid Reducer**

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Strawberry Flavor




Actual
size

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



8U155 FA C8-S 100422



Tips for Managing Heartburn

- Many foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and
- fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- If you are overweight, lose weight. ■ If you smoke, quit smoking.
- Wear loose-fitting clothing around your stomach.
- Raise the head of your bed.
- Do not lie flat or bend over soon after eating.

© 2007, Janssen Pharmaceutica, Inc. All rights reserved. This product is not manufactured or distributed by Janssen Pharmaceutica U.S.A., Inc., owner of the registered trademark Prevacid®.

Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

Made in Israel

Drug Facts

Active ingredient (in each tablet) lansoprazole 15 mg.

Purpose Acid Reducer

Use

- Treats frequent heartburn (occurs **2 or more** days a week)
- Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- Do not use if you are allergic to lansoprazole
- Lansoprazole may cause severe skin reactions. Symptoms may include:
 - rash
 - blisters
 - hives
- If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask your doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions • adults 18 years of age and older
This product is to be used once a day (every 24 hours), every day for 14 days
• I may take 1 to 3 days for full effect (though some people get complete relief of symptoms within 24 hours)

14-Day Course of Treatment

- Take 1 tablet before eating in the morning
- Do not eat crust or chew tablets
- Place the tablet on tongue; tablet disintegrates, with or without water.
- Do not take more than 1 tablet a day
- Take every day for 14 days
- Do not take more than 1 tablet a day
- Do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed)

- You may repeat a 14-day course every 4 months
- Do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Children under 18 years of age: ask a doctor before use. Headturn in children may sometimes be caused by a serious condition.

Other information


- Read the directions and warnings before use
- Keep the carton. It contains important information.
- Store at 20-25°C (68-77°F)
- Protect product from moisture
- Take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscapellone, flavor, hypromellose, hypromellose phthalate, maize polyphosphate, 80 polyphosphite glycol, silicon dioxide, sodium stearoyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg



Lansoprazole

**Delayed Release
Orally Disintegrating Tablets 15 mg**

Safety Feature – Do not use if printed tablet blister unit is open or torn.

**KEEP THE CARTON. IT CONTAINS
IMPORTANT INFORMATION.**



NDC 64239-181-03

Treats Frequent Heartburn

MELTech™
Melts In Your Mouth

**Compare to the active ingredient
in Prevacid® 24 HR***

Bonus! 14 Tablets FREE

 **24 HR**

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg
Acid Reducer

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Strawberry Flavor



Actual
size

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
11/01/2022 01:46:06 PM