

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use SPY AGENT™ GREEN safely and effectively. See full prescribing information for SPY AGENT™ GREEN.

SPY AGENT™ GREEN (indocyanine green for injection), for intravenous or interstitial use.
Initial U.S. Approval: 1959

INDICATIONS AND USAGE

SPY AGENT™ GREEN is an optical imaging agent indicated for use with a fluorescence imaging device for:

- Visualization of vessels (micro and macro vasculature), blood flow and tissue perfusion before, during, and after vascular, gastrointestinal, organ transplant, and plastic, micro- and reconstructive surgeries including general minimally invasive surgical procedures in adults and pediatric patients one month of age and older. (1.1)
- Visualization of extrahepatic biliary ducts in adults and pediatric patients aged 12 to 17 years. (1.2)
- Visualization of lymph nodes and lymphatic vessels during lymphatic mapping in women with cervical and uterine tumors. (1.3)

DOSAGE AND ADMINISTRATION

Visualization of Vessels, Blood Flow and Tissue Perfusion (2.5 mg/mL solution):

- The recommended single intravenous dose in adults and pediatric patients one month and older for a surgical procedure is 1.25 mg to 5 mg.
- The recommended single intravenous dose in adults for visualization of perfusion in extremities through the skin for plastic, micro, and reconstructive surgeries is 3.75 mg to 10 mg.

Additional doses may be administered to obtain additional imaging sequences during the procedure, however, do not exceed a total dose of 2 mg/kg. (2.1)

Visualization of Extrahepatic Biliary Ducts (2.5 mg/mL solution):

- The recommended single intravenous dose in adults and pediatric patients aged 12 to 17 years is 2.5 mg SPY AGENT GREEN.
- Inject SPY AGENT GREEN approximately 45 minutes prior to surgery to allow indocyanine green to collect in the biliary anatomy.

Additional doses may be administered to obtain additional imaging sequences during the procedure, however, do not exceed a total dose of 2 mg/kg. (2.2)

Visualization of Lymph Nodes and Lymphatic Vessels During Lymphatic Mapping for cervical and uterine tumors (1.25 mg/mL solution):

- The recommended single interstitial dose in women is four 1.25 mg injections for a total dose of 5 mg. Inject at the 3 and 9 o'clock positions of the cervix with a superficial (1 to 3 mm) and deep (1 to 3 cm) injection at each position. (2.3)

See Full Prescribing Information for instructions on reconstitution of the lyophilized powder, and preparation and administration of injections. (2.4).

DOSAGE FORMS AND STRENGTHS

For injection: SPY AGENT GREEN is a sterile lyophilized green powder containing 25 mg indocyanine green in a single-patient use vial for reconstitution. (3)

CONTRAINDICATIONS

History of hypersensitivity to indocyanine green. (4)

WARNINGS AND PRECAUTIONS

- Hypersensitivity reactions: Hypersensitivity reactions including anaphylaxis and urticaria have occurred with administration of indocyanine green. Always have cardiopulmonary resuscitation personnel and equipment readily available and monitor all patients for hypersensitivity reactions. (5.1)
- Interference with Radioactive Iodine Uptake Studies: Do not perform radioactive iodine uptake studies for at least a week following the use of SPY AGENT GREEN (5.2)

To report SUSPECTED ADVERSE REACTIONS, contact Stryker Corp. at 1-800-624-4422 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Revised: 03/2022

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE**
 - 1.1 Visualization of Vessels, Blood Flow, and Tissue Perfusion
 - 1.2 Visualization of Extrahepatic Biliary Ducts
 - 1.3 Visualization of Lymph Nodes and Lymphatic Vessels During Lymphatic Mapping for Cervical and Uterine Tumors
- 2 DOSAGE AND ADMINISTRATION**
 - 2.1 Recommended Dose and Administration for Visualization of Vessels, Blood Flow, and Tissue Perfusion
 - 2.2 Recommended Dose and Administration for Visualization of Extrahepatic Biliary Ducts
 - 2.3 Recommended Dose and Administration for Visualization of Lymph Nodes and Lymphatic Vessels During Lymphatic Mapping for Cervical and Uterine Tumors
 - 2.4 Reconstitution Instructions
- 3 DOSAGE FORMS AND STRENGTHS**
- 4 CONTRAINDICATIONS**
- 5 WARNINGS AND PRECAUTIONS**
 - 5.1 Hypersensitivity reactions
 - 5.2 Interference with Radioactive Iodine Uptake Studies

- 6 ADVERSE REACTIONS**
 - 7 DRUG INTERACTIONS**
 - 8 USE IN SPECIFIC POPULATIONS**
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 11 DESCRIPTION**
 - 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
 - 14 CLINICAL STUDIES**
 - 16 HOW SUPPLIED/STORAGE AND HANDLING**
 - 17 PATIENT COUNSELING INFORMATION**
- * Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Visualization of Vessels, Blood Flow and Tissue Perfusion

SPY AGENT™ GREEN is indicated in adults and pediatric patients one month of age and older for:

- fluorescence imaging of micro- and macro-vasculature, blood flow and tissue perfusion before, during and after vascular, gastrointestinal, organ transplant, and plastic, micro- and reconstructive surgeries, including general minimally invasive surgical procedures.

1.2 Visualization of Extrahepatic Biliary Ducts

SPY AGENT™ GREEN is indicated in adults and pediatric patients aged 12 to 17 years for:

- fluorescence imaging of extrahepatic biliary ducts.

1.3 Visualization of Lymph Nodes and Lymphatic Vessels During Lymphatic Mapping for Cervical and Uterine Tumors

SPY AGENT™ GREEN is indicated in women for:

- fluorescence imaging of lymph nodes and delineation of lymphatic vessels in the cervix and uterus during lymphatic mapping in patients with solid tumors for which this procedure is a component of intraoperative management.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dose and Administration for Visualization of Vessels, Blood Flow and Tissue Perfusion

Dosing

Adults:

The recommended dose of SPY AGENT GREEN for a single image sequence is 1.25 mg to 5 mg of a 2.5 mg/mL solution.

For visualization of perfusion in extremities through the skin, the recommended dose is 3.75 mg to 10 mg of a 2.5 mg/mL solution.

Immediately flush with a 10 mL bolus of 0.9% Sodium Chloride, USP.

Pediatric patients:

The recommended dose for a single image sequence is 1.25 mg to 5 mg SPY AGENT GREEN of a 2.5 mg/mL solution. Lower doses may be administered in younger patients and in those with lower body weight. Adjust the amount and type of flush to avoid volume and/or sodium overload.

Additional doses may be administered to obtain imaging sequences during the procedure. Do not exceed the maximum total dose of 2 mg/kg.

Administration

Prior to the imaging procedure, draw up the desired dosage of SPY AGENT GREEN solution into appropriate syringes and prepare a 10 mL syringe of 0.9% Sodium Chloride, USP.

Administer via a central or peripheral venous line using a three-way stopcock attached to an injection port on the infusion line.

Inject the prepared SPY AGENT GREEN into the line as a tight bolus. Immediately switch the access on the stopcock and inject the prepared flush.

Imaging Instructions for Visualization of Vessels, Blood Flow and Tissue Perfusion

SPY AGENT GREEN may be used with either the SPY® Elite or PINPOINT® Fluorescence Imaging Systems or with a FDA cleared or approved imaging device that is specifically intended for fluorescence imaging with indocyanine green.

A fluorescence response should be visible in blood vessels within 5 to 15 seconds after injection.

2.2 Recommended Dose and Administration for Visualization of Extrahepatic Biliary Ducts

Dosing and Administration

The recommended single dose of SPY AGENT GREEN for adults and pediatric patients aged 12 to 17 years is 2.5 mg of a 2.5 mg/mL solution administered intravenously as a single dose at least 45 minutes prior to surgery. Additional doses may be administered to obtain imaging sequences during the procedure.

Do not exceed a total dose of 2 mg/kg.

Imaging Instructions for Visualization of Extrahepatic Biliary Ducts

SPY AGENT GREEN may be used with the PINPOINT® Fluorescence Imaging System or with a FDA cleared or approved imaging device that is specifically intended for fluorescence imaging with indocyanine green.

Fluorescence is visible in the biliary tree within 45 minutes after injection.

2.3 Recommended Dose and Administration for Visualization of Lymph Nodes and Lymphatic Vessels During Lymphatic Mapping for Cervical and Uterine Tumors

Dosing and Administration

The recommended single dose of SPY AGENT GREEN is 5 mg of a 1.25 mg/mL solution (four 1 mL injections administered interstitially into the cervix, at the three o'clock and the nine o'clock positions with a superficial (1 mm to 3 mm) and a deep (1 cm to 3 cm) injection at each position).

Imaging Instructions for Visualization of Lymph Nodes and Lymphatic Vessels During Lymphatic Mapping

SPY AGENT GREEN may be used with the PINPOINT® Fluorescence Imaging System or with a FDA cleared or approved imaging device that is specifically intended for fluorescence imaging with indocyanine green.

Fluorescent lymphatic vessels and lymph nodes should begin to be visible within 1 minute after injection.

2.4 Reconstitution Instructions

General

- Prepare SPY AGENT GREEN under sterile conditions prior to surgery.
- Inspect the reconstituted solution for particulate matter, the reconstituted solution should be a clear, green solution.
- Use the prepared solution within 6 hours.
- Discard any unused product.

Visualization of Vessels, Blood Flow, and Tissue Perfusion and Visualization of Extrahepatic Biliary Ducts

- Dissolve SPY AGENT GREEN with 10 mL Sterile Water for Injection, USP to form a concentration of 2.5 mg indocyanine green/mL.

Visualization of Lymph Nodes and Lymphatic Vessels During Lymphatic Mapping

- Dissolve SPY AGENT GREEN with 20 mL Sterile Water for Injection, USP to form a concentration of 1.25 mg indocyanine green/mL.

3 DOSAGE FORMS AND STRENGTHS

SPY AGENT GREEN (indocyanine green for injection, USP) is a sterile lyophilized green powder containing 25 mg of indocyanine green with no more than 5% sodium iodide provided in a 20 mL single-patient use vial.

4 CONTRAINDICATIONS

SPY AGENT GREEN is contraindicated in patients with a history of hypersensitivity to indocyanine green. Reactions have included anaphylaxis [see *Warnings and Precautions* (5.1)].

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

Hypersensitivity reactions including anaphylaxis, urticaria, and deaths due to anaphylaxis have been reported following intravenous administration of indocyanine green. [see *Adverse Reactions* (6)]. Always have cardiopulmonary resuscitation personnel and equipment readily available and monitor all patients for hypersensitivity reactions.

5.2 Interference with Radioactive Iodine Uptake Studies

Because SPY AGENT GREEN contains iodine, the iodine-binding capacity of thyroid tissue may be reduced for at least one week following administration. Do not perform radioactive iodine uptake studies for at least a week following administration of SPY AGENT GREEN [see *Drug Interactions* (7)].

6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the label:

- Hypersensitivity Reactions [see *Warnings and Precautions* (5.1)].

The following adverse reactions have been identified during post-approval use of indocyanine green. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Immune System Disorders: Anaphylaxis, urticaria.

7 DRUG INTERACTIONS

Effect on Radioactive Iodine Uptake Studies

Because SPY AGENT GREEN contains iodine, the iodine-binding capacity of thyroid tissue may be reduced for at least one week following administration. Do not perform radioactive iodine uptake studies for at least one week following administration of SPY AGENT GREEN [see *Warnings and Precautions* (5.2)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no adequate and well-controlled studies of SPY AGENT GREEN in pregnant women. Available data from a very small number of scientific literature studies with indocyanine green use in pregnant women over several decades have not reported any drug associated risks for major birth defects, miscarriage, or adverse maternal or fetal outcomes. Data from one small study in which indocyanine green was administered intravenously to pregnant women during labor suggest there is no placental transfer of the drug. Animal reproduction studies have not been conducted with indocyanine green.

All pregnancies have a background risk of birth defects, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

8.2 Lactation

Risk Summary

Seventeen cases of indocyanine green use in lactating women have been reported in the scientific literature with no adverse events observed in the breastfed infant. However, there are

no data on the presence of indocyanine green in human milk or the effects on milk production. Therefore, the developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for SPY AGENT GREEN and any potential adverse effects on the breastfed infant from SPY AGENT GREEN or from the underlying maternal condition.

8.4 Pediatric Use

Use of indocyanine green for visualization of vessels, blood flow and tissue perfusion has been established in pediatric patients one month and older. Pediatric use is supported by published data in 49 pediatric patients who received indocyanine green for assessment of blood flow and tissue perfusion in cardiovascular, vascular and plastic, micro and reconstructive procedures, and by clinical trials in adults. No overall differences in safety or effectiveness have been observed between pediatric patients and adults. The dose range was similar to the effective dose range in adults [See *Dosage and Administration* (2.1)]. The use of indocyanine green for visualization of vessels, blood flow and tissue perfusion has not been established in pediatric patients less than one month of age.

Use of indocyanine green for visualization of extrahepatic biliary ducts has been established in pediatric patients aged 12 to 17 years. Pediatric use is supported by clinical trials in adults in addition to clinical use in pediatric patients. No overall differences in safety or effectiveness have been observed between pediatric patients and adults. The dose range was similar to the effective dose range in adults [See *Dosage and Administration* (2.2)]. The use of indocyanine green for visualization of extrahepatic biliary ducts has not been established in pediatric patients less 12 years of age.

The safety and efficacy of indocyanine green for visualization of lymph nodes and lymphatic vessels during lymphatic mapping for cervical and uterine tumors has not been established in pediatric patients.

8.5 Geriatric Use

Of the total number of subjects in clinical studies of SPY AGENT GREEN in visualization of vessels, blood flow, and tissue perfusion, 7 percent were 65 and over, while 1 percent were 75 and over and of the total number of subjects in clinical studies of SPY AGENT GREEN in visualization of lymph nodes and lymphatic vessels during lymphatic mapping, 9 percent were 65 and over, while 2 percent were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.


Clinical studies of SPY AGENT GREEN in visualization of extrahepatic biliary ducts did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

11 DESCRIPTION

SPY AGENT GREEN (indocyanine green for injection, USP) is a water soluble, optical imaging agent which is reconstituted with Sterile Water for Injection, USP, for intravenous or interstitial use. Each vial contains a sterile, lyophilized green powder containing 25 mg of indocyanine green with not more than 5% sodium iodide. Hydrochloric acid or sodium hydroxide may be used to adjust the pH prior to lyophilization.

The chemical name for Indocyanine Green is 1 HBenz[e]indolium, 2-[7-[1,3-dihydro-1,1-dimethyl-3-(4-sulfobutyl)-2H-benz[e] indol-2-ylidene]-1,3,5-heptatrienyl]-1,1-dimethyl-3-(4-sulfobutyl)-, hydroxide, inner salt, sodium salt.

Molecular Formula: $C_{43}H_{47}N_2NaO_6S_2$; Molecular Mass: 774.96 g/mol, with the following structural formula:



Indocyanine green has a peak spectral absorption at 800 nm. SPY AGENT GREEN has a pH of 5.5 to 7.5 when reconstituted with Sterile Water for Injection, USP.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

When bound to proteins in plasma or in lymph fluid, indocyanine green absorbs light in the near-infrared region at 805 nm, and emits fluorescence (light) at a slightly longer wavelength, with peak emission at 830 nm. Fluorescence imaging devices provide external energy as near infrared light for ICG to absorb, resulting in excitation of the ICG and the emitted light (fluorescence) is transferred from the field of view to an image on a monitor. These optical properties of indocyanine green are utilized in fluorescence imaging of the micro and macro vasculature, blood flow and tissue perfusion, the extrahepatic biliary ducts, and for lymphatic mapping of lymph nodes and lymphatic vessels.

12.2 Pharmacodynamics

There is no relevant pharmacodynamic data.

12.3 Pharmacokinetics

Distribution

Following intravenous injection, indocyanine green binds to plasma proteins (98%) and is largely confined to the intravascular compartment. Indocyanine green undergoes no significant extrahepatic or enterohepatic circulation; simultaneous arterial and venous blood estimations have shown negligible renal, peripheral, lung or cerebro-spinal uptake of the dye. After biliary obstruction, the dye appears in the hepatic lymph, independently of the bile, suggesting that

the biliary mucosa is sufficiently intact to prevent diffusion of the dye, though allowing diffusion of bilirubin.

Following interstitial injection, indocyanine green is taken up by the lymphatic vessels and lymph nodes.

Elimination

Indocyanine green is taken up from the plasma almost exclusively by the hepatic parenchymal cells and is secreted entirely into the bile.

The plasma fractional disappearance rate at a 0.5 mg/kg dose has been reported to be significantly greater in women than in men, although there was no significant difference in the calculated value for clearance.

14 CLINICAL STUDIES

Visualization of Lymph Nodes and Lymphatic Vessels During Lymphatic Mapping of Cervical and Uterine Tumors

FILM Study (NCT 02209532)

FILM was a randomized, prospective, multi-center, open-label study in patients with early stage uterine or cervical cancers and no known regional nodal or metastatic disease by standard clinical evaluation. SPY AGENT GREEN and a blue dye comparator were injected into the cervix of patients at the beginning of the operative procedure. A total of 176 patients were randomized to receive either SPY AGENT GREEN followed by blue dye or blue dye followed by SPY AGENT GREEN. A total of four 1 mL injections of a 1.25 mg/mL solution of SPY AGENT GREEN were administered into the cervix at the 3 and 9 o'clock positions with a superficial (1 to 3 mm) and a deep (1 to 3 cm) injection at each position for a total dose of 5 mg per patient. Lymphatic mapping was performed intraoperatively using the PINPOINT® Fluorescence Imaging System and standard light, followed by excision of tissues identified by SPY AGENT GREEN, blue dye, or the surgeon's visual and palpation examination. The resected tissues were evaluated by histopathology to confirm presence of lymph nodes. The efficacy of SPY AGENT GREEN and the PINPOINT® Fluorescence Imaging System in the detection of lymphatic vessels and lymph nodes during lymphatic mapping procedures was determined by the number of histology-confirmed lymph nodes detected by SPY AGENT GREEN and/or the blue dye comparator.

[Table 1](#) shows the distribution of resected, confirmed lymph nodes by the presence or absence of SPY AGENT GREEN or blue dye in the modified intent-to-treat population (mITT). Among the confirmed lymph nodes identified, 93% were identified using SPY AGENT GREEN, and 43% were identified using blue dye, a difference of 50% [95% confidence interval 39% to 60%].

Table 1: Distribution of Resected Confirmed Lymph Nodes Detected by SPY AGENT GREEN or Blue Dye (BD)

Analysis population	Nodes (n)	All Lymph Nodes Detected with SPY AGENT GREEN	All Lymph Nodes Detected with BD	Lymph Nodes Detected with SPY AGENT GREEN Only	Lymph Nodes Detected with BD Only	Lymph Nodes Detected with Neither
mITT	513	(476/513) 93%	(220/513) 43%	(262/513) 51%	(6/513) 1%	(31/513) 6%

Table 2 shows the number of patients with at least one resected, confirmed lymph node and the number of patients with at least one bilateral lymph node pair detected by SPY AGENT GREEN or blue dye. With SPY AGENT GREEN, approximately 97% of patients had at least one resected, confirmed lymph node detected and 73% had at least one bilateral lymph node pair detected, compared with 68% and 28%, respectively, with blue dye [p-values for each analysis <0.0001].

Table 2: Distribution of Patients with at Least One Confirmed Unilateral Lymph Node / Bilateral Pair Detected by SPY AGENT GREEN or Blue Dye (BD)

Analysis population	Patients (n)	Patients with All Lymph Nodes Detected with SPY AGENT GREEN	Patients with All Lymph Nodes Detected with BD	Patients with Lymph Nodes Detected with SPY AGENT GREEN only	Patients with Lymph Nodes Detected with BD only	Patients with Lymph Nodes detected with Neither
mITT Unilateral*	172	(167/172) 97%	(118/172) 68%	(51/172) 30%	(2/172) 1%	(3/172) 3%
mITT Bilateral**		(126/172) 73%	(49/172) 28%	(79/172) 46%	(2/172) 1%	(44/172) 26%

*: patients with at least one resected confirmed lymph node detected unilaterally

**: patients with at least one resected confirmed lymph node detected bilaterally

16 HOW SUPPLIED/STORAGE AND HANDLING

NDC 66259-146-01

SPY AGENT GREEN (indocyanine green for injection, USP) is supplied as 25 mg of

lyophilized powder in a 20 mL or 25 mL single patient use vial in the following configurations:

	Components
Spy Elite Kit NDC 66259-306-13 Kit for use with SPY Elite System For visualization of vessels, blood flow, and tissue perfusion	One 25 mg SPY AGENT GREEN (indocyanine green for injection, USP) vial, One 10 mL Sterile Water for Injection, USP plastic vial, One ND8000 sterile drape

SPY Elite Pack NDC 66259-306-23	6 SPY Elite kits: Six 25 mg SPY AGENT GREEN (indocyanine green for injection, USP) vials, Six 10 mL Sterile Water for Injection, USP plastic vials, Six ND8000 sterile drapes
SPY-PHI Kit NDC 66259-903-14 Kit for use with SPY-PHI System For visualization of vessels, blood flow, and tissue perfusion	One 25 mg SPY AGENT GREEN (indocyanine green for injection, USP) vial, One 10 mL Sterile Water for Injection, USP plastic vial, One SPY-PHI (HH2000) sterile drape
SPY-PHI Pack NDC 66259-903-24	6 SPY-PHI kits: Six 25 mg SPY AGENT GREEN (indocyanine green for injection, USP) vials, Six 10 mL Sterile Water for Injection, USP plastic vials, Six SPY-PHI (HH2000) sterile drapes
SPY Minimally Invasive Surgery (SPY-MIS) Kit NDC 66259-936-15 Kit for use with Advanced Imaging Modality (AIM) and PINPOINT Systems For visualization of vessels, blood flow, and tissue perfusion For visualization of extrahepatic biliary ducts	One 25 mg SPY AGENT GREEN (indocyanine green for injection, USP) vial, One 10 mL Sterile Water for Injection, USP plastic vial, Two 3 mL syringes (sterile), Two 10 mL syringes (sterile), Two 18G, 1 inch needles (sterile), Labels for syringes
SPY Minimally Invasive Surgery (SPY-MIS) Pack NDC 66259-936-25	6 SPY-MIS kits: Six 25 mg SPY AGENT GREEN (indocyanine green for injection, USP) vials, Six 10 mL Sterile Water for Injection, USP plastic vials, Twelve 3 mL syringes (sterile), Twelve 10 mL syringes (sterile), Twelve 18G, 1 inch needles (sterile), Labels for syringes
SPY Lymphatics Kit NDC 66259-937-16 Kit for use with the Advanced Imaging Modality (AIM) L10 and L11 Light Sources and 1688 Camera System, and the PINPOINT System For lymphatic mapping	One 25 mg SPY AGENT GREEN (indocyanine green for injection, USP) vial, Two 10 mL Sterile Water for Injection, USP plastic vials, One 10 mL syringe (sterile), One luer-lock 10 mL syringe with controlled handle (sterile), Two spinal needles 22G, 3.5 inch (sterile), Labels for syringes

SPY Lymphatics Pack NDC 66259-937-26	6 SPY Lymphatics kits: Six 25 mg SPY AGENT GREEN (indocyanine green for injection, USP) vials, Twelve 10 mL Sterile Water for Injection, USP plastic vials, Six 10 mL syringes (sterile), Six luer-lock 10 mL syringes with controlled handle (sterile), Twelve spinal needles 22G, 3.5 inch (sterile) Labels for syringes
--	--

NDC 63323-185-10 (or NDC 0409-4887-10) Sterile Water for Injection, USP, 10 mL fill in 10 mL plastic vials.

STORAGE: Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].

PATIENT COUNSELING INFORMATION

Hypersensitivity Reactions

Advise patients to seek medical attention for reactions following injection of SPY AGENT GREEN such as difficulty breathing, swollen tongue or throat, skin reactions including hives, itching and flushed or pale skin, low blood pressure, a weak and rapid pulse and other symptoms or signs of an anaphylactic reaction [see *Warnings and Precautions* ([5.1](#))].

Manufactured by:
Patheon Italia S.p.A.
Ferentino (FR), Italy

Distributed by:
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101,
Burnaby, BC, Canada V5A 4W2

Sterile Water for Injection, USP is manufactured by:
Fresenius Kabi USA, LLC
Grand Island, NY 14072 U.S.A.
or
Hospira, Inc.
Rocky Mount, NC 27804 U.S.A.

LB0016-04

SPYAGENT GREEN NDC 66259-146-01
(indocyanine green for injection, USP)

25 mg/vial

Sterile, single dose vial, discard unused portion
Use within 6 hours of reconstitution.
Usual dosage: See prescribing information

stryker®

Rx Only

Manufactured by: Patheon Italia S.p.A.
Distributed by: Novadaq Technologies ULC



3
5/06/21
05/2021

Lot
Exp.

SPY Elite Pack

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the SPY Elite System
For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-306-23	Use-by Date	DD-MMM-YYYY
REF	Catalogue Number LC3006		
LOT	Batch Code YYMMN		

Manufactured for:
Novadig Technologies U.L.C.
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free
Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].



6625930623



CAUTION



CONSULT
IFU

Rx ONLY

DO NOT REUSE

stryker®

SPY Elite Pack

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the SPY Elite System
For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-306-23	Use-by Date	DD-MMM-YYYY
REF	Catalogue Number LC3006		
LOT	Batch Code YYMMN		

Manufactured for:
Novadig Technologies U.L.C.
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free
Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].



6625930623



CAUTION



CONSULT
IFU

Rx ONLY

DO NOT REUSE

stryker®

SPY Elite Pack

with


SPY AGENT™ GREEN (indocyanine green for injection, USP)

25 mg per vial

Kits for use with the SPY Elite System

For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-306-23		
REF	Catalogue Number LC3006		
LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY
Contents	<div>Components for six (6) SPY Elite Procedures (packaged as six (6) SPY Elite Kits):</div> <ul style="list-style-type: none">• Six 25 mg vials of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, <i>sterile</i>• Six 10 ml vials of Sterile Water for Injection• Six SPY Elite Drapes (ND8000), <i>sterile</i>• Six SPY Elite Kit IFUs• Six SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Inserts		

DO NOT REUSE
Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged.

**Manufactured for:**
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].











6625930623

stryker®

P42573 Rev C

SPY Elite Pack

with



SPY AGENT™ GREEN (indocyanine green for injection, USP)

25 mg per vial

Kits for use with the SPY Elite System

For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-306-23		
REF	Catalogue Number LC3006		
 LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY

Manufactured for:
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].



6625930623

Rx ONLY



P42573 Rev C

DO NOT REUSE

CAUTION


CONSULT
IFU

stryker®

SPY Elite Kit
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kit for use with the SPY Elite System
For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-306-13		
REF	Catalogue Number LC3006		
LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY
Contents	<ul style="list-style-type: none">• One 25 mg vial of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, <i>sterile</i>• One 10 ml vial of Sterile Water for Injection, USP• One SPY Elite Drape (ND8000), <i>sterile</i>• One SPY Elite Kit IEL• One SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Insert		

DO NOT REUSE
Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged.

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].



Manufactured for:
Novadaq Technologies LLC
8329 Eastview Drive, Unit 101
Baltimore, MD 21236, USA
Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

6625930613
Customer Support
1-800-624-4422 Toll Free



stryker®



CAUTION



DO NOT REUSE



Rx ONLY




CONSULT
IEL

SPY Elite Kit
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kit for use with the SPY Elite System
For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-306-13		
REF	Catalogue Number LC3006		
LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY
Contents	<ul style="list-style-type: none">• One 25 mg vial of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, <i>sterile</i>• One 10 ml vial of Sterile Water for Injection, USP• One SPY Elite Drape (ND8000), <i>sterile</i>• One SPY Elite Kit IFLU• One SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Insert		

DO NOT REUSE
Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged.

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].



Manufactured for:
Novadaq Technologies LLC
8329 Eastview Drive, Unit 101
Baltimore, MD 21236, USA
Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

6625930613
Customer Support
1-800-624-4422 Toll Free



stryker®



CAUTION



DO NOT REUSE



Rx ONLY





CONSULT
IEL

SPY Lymphatics Pack

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per Vial

Kits for use with the Advanced Imaging Modality (AIM) L10 and L11 Light Sources and 1688 Camera System, and the PINPOINT System
For lymphatic mapping

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

	NDC	66259-937-26		Use-by Date DD-MMM-YYYY
	REF	Catalogue Number PP9037		
LOT	Batch Code YYMMN			

Manufactured for:
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada
Assembled and distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].



6625993726



CAUTION



CONSULT
IFU

Rx ONLY

DO NOT REUSE

stryker®



P42578 Rev D

SPY Lymphatics Pack

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the Advanced Imaging Modality (AIM) L10 and L11 Light Sources and 1688 Camera System, and the PINPOINT System
For lymphatic mapping

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

	NDC	66259-937-26		Use-by Date DD-MMM-YYYY
	REF	Catalogue Number PP9037		
LOT	Batch Code YYMMIN			

Manufactured for:
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada
Assembled and distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].



6625993726



CAUTION



CONSULT
IFU

Rx ONLY

DO NOT REUSE


stryker®

P42578 Rev D


SPY Lymphatics Pack
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the Advanced Imaging Modality (AIM) L10 and L11 Light Sources and 1688 Camera System, and the PINPOINT System
For lymphatic mapping

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-937-26		
REF	Catalogue Number PP9037		
LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY
Contents	Components for six (6) SPY Lymphatics Procedures (packaged as six (6) SPY Lymphatics Kits): <ul style="list-style-type: none">• Six 25 mg vials of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, <i>sterile</i>• Twelve 10 ml vials of Sterile Water for Injection, USP• Six 10 ml syringes, <i>sterile</i>• Six luer-lock 10 ml syringes with control handles, <i>sterile</i>• Twelve 22-gauge, 3.5-inch needles, <i>sterile</i>• Labels for syringes• Six SPY Lymphatics Kit IFUs• Six SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Inserts		

DO NOT REUSE
Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged.

 Manufactured for:
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada
Assembled and distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].

 ONLY



DO NOT REUSE



CAUTION



CONSULT
IFU




6625993726


stryker®

SPY Lymphatics Pack
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the Advanced Imaging Modality (AIM) L10 and L11 Light Sources and 1688 Camera System, and the PINPOINT System
For lymphatic mapping

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-937-26		
REF	Catalogue Number PP9037		
LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY

 Manufactured for:
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada
Assembled and distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].

 ONLY



DO NOT REUSE



CAUTION



CONSULT
IFU



6625993726

stryker®

SPY Lymphatics Kit

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kit for use with the Advanced Imaging Modality (AIM) L10 and L11 Light Sources and 1688 Camera System, and the PINPOINT System
For lymphatic mapping

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-937-16		
REF	Catalogue Number PP9037		
LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY	
Contents	Components for one (1) SPY Lymphatics Procedure • One 25 mg vial of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, sterile • Two 10 ml vials of Sterile Water for Injection, USP • One 10 ml syringe, sterile • One luer-lock 10 ml syringe with control handle, sterile • Two 22 gauge, 3.5-inch needles, sterile • Labels for syringes • One SPY Lymphatics Kit IFU • One SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Insert		

DO NOT REUSE

Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].



Manufactured for:
Novadaq Technologies LLC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada
Assembled and distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

6625993716
Customer Support
1-800-624-4422 Toll Free

P42579 Rev D

SPY Lymphatics Kit

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kit for use with the Advanced Imaging Modality (AIM) L10 and L11 Light Sources and 1688 Camera System, and the PINPOINT System
For lymphatic mapping

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-937-16		
REF	Catalogue Number PP9037		
LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY	
Contents	Components for one (1) SPY Lymphatics Procedure • One 25 mg vial of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, sterile • Two 10 ml vials of Sterile Water for Injection, USP • One 10 ml syringe, sterile • One luer-lock 10 ml syringe with control handle, sterile • Two 22 gauge, 3.5-inch needles, sterile • Labels for syringes • One SPY Lymphatics Kit IFU • One SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Insert		

DO NOT REUSE

Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].



Manufactured for:
Novadaq Technologies LLC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada
Assembled and distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625


6625993716
Customer Support
1-800-624-4422 Toll Free

P42579 Rev D

SPY Minimally Invasive Surgery (SPY-MIS) Pack
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the Advanced Imaging Modality (AIM) and PINPOINT Systems
For visualization of vessels, blood flow and tissue perfusion
For visualization of extrahepatic biliary ducts

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-936-25	
REF	Catalogue Number PP9036	
LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY

Manufactured for:
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada
Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free
Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].



6625993625



CAUTION



CONSULT
IFU

Rx ONLY

DO NOT REUSE


stryker®

P43124 Rev C

SPY Minimally Invasive Surgery (SPY-MIS) Pack
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the Advanced Imaging Modality (AIM) and PINPOINT Systems
For visualization of vessels, blood flow and tissue perfusion
For visualization of extrahepatic biliary ducts

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-936-25	
REF	Catalogue Number PP9036	
LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY

Manufactured for:
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada
Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free
Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].



6625993625



CAUTION



CONSULT
IFU

Rx ONLY

DO NOT REUSE

stryker®


P43124 Rev C

13.5

SPY Minimally Invasive Surgery (SPY-MIS) Pack
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial


Kits for use with the Advanced Imaging Modality (AIM) and PINPOINT Systems
For visualization of vessels, blood flow and tissue perfusion
For visualization of extrahepatic biliary ducts

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-936-25		
REF	Catalogue Number PP9036		
LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY
Contents	Components for six (6) SPY-MIS Procedures (packaged as six (6) SPY MIS Kits): <ul style="list-style-type: none">• Six 25 mg vials of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, <i>sterile</i>• Six 10 ml vials of Sterile Water for Injection, USP• Twelve 3 ml syringes, <i>sterile</i>• Twelve 10 ml syringes, <i>sterile</i>• Twelve 18 G, 1 inch needles, <i>sterile</i>• Labels for syringes• Six SPY-MIS Kit IFUs• Six SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Inserts		



DO NOT REUSE
Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged.

 **Manufactured for:**
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Rx ONLY



DO NOT REUSE



CAUTION



CONSULT IFU


stryker®

P43123 Rev E

SPY Minimally Invasive Surgery (SPY-MIS) Pack
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the Advanced Imaging Modality (AIM) and PINPOINT Systems
For visualization of vessels, blood flow and tissue perfusion
For visualization of extrahepatic biliary ducts

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-936-25		
REF	Catalogue Number PP9036		
LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY



6625993625

Manufactured for:
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada
Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].

Rx ONLY



DO NOT REUSE



CAUTION



CONSULT IFU

stryker®

P43123 Rev E

SPY Minimally Invasive Surgery (SPY-MIS) Kit
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kit for use with the Advanced Imaging Modality (AIM) and PINPOINT Systems
For visualization of vessels, blood flow and tissue perfusion
For visualization of extrahepatic biliary ducts

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-936-15		
REF	Catalogue Number PP9036		
LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY	
Contents	Components for one (1) SPY-MIS Procedure: • One 25 mg vial of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, <i>sterile</i> • One 10 ml vial of Sterile Water for Injection, USP • Two 3 ml syringes, <i>sterile</i> • Two 10 ml syringes, <i>sterile</i> • Two 18 G, 1 inch needles, <i>sterile</i> • Labels for syringes • One SPY-MIS Kit IFU • One SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Insert		

DO NOT REUSE
Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged.

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].



Manufactured for:
Mediag Technology LLC
8329 East 1st Avenue, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

66259936 15
Customer Support
1-800-624-4422 Toll Free



stryker®
IFU
DO NOT REUSE
CONSULT



CAUTION



Rx ONLY

SPY Minimally Invasive Surgery (SPY-MIS) Kit
with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kit for use with the Advanced Imaging Modality (AIM) and PINPOINT Systems
For visualization of vessels, blood flow and tissue perfusion
For visualization of extrahepatic biliary ducts

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-936-15		
REF	Catalogue Number PP9036		
LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY	
Contents	Components for one (1) SPY-MIS Procedure: • One 25 mg vial of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, <i>sterile</i> • One 10 ml vial of Sterile Water for Injection, USP • Two 3 ml syringes, <i>sterile</i> • Two 10 ml syringes, <i>sterile</i> • Two 18 G, 1 inch needles, <i>sterile</i> • Labels for syringes • One SPY-MIS Kit IFU • One SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Insert		

DO NOT REUSE
Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged.

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].



Manufactured for:
Mediag Technology LLC
8329 East 1st Avenue, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

66259936 15
Customer Support
1-800-624-4422 Toll Free



stryker®
IFU
DO NOT REUSE
CONSULT



CAUTION



Rx ONLY

SPY-PHI Pack

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the SPY-PHI System
For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-903-24	LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY
REF	Catalogue Number HH9006			

Manufactured for:
Novadac Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].



Rx ONLY



DO NOT RELISE



CAUTION

6625990324



CONSULT
IFU

stryker®

SPY-PHI Pack

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kits for use with the SPY-PHI System
For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-903-24	LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY
REF	Catalogue Number HH9006			

Manufactured for:
Novadac Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].



Rx ONLY



DO NOT RELISE



CAUTION

6625990324



CONSULT
IFU

stryker®

SPY-PHI Pack

with


SPY AGENT™ GREEN (indocyanine green for injection, USP)

25 mg per vial

Kits for use with the SPY-PHI System

For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-903-24		
REF	Catalogue Number HH9006		
LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY
Contents	Components for six (6) SPY-PHI Procedures (packaged as six (6) SPY-PHI Kits): <ul style="list-style-type: none">• Six 25 mg vials of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, <i>sterile</i>• Six 10 ml vials of Sterile Water for Injection• Six SPY-PHI Drapes (HH2000), <i>sterile</i>• Six SPY-PHI Kit IFUs• Six SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Inserts		

DO NOT REUSE
Outside packaging and tray are not sterile – contents of the individual components are sterile.
Do not use if immediate packaging is damaged.

**Manufactured for:**
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].

**Rx ONLY**


DO NOT REUSE


CAUTION


CONSULT
IFU

stryker®

SPY-PHI Pack

with


SPY AGENT™ GREEN (indocyanine green for injection, USP)

25 mg per vial

Kits for use with the SPY-PHI System

For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-903-24		
REF	Catalogue Number HH9006		
LOT	Batch Code YYMMN		Use-by Date DD-MMM-YYYY

**Manufactured for:**
Novadaq Technologies ULC
8329 Eastlake Drive, Unit 101
Burnaby, BC V5A 4W2, Canada

Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

Customer Support
1-800-624-4422 Toll Free

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].

**Rx ONLY**


DO NOT REUSE


CAUTION


CONSULT
IFU

stryker®

SPY-PHI Kit

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kit for use with the SPY-PHI System

For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-903-14		
REF	Catalogue Number HH9006		
LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY	
Contents	<ul style="list-style-type: none">• One 25 mg vial of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, sterile• One 10 ml vial of Sterile Water for Injection, USP• One SPY-PHI Drape (HH2000), sterile• One SPY-PHI Kit IFU• One SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Insert		

DO NOT REUSE

Outside packaging and tray are not sterile – contents of the individual components are sterile.

Do not use if immediate packaging is damaged.

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].



Manufactured for:
Novadaq Technologies LLC
2500 East Beaver Creek Road, Unit 101
Burnaby, BC V5A 4W2, Canada
Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

6625990314
Customer Support
1-800-624-4422 Toll Free



Rx ONLY



CAUTION



DO NOT REUSE



SPY-PHI Kit

with
SPY AGENT™ GREEN (indocyanine green for injection, USP)
25 mg per vial

Kit for use with the SPY-PHI System

For visualization of vessels, blood flow and tissue perfusion

Reconstitute with Accompanying Sterile Water Diluent Before Use
Use within 6 hours of reconstitution

NDC	66259-903-14		
REF	Catalogue Number HH9006		
LOT	Batch Code YYMMN	Use-by Date DD-MMM-YYYY	
Contents	<ul style="list-style-type: none">• One 25 mg vial of SPY AGENT™ GREEN (indocyanine green for injection, USP), lyophilized, sterile• One 10 ml vial of Sterile Water for Injection, USP• One SPY-PHI Drape (HH2000), sterile• One SPY-PHI Kit IFU• One SPY AGENT™ GREEN (indocyanine green for injection, USP) Package Insert		

DO NOT REUSE

Outside packaging and tray are not sterile – contents of the individual components are sterile.

Do not use if immediate packaging is damaged.

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].



Manufactured for:
Novadaq Technologies LLC
2500 East Beaver Creek Road, Unit 101
Burnaby, BC V5A 4W2, Canada
Distributed by:
Carestream Health, Inc.
882 Linden Avenue
Rochester, NY USA 14625

6625990314
Customer Support
1-800-624-4422 Toll Free



Rx ONLY



CAUTION



DO NOT REUSE





Ramesh
Raghavachari

Digitally signed by Ramesh Raghavachari
Date: 3/16/2022 05:57:33PM
GUID: 502d0913000029f375128b0de8c50020