

CYLTEZO®
(adalimumab-adbm)
Injection

NDC 0597-0370-72

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Syringe

NOT FOR SALE
Rx only

For Subcutaneous Use Only

1 Single-Dose Prefilled Syringe

Boehringer
Ingelheim

40 mg/0.8 mL
Injection (adalimumab-adbm)

NDC 0597-0370-72

ATTENTION PHARMACIST:
Each Patient is Required to Receive the
Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.

NDC 0597-0370-72

CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Syringe

GTIN
SN
EXP
LOT

00305970370727
12345678901234
MMYYYY
1234567

50-51601001

NDC 0597-0370-72

CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Syringe

Citrate Free

NOTE TO PHARMACIST:
(Place Label Here)

Do not use beyond the expiration date.

10010943-03

NDC 0597-0370-72

CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Syringe

Citrate Free

For more information about
CYLTEZO, including current
prescribing information and
Medication Guide, go to
www.cyltezo.com, scan the code,
or call 1-800-452-6257.
Boehringer Ingelheim
Pharmaceuticals, Inc. at

NDC 0597-0370-72

CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Syringe

Product and trademark licensed from:
Boehringer Ingelheim
International GmbH

© 2022 Boehringer Ingelheim
Pharmaceuticals, Inc.
ALL RIGHTS RESERVED

Manufactured by:
Boehringer Ingelheim
Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7594G

NDC 0597-0370-72

CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL


For Subcutaneous Use Only

1 Single-Dose Prefilled Syringe

Citrate Free

1010005970370727






front/outside

10 mg/0.2 mL
For Subcutaneous Use Only
2 Single-Dose Prefilled Syringes

CYLTEZO[®]
(adalimumab-adbm)
Injection



2 Single-Dose Prefilled Syringes



Rx only
For Subcutaneous Use Only

10 mg/0.2 mL

CYLTEZO[®]
(adalimumab-adbm)
Injection

ATTENTION PHARMACIST:
Each Patient is Required to Receive the
Enclosed Medication Guide.
The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.

NDC 0597-0400-89

NDC 0597-0400-89

CYLTEZO[®]
(adalimumab-adbm)
Injection

10 mg/0.2 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes



GTIN
SN
EXP
LOT

00305970400899
12345678901234
MMMMYYYY
1234567



1041311001

NDC 0597-0400-89

CYLTEZO[®]
(adalimumab-adbm)
Injection

10 mg/0.2 mL

For Subcutaneous Use Only

This Carton Contains:

- 2 dose trays (each containing 1 single-dose prefilled syringe with a fixed 1/8 inch length needle)
- 2 alcohol preps
- 1 Package Insert
- 1 Medication Guide
- 1 Instructions for Use

See package insert for full Prescribing Information.
Needle Cap for Syringe Contains Dry Natural Rubber.



01700305970400899

NOTE TO PHARMACIST:
{Place Prescription Label Here}

Each Sterile Single-Dose Prefilled Syringe Contains:
Adalimumab-adbm..... 10 mg
Glacial acetic acid..... 0.03 mg
Polysorbate 80..... 0.2 mg
Sodium acetate trihydrate..... 0.61 mg
Trehalose dihydrate..... 16.3 mg
Water for Injection, USP.

Contains no preservatives. No U.S. standard of potency.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE.
If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator ____/____/____.

Do not use beyond the expiration date.



10011318-01

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
Boehringer Ingelheim
Birkbeck Drive
Bismarck, ND 58503
© 2023, Boehringer Ingelheim
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number: 2006
CT199948

Citrate
Free

NDC 0597-0400-89

CYLTEZO[®]
(adalimumab-adbm)
Injection

10 mg/0.2 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

2 Single-Dose Prefilled Syringes
For Subcutaneous Use Only
10 mg/0.2 mL





CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0400-89

For more information about CYLTEZO, including
current prescribing information and Medication
Guide, go to www.cyltezo.com, scan the code, or
call Boehringer Ingelheim Pharmaceuticals, Inc.
at 1-800-545-2427.



QUICKTIPS for USING CYLTEZO® (adalimumab-adbm) Injection
For complete instructions refer to the Instructions for Use in the carton.
Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
			
<ul style="list-style-type: none">• Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.• Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.• Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">• Inspect the prefilled syringe, medicine, and expiration date.• Pull the cap straight off the syringe. Do Not let the needle touch anything.• Do Not recap the needle.	<ul style="list-style-type: none">• Squeeze the skin around the injection site and hold firmly.• Hold the syringe at about a 45 degree angle to the injection site.• Insert the needle into the injection site with a quick, smooth motion.	<ul style="list-style-type: none">• Slowly push the plunger down fully until the plunger reaches bottom of syringe.• Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.


CUTSCORE

CUTSCORE

2 Single-Dose Prefilled Syringes For Subcutaneous Use Only

20 mg/0.4 mL

CYLTEZO®
(adalimumab-adbm)
Injection



2 Single-Dose Prefilled Syringes



Rx only
For Subcutaneous Use Only

20 mg/0.4 mL

CYLTEZO®
(adalimumab-adbm)
Injection

ATTENTION PHARMACIST:
Each Patient is Required to Receive the
Enclosed Medication Guide.
The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.

NDC 0597-0405-80

NDC 0597-0405-80

CYLTEZO®
(adalimumab-adbm)
Injection

20 mg/0.4 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

GTIN 12345678901234
SN 12345678901234
EXP MMMYYYY
LOT 1234567



60-9111001

NDC 0597-0405-80

CYLTEZO®
(adalimumab-adbm)
Injection

20 mg/0.4 mL

For Subcutaneous Use Only

This Carton Contains:

- 2 dose trays (each containing 1 single-dose prefilled syringe with a fixed 1/8 inch length needle)
- 2 alcohol preps
- 1 Package Insert
- 1 Medication Guide
- 1 Instructions for Use

See package insert for full Prescribing Information.
Needle Cap for Syringe Contains Dry Natural Rubber.



01700305970405801

NOTE TO PHARMACIST:
{Place Prescription Label Here}

Each Sterile Single-Dose Prefilled Syringe Contains:

Adalimumab-adbm.....	20 mg
Glacial acetic acid.....	0.06 mg
Polysorbate 80.....	0.4 mg
Sodium acetate trihydrate.....	1.21 mg
Trehalose dihydrate.....	32.5 mg
Water for Injection, USP.	

Contains no preservatives. No U.S. standard of potency.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator ____/____/____.

Do not use beyond the expiration date.



10011161-03

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
Boehringer Ingelheim
Bismarck, CA, USA
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number: 2006
CT7974F

Citrate Free

NDC 0597-0405-80

CYLTEZO®
(adalimumab-adbm)
Injection

20 mg/0.4 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

NDC 0597-0405-80


CYLTEZO®
(adalimumab-adbm)
Injection

20 mg/0.4 mL

For Subcutaneous Use Only
2 Single-Dose Prefilled Syringes





For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call [1-800-542-2527](tel:1-800-542-2527).






QUICKTIPS for USING CYLITEZO® (adalimumab-adbm) Injection

For complete instructions refer to the **Instructions for Use** in the carton. Do not try to inject CYLITEZO® yourself until you have been shown the right way to give the injection.

<p>1 Choose Injection Site and Clean</p>  <p>Abdomen or Thighs</p>	<p>2 Inspect and Uncap</p> 	<p>3 Prepare Injection</p>  <p>45°</p>	<p>4 Administer Dose</p>  <p>Plunger</p>
<ul style="list-style-type: none">Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">Inspect the prefilled syringe, medicine, and expiration date.Pull the cap straight off the syringe. Do Not let the needle touch anything.Do Not recap the needle.	<ul style="list-style-type: none">Squeeze the skin around the injection site and hold firmly.Hold the syringe at about a 45 degree angle to the injection site.Insert the needle into the injection site with a quick, smooth motion.	<ul style="list-style-type: none">Slowly push the plunger down fully until the plunger reaches bottom of syringe.Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

For more information about CYLITEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

NDC 0597-0370-82



CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes


Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH

© 2022 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7422G

NDC 0597-0370-82



CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

NOTE TO PHARMACIST:
(Place Prescription Label Here)

For Subcutaneous Use Only

This Carton Contains:

- 2 dose trays (each containing 1 single-dose prefilled syringe with a fixed 1/2 inch length needle)
- 2 alcohol preps
- 1 Package Insert
- 1 Medication Guide
- 1 Instructions for Use

See package insert for full Prescribing Information. Needle Cap for Syringe Contains Dry Natural Rubber.

Each Sterile Single-Dose Prefilled Syringe Contains:

Adalimumab-adbm.....	40 mg
Glacial acetic acid.....	0.13 mg
Poly sorbate 80.....	0.8 mg
Sodium acetate trihydrate.....	2.4 mg
Trehalose dihydrate.....	65.0 mg
Water for Injection, USP.....	


Contains no preservatives. No U.S. standard of potency.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator / / .

(01)700303970370824

NDC 0597-0370-82





CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only


2 Single-Dose Prefilled Syringes

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

1001094443

NDC 0597-0370-82




CYLTEZO®
(adalimumab-adbm)
Injection


40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes




GTN 00305970370826
SN 12345678901234
EXP MMYYY
LOT 1234567



30-79601001

NDC 0597-0370-82



CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

Rx only

2 Single-Dose Prefilled Syringes

Boehringer Ingelheim

CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only


2 Single-Dose Prefilled Syringes

NDC 0597-0370-82

ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.

NDC 0597-0370-82




CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

Rx only



2 Single-Dose Prefilled Syringes

Boehringer Ingelheim


CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

NDC 0597-0370-82




CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only





2 Single-Dose Prefilled Syringes

S
S
S
S
S



QUICKTIPS for USING CYLITEZO® (adalimumab-adbm) Injection

For complete instructions refer to the **Instructions for Use** in the carton. Do not try to inject CYLITEZO® yourself until you have been shown the right way to give the injection.

<p>1 Choose Injection Site and Clean</p>  <p>Abdomen or Thighs</p>	<p>2 Inspect and Uncap</p> 	<p>3 Prepare Injection</p>  <p>45°</p>	<p>4 Administer Dose</p>  <p>Plunger</p>
<ul style="list-style-type: none">Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">Inspect the prefilled syringe, medicine, and expiration date.Pull the cap straight off the syringe. Do Not let the needle touch anything.Do Not recap the needle.	<ul style="list-style-type: none">Squeeze the skin around the injection site and hold firmly.Hold the syringe at about a 45 degree angle to the injection site.Insert the needle into the injection site with a quick, smooth motion.	<ul style="list-style-type: none">Slowly push the plunger down fully until the plunger reaches bottom of syringe.Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

For more information about CYLITEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



Susan
Kirshner

Digitally signed by Susan Kirshner

Date: 10/03/2022 10:28:24AM

GUID: 508da6db000266b77da0ba4bfa620030