

The trade dress of this ZYRTEC<sup>®</sup> package is subject to trademark protection.  
Active ingredient made in Belgium.

Indoor & Outdoor Allergies

**ZYRTEC<sup>®</sup>**  
Cetirizine HCl tablets  
5 mg antihistamine

NDC 0000-000-00

**ZYRTEC<sup>®</sup>**  
Cetirizine HCl tablets  
5mg antihistamine

Indoor & Outdoor Allergies  
5 mg Strength  
5 mg Tablets

Take 1 to 2 tablets\* once daily depending on the strength of the product.  
\*Adults 65 and older take only one tablet per day.

24 Hour relief of

Runny Nose

Itchy, Watery Eyes

Itchy Throat or Nose

Sneezing

35 Tablets 5 mg each

Actual Size



Actual Size

LOT:  
EXP:



Active ingredient made in Belgium.  
Distributed by:  
Janssen & Johnson Consumer Inc.  
Janssen Pharmaceutica Division  
P.O. Box 2001  
Kenilworth, NJ 07033 USA  
©J&J 2014 zyrtec.com



00000000

<p><b>Directions</b></p> <p>Swallow whole. Do not crush, chew, or break. Take with water.</p> <p>Children under 6 years of age: 5 mg once daily.</p> <p>Children 6 to 11 years of age: 5 mg once daily.</p> <p>Children 12 to 17 years of age: 5 mg once daily.</p> <p>Adults 18 years of age and older: 5 mg once daily.</p> <p>Adults 65 years of age and older: 5 mg once daily.</p> <p>Take 1 to 2 tablets once daily depending on the strength of the product.</p>	<p><b>Warnings</b></p> <p>Do not take if you are allergic to cetirizine HCl or any of the ingredients.</p> <p>Do not take if you are taking other antihistamines.</p> <p>Do not take if you are taking other sedating drugs.</p> <p>Do not take if you are taking other drugs that may interact with cetirizine HCl.</p> <p>Do not take if you are taking other drugs that may increase the risk of drowsiness.</p> <p>Do not take if you are taking other drugs that may increase the risk of dizziness.</p> <p>Do not take if you are taking other drugs that may increase the risk of dry mouth.</p> <p>Do not take if you are taking other drugs that may increase the risk of constipation.</p> <p>Do not take if you are taking other drugs that may increase the risk of urinary retention.</p> <p>Do not take if you are taking other drugs that may increase the risk of blurred vision.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty swallowing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty breathing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty urinating.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty seeing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty hearing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty speaking.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty thinking.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty remembering.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty concentrating.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty focusing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty understanding.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty communicating.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty interacting.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty relating.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty connecting.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty cooperating.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty assisting.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty supporting.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty helping.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty finishing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty completing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty performing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty executing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty implementing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty putting into effect.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty carrying out.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty accomplishing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty achieving.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty attaining.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty obtaining.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty securing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty gaining.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty earning.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty making.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty doing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty acting.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty behaving.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty performing.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty reacting.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty responding.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty replying.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty answering.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty returning.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty restoring.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty reviving.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing back.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing up.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing forward.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing in.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing out.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing down.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing back.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing up.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing forward.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing in.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing out.</p> <p>Do not take if you are taking other drugs that may increase the risk of difficulty bringing down.</p>
---	---

Important: Read all product information before using. Keep this card for important information.

LOT: XXXXXX  
EXP: XXXXXX

NDC 0000-000-00

**ZYRTEC**  
ALLERGY

Cetirizine HCl/  
antihistamine  
5 mg tablets

**35**  
Tablets  
5 mg each

**Active ingredient (in each tablet)**

Cetirizine HCl 5 mg ..... **Purpose**  
Antihistamine  
Uses temporarily relieves these symptoms due to hay fever or other upper  
respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching  
of the nose or throat

**Warnings**

Do not use if you have ever had an allergic reaction to this product or any of its ingredients  
or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or  
kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or  
pharmacist before use if you are taking tranquilizers or sedatives. When using this product  
■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers  
may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery.  
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help  
right away, if pregnant or breast-feeding, ■ if breast-feeding, not recommended if  
pregnant, ask a health professional before use. Keep out of reach of children. In case of  
overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Purpose**  
Antihistamine



**Directions** Adults and children 6 years and over: 1 tablet (5 mg) or 2 tablets  
(10 mg) once daily depending upon severity of symptoms; do not take more  
than 2 tablets (10 mg) in 24 hours. Adults 65 years and over: 1 tablet (5 mg)  
once daily; do not take more than 1 tablet (5 mg) in 24 hours. Children UNDER  
6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a  
doctor. Other information ■ store between 20° to 25°C (68° to 77°) ■ do not  
use if foil liner seal imprinted with "ZYRTEC®" is broke or missing  
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium,  
hydroxyethylcellulose, lactose monohydrate, magnesium stearate, microcrystalline  
cellulose, polyethylene glycol, titanium dioxide

**Questions?** call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)  
The trade dress of this ZYRTEC® package is subject to trademark protection.  
Active ingredient made in Belgium.  
Distributed by: JOHNSON & JOHNSON CONSUMER INC.  
McNeil Consumer Healthcare Division Fort Washington, PA 19034 USA  
©J&J 200X **zyrtec.com**

00000000



**ZYRTEC**  
Cetirizine HCl tablets **Allergy**  
5 mg antihistamine

Indoor & Outdoor Allergies  
24 Hour relief of

- Sneezing
- Runny nose
- Itchy, watery eyes
- Itchy throat or nose

5 mg Strength

**Drug Facts**  
Active ingredient (in each tablet): Cetirizine HCl 5 mg. Purpose: Antihistamine. How it works: orally reduces these symptoms due to hay fever or other upper respiratory allergies: runny nose, sneezing, itchy, watery eyes, itching of the nose or throat.  
Warnings: Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

To Open: While Folded on Line, Tear At Slit

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product, no drowsiness may occur, so avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness. Do not drink or drive a motor vehicle or operate machinery. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breast-feeding, or if breast-feeding not recommended, or if pregnant, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Warnings: Adults and children 6 years and over: 1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours. Adults 65 years and over: 1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor. Other information: Store between 20° to 25°C (68° to 77°F); do not use if pouch is torn or damaged. Each the ingredients: or hydroxyzine hydrochloride, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, titanium dioxide, croscarmellose sodium. Question? call 1-800-222-1222 (10-10mg) or 1-800-222-1222 (5-mg). Active ingredient made in Belgium. Distributed by: JOHNSON & JOHNSON CONSUMER INC., Medical Consumer Products Division, Fort Washington, PA 19074 USA. Call 1-800-222-1222 www.zyrtec.com Pat. www.zyrtec.com 00000000