

NDC 16837-889-20

MAXIMUM STRENGTH

Pepcid[®] AC


Famotidine Tablets 20 mg Acid Reducer

ICY COOL MINT
Flavored Coating

Just One Tablet!
Prevents & Relieves Heartburn
Due to Acid Indigestion

actual size

20 Tablets



Questions or comments?
1-800-755-4008 (toll-free) or
215-273-8755 (local)

Drug Facts (continued)
Other information
• read the directions and warnings before use
• keep the carton in a cool, dry place
• store at 20°-25°C (68°-77°F)
• protect from moisture
Inactive ingredients
carnauba wax, FD&C blue #1, aluminum lake, flavors, hypromellose 29 mg, magnesium stearate, microcrystalline cellulose, polyethylene glycol, poly sorbate 80, pregelatinized starch, sucralose, talc, titanium dioxide

20 Tablets
Pepcid[®] AC
MAXIMUM STRENGTH

Drug Facts (continued)
Directions
• adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
• to prevent symptoms, swallow 1 tablet with a glass of water at any time from 30 minutes to 1 hour before eating food or drinking beverages that cause heartburn.
• do not use more than 2 tablets in 24 hours.
• children under 12 years: ask a doctor.
Warnings
• frequent heartburn, particularly with heartburn over time, get medical help or contact a poison control center. Call right away, (1-800-222-1222).
• stomach pain
• kidney disease
• stomach pain
• frequent heartburn, particularly with heartburn over time, get medical help or contact a poison control center. Call right away, (1-800-222-1222).
Drug Facts (continued)
Directions
• adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
• to prevent symptoms, swallow 1 tablet with a glass of water at any time from 30 minutes to 1 hour before eating food or drinking beverages that cause heartburn.
• do not use more than 2 tablets in 24 hours.
• children under 12 years: ask a doctor.
Warnings
• frequent heartburn, particularly with heartburn over time, get medical help or contact a poison control center. Call right away, (1-800-222-1222).
• stomach pain
• kidney disease
• stomach pain
• frequent heartburn, particularly with heartburn over time, get medical help or contact a poison control center. Call right away, (1-800-222-1222).

Drug Facts (continued)
Adverse reaction (in each tablet) Purpose
Famotidine 20 mg. Acid reducer
Do not use
• If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor for serious conditions. See your doctor for relief of heartburn associated with acid indigestion and sour stomach.
• Ask a doctor before use if you have had heartburn over 3 months. This may be a sign of a more serious condition.
• heartburn with lightheadedness, sweating, or dizziness
• chest pain or shoulder pain with shortness of breath
• neck or shoulder, or lightheadedness, or fainting or other allergic reactions
Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.
Uses
relieves heartburn associated with acid indigestion and sour stomach
prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages.
Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.

Tips for Managing Heartburn
• Do not lie flat or bend over after eating. • Do not wear tight-fitting clothing around the stomach. • Do not eat before bedtime. • Raise the head of your bed. • Avoid heartburn-causing foods such as rich, spicy, fatty, or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables. • Eat slowly and avoid big meals.
• If overweight, lose weight. • Quit smoking.

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Do not use if foil inner seal is broken or missing
SAFETY IS BROKEN OR MISSING

LOT:
EXP:

Famotidine (active ingredient) made in Hungary.
Manufactured by: **JOHNSON & JOHNSON CONSUMER INC.**
Mylan Consumer Healthcare Division
Fort Washington, PA, 19034 USA
Pepcid[®] and Pepcid AC[®] are registered trademarks.
Pat. www.jnj.com

300056654730002

Do not use: If the inner seal imprinted with **SEALED FOR YOUR SAFETY** is broken or missing.
Active ingredient (in each tablet)

Famotidine 20 mg Acid reducer

Uses: • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages. **Warnings:** Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers. Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness

• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease. Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-232-1223).

MAXIMUM STRENGTH

NDC 16837-889-20

Pepcid[®]
AC

Famotidine Tablets 20 mg / Acid Reducer

ICY COOL MINT
Flavored Coating

Just One Tablet!

Prevents & Relieves Heartburn
Due to Acid Indigestion

actual size

20 Tablets

Directions: • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor

Other information: • read the directions and warnings before use • keep the carton. It contains important information. • store at 20°-25°C (68°-77°F) • protect from moisture
Inactive ingredients: croscarmellose, FD&C blue no. 1 aluminum lake, flavors, hypromellose 2910, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyacrylate 60, pregelatinized starch, sucralose, talc, titanium dioxide
Questions or comments? 1-800-755-4000 (to 1-free) or 215-233-0755 (no text)

Famotidine (active ingredient) made in Hungary
Distributed by: JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA © McNeil Consumer
Pharmaceuticals Co. 2023 PEPCID[®] and PEPCID A[®]
are registered trademarks. Pat. www.janvape.com

EXP.

LOT:

10/2023 (USA) 01/2023

Do not use if foil inner seal imprinted with **SEALED FOR YOUR SAFETY** is broken or missing
Ac: inactive ingredient for each tablet

Famotidine 20 mg Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages. **Warnings:** A large alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor • with other acid reducers. Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness

• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease. Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor or if • your heartburn continues or worsens • you need to take this product or more than 14 days. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).



• frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease. Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor or if • your heartburn continues or worsens • you need to take this product or more than 14 days. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

• frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease. Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor or if • your heartburn continues or worsens • you need to take this product or more than 14 days. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

MAXIMUM STRENGTH

NDC 16 837-889-40

Pepcid[®]
AC

Famotidine Tablets 20 mg / Acid Reducer

ICY COOL MINT
Flavored Coating

Just One Tablet!
Prevents & Relieves Heartburn
Due to Acid Indigestion



actual size

40 Tablets

Directions • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor

Other information • read the directions and warnings before use • keep the carton. It contains important information. • store at 20°-25°C (68°-77°F) • protect from moisture

Inactive ingredients contains wax, FD&C blue no. 1 aluminum lake, flavors, hypromellose 2910, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyacrylate E1, pregelatinized starch, sucralose, talc, titanium dioxide

Questions or comments? 1-800-755-4888 (ask-free) or 215-233-0755 (collect)
Famotidine (active ingredient) made in Hungary
Distributed by: JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA

©McNeil Consumer Pharmaceuticals Co. 2023 PEPCID[®] and PEPCID AC[®] are registered trademarks. Pat. www.janvapephs.com

30 (15 05 25) 12/15/23

EXP.

LOT:

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
12/14/2023 06:41:59 PM