

# EXCEDRIN

## MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine  
Pain Reliever/Pain Reliever Aid

8  
CAPLETS\*

\*CAPSULE-SHAPED TABLETS



Drug Facts	
<b>Active ingredients (in each caplet)</b>	<b>Purposes</b>
Acetaminophen 250 mg.....Pain reliever	
Aspirin 250 mg (NSAID)*.....Pain reliever	
Caffeine 65 mg.....Pain reliever aid	
*nonsteroidal anti-inflammatory drug	
<b>Use</b>	
• treats migraine	
<b>Warnings</b>	
<b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.	
<b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.	

Drug Facts (continued)	
<b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock	
<b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product	
<b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed	

DO NOT USE IF EXCEDRIN  
SAFETY SEAL AROUND CAP  
IS BROKEN OR MISSING.



Drug Facts (continued)	
<b>Medication overuse headache warning:</b> Headaches may worsen if this product is used for 10 or more days per month.	
<b>Caffeine warning:</b> The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.	
<b>Do not use</b>	
• if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer	
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
<b>Ask a doctor before use if</b>	
• you have never had migraines diagnosed by a health professional	
• you have a headache that is different from your usual migraines	
• you have the worst headache of your life	
• you have fever and stiff neck	
• you have headaches beginning after or caused by head injury, exertion, coughing or bending	
• you experienced your first headache after the age of 50	
• you have daily headaches	
• you have a migraine so severe as to require bed rest	
• you have liver disease	
• stomach bleeding warning applies to you	
• you have a history of stomach problems, such as heartburn	
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease	
• you are taking a diuretic	
• you have asthma	
• you have problems or serious side effects from taking pain relievers or fever reducers	
• you have vomiting with your migraine headache	
<b>Ask a doctor or pharmacist before use if you are</b>	
• taking a prescription drug for: • diabetes • gout • arthritis	
• under a doctor's care for any serious condition	
• taking any other drug	
• taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer	
<b>Stop use and ask a doctor if</b>	
• an allergic reaction occurs. Seek medical help right away.	
• you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better	

Drug Facts (continued)	
• your migraine is not relieved or worsens after first dose	
• new or unexpected symptoms occur	
• ringing in the ears or loss of hearing occurs	
<b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.	
<b>Keep out of reach of children.</b>	
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
<b>Directions</b>	
• do not use more than directed	
• adults: take 2 caplets with a glass of water	
• if symptoms persist or worsen, ask your doctor	
• do not take more than 2 caplets in 24 hours, unless directed by a doctor	
• under 18 years of age: ask a doctor	
<b>Other information</b>	
• store at or below 25°C (77°F)	
• close cap tightly after use	
• read all product information before using	
<b>Inactive ingredients</b>	benzoic acid, carnauba wax, FD&C blue no. 1, hypromellose, light mineral oil, low-substituted hydroxypropyl cellulose, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone, sorbitan monolaurate, stearic acid, titanium dioxide
<b>Questions or comments?</b>	1-800-468-7746

Distributed by:  
GSK Consumer Healthcare  
Warren, NJ 07059  
©2022 GSK or licensor.  
Trademarks owned or licensed by GSK.  
1001792

GSK Regulatory Spec Box Verified Date: 08.03.22

Drug Facts Info		
Drug Facts	Helvetica Neue LT Std Bold Cond Oblique	9 pt.
Drug Facts (continued)	Helvetica Neue LT Std Bold Cond Oblique / Cond	8 pt.
Headings	Helvetica Neue LT Std Bold Cond Oblique	8 pt.
Subheadings	Helvetica Neue LT Std Bold Cond	6 pt.
Body text	Helvetica Neue LT Std Bold Cond / Cond	6 pt.
Bullets	Helvetica Neue LT Std Roman	5 pt.
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hair lines from edge of box-i.e. Minimum of 2 spaces either side of Drug Fact Box		Yes
Tracking:	0	Horizontal Scale: 85%
Leading:	6.25 pt.	Maximum Characters/Inch: < 39
Barlines:	1.5 pt.	Hairlines: 0.5 pt.
Primary Display Panel Info		
Size of Net Wt/Contents (Smallest character height in inches)*		> 0.125 in.
PDP dimensions (in square inches)		5-25 sq. in.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		
Vertical PDP: N/A		Horizontal PDP: 35%

EXCEDRIN SAFETY SEALED • EXCEDRIN SAFETY SEALED  
TO OPEN PUSH DOWN THEN TURN • TO OPEN PUSH DOWN THEN TURN

**Drug Facts (continued)**

- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for:
  - diabetes
  - gout
  - arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
  - new or unexpected symptoms occur
  - ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

**Other information**

- store at or below 25°C (77°F)
- close cap tightly after use
- read all product information before using

**Inactive ingredients** benzoic acid, carnauba wax, FD&C blue no. 1, hypromellose, light mineral oil, low-substituted hydroxypropyl cellulose, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone, sorbitan monooleate, stearic acid, titanium dioxide

**Questions or comments?** 1-800-468-7746

Distributed by GSK Consumer Healthcare, Warren, NJ 07059  
©2022 GSK or its subsidiaries or its licensee.  
Trademarks are owned by or licensed to the GSK group of companies.

**DO NOT USE IF EXCEDRIN SAFETY SEAL AROUND CAP IS BROKEN OR MISSING.**



**EXCEDRIN**

**MIGRAINE**  
Acetaminophen, Aspirin (NSAID) and Caffeine  
Pain Reliever/Pain Reliever Aid

**Drug Facts**  
**Active ingredients (in each caplet)**

Acetaminophen 250 mg ..... Pain reliever  
Aspirin 250 mg (NSAID) ..... Pain reliever aid  
Caffeine 65 mg ..... Nonsteroidal anti-inflammatory drug

**Purposes**  
Pain reliever  
Pain reliever aid  
Nonsteroidal anti-inflammatory drug

PEEL RIGHT CORNER FOR FULL DRUG FACTS

**Drug Facts (continued)**

**Use** • treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Medication overdose headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life

GSK Regulatory Spec Box		Verified Date: 08.02.22
<b>Drug Facts Info</b>		
<b>Drug Facts</b>	Helvetica Neue LT Std Bold Cond Oblique	9 pt.
<b>Drug Facts (continued)</b>	Helvetica Neue LT Std Bold Cond Oblique / Cond	8 pt.
<b>Headings</b>	Helvetica Neue LT Std Bold Cond Oblique	8 pt.
<b>Subheadings</b>	Helvetica Neue LT Std Bold Cond	6 pt.
<b>Body text</b>	Helvetica Neue LT Std Bold Cond / Cond	6 pt.
<b>Bullets</b>	Helvetica Neue LT Std Roman	5 pt.
<b>Bullets on same lines: end of statement separated from bulleted statement by two ems</b>		Yes
<b>Spacing of the hair lines from edge of box-i.e. Minimum of 2 spaces either side of Drug Fact Box</b>		Yes
<b>Tracking:</b>	0 pt.	<b>Horizontal Scale:</b> 93%
<b>Leading:</b>	6 pt.	<b>Maximum Characters/Inch:</b> < 39
<b>Barlines:</b>	1.5 pt.	<b>Hairlines:</b> 0.5 pt.
<b>Primary Display Panel Info</b>		
<b>Size of Net Wt/Contents (Smallest character height in inches)*</b>		>0.0625 in.
<b>PDP dimensions (in square inches)</b>		< 5 sq. in.
<b>Ratio of Statement of Identity to Logo/Largest Copy on PDP</b>		
<b>Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)</b>		
<b>Vertical PDP:</b>	N/A	<b>Horizontal PDP:</b> 42%