

Contents: 1 mg treprostinil per mL.
Each mL also contains 0.5 mg sodium citrate dihydrate, 5.3 mg sodium chloride, 3 mg metacresol, 0.24 mg sodium hydroxide, and Water for Injection. Hydrochloric acid and sodium hydroxide may have been added to adjust pH.
Prior to intravenous infusion, Remodulin® may be diluted.

See Package Insert for Dosage and Administration.

United Therapeutics Corp., RTP, NC USA 27709




REMODULIN®
(treprostinil) Injection

20 mg/20 mL
(1 mg/mL)

For Subcutaneous
and Intravenous Infusion

20 mL multidose vial



NDC 68302-101-01

Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).

Keep out of reach of children.

Rx Only 10050 Rev. 01/2023

LOT XXXXXXX

EXP MM-YYYY

(b) (4)

Contents: 2.5 mg treprostinil per mL.
Each mL also contains 6.3 mg sodium citrate dihydrate, 5.3 mg sodium chloride, 3 mg metacresol, 0.32 mg sodium hydroxide, and Water for Injection. Hydrochloric acid and sodium hydroxide may have been added to adjust pH.
Prior to intravenous infusion, Remodulin® may be diluted.
See Package Insert for Dosage and Administration.
United Therapeutics Corp., RTP, NC USA 27709



REMODULIN®
(treprostinil) Injection

50 mg/20 mL
(2.5 mg/mL)

For Subcutaneous
and Intravenous Infusion

20 mL multidose vial

**United
Therapeutics**

NDC 66302-102-01
Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).
Keep out of reach of children.

Rx Only 10051 Rev. 01/2023

LOT XXXXXXXX
EXP MM-YYYY

(b) (4)

Contents: 5 mg treprostinil per mL.
Each mL also contains 6.3 mg sodium citrate dihydrate, 5.3 mg sodium chloride, 3 mg metacresol, 0.62 mg sodium hydroxide, and Water for Injection. Hydrochloric acid and sodium hydroxide may have been added to adjust pH.
Prior to intravenous infusion, Remodulin® may be diluted.
See Package Insert for Dosage and Administration.

United Therapeutics Corp., RTP, NC USA 27709



REMODULIN®
(treprostinil) Injection
100 mg/20 mL
(5 mg/mL)

For Subcutaneous and Intravenous Infusion

20 mL multidose vial



NDC 68302-105-01

Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).

Keep out of reach of children.

Rx Only 100852 Rev. 01/2023

LOT XXXXXXXX

EXP MM-YYYY

(b) (4)

Contents: 10 mg treprostinil per mL.
Each mL also contains 6.3 mg sodium citrate
dihydrate, 4 mg sodium chloride, 3 mg
metacresol, 1.2 mg sodium hydroxide, and
Water for Injection. Hydrochloric acid and sodium
hydroxide may have been added to adjust pH.
Prior to intravenous infusion, Remodu i[®] may
be diluted.

See Package Inset for Dosage and Administration.


United Therapeutics Corp., RTP, NC USA 27709



REMODULIN[®]
(treprostinil) Injection
200 mg/20 mL
(10 mg/mL)

For Subcutaneous
and Intravenous Infusion

20 mL multidose vial



NDC 66302-110-01

Store at 25°C (77°F) with excursions
permitted to 2-30°C (36-86°F).

Keep out of reach of children.

Rx Only 10053 Rev. 01/2023

LOT XXXXXXX

EXP MM-YYYY

(b) (4)

Contents: 20 mg treprostinil per mL.
Each mL also contains 6.3 mg sodium citrate dihydrate, 4 mg sodium chloride, 3 mg metacresol, 2.25 mg sodium hydroxide, and Water for Injection. Hydrochloric acid and sodium hydroxide may have been added to adjust pH.

Prior to intravenous infusion, Remodu ^{tr} may be diluted.

See Package Insert for Dosage and Administration.
United Therapeutics Corp., RTP, NC USA 27709




REMODULIN^{tr}
(treprostinil) Injection

400 mg/20 mL
(20 mg/mL)

For Subcutaneous
and Intravenous Infusion

20 mL multidose vial



NDC 68302-120-01

Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).


Keep out of reach of children.

Rx Only 101749 Rev. 01/2023


LOT XXXXXXX

EXP MM-YYYY

(b) (4)



(b) (4)




2D
Barcode
Space


LOT XXXXXX
EXP MM-YYYY
SN XXXXXXXXXXXXX
GTIN XXXXXXXXXXXXX

MADE IN USA

(b) (4)



(b) (4)



REMODULIN[®]
(treprostinil) Injection

**200 mg/20 mL
(10 mg/mL)**

**For Subcutaneous
and Intravenous Infusion**

20 mL multidose vial



NDC 66302-110-01

Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).

Keep out of reach of children.

Rx Only



REMODULIN[®]
(treprostinil) Injection

**200 mg/20 mL
(10 mg/mL)**

**For Subcutaneous
and Intravenous Infusion**

20 mL multidose vial




X5346819

2D Barcode Space
LOT XXXXXX
EXP MM-YYYY
SN XXXXXXXXXXXXX
GTIN XXXXXXXXXXXXX

MADE IN USA

(b) (4)





(b) (4)

