







Reference ID: 5277817





Reference ID: 5277817



Drug Facts (continued)	Drug Facts (continued)	Drug Facts (continued)	Drug Facts (continued)	Drug Facts (continued)
Uses Temporarily relieves these symptoms of hay fever or other upper respiratory allergies: • nasal congestion • runny nose • sneezing • lichy nose • lichy, watery eyes (for ages 12 and older)	Ask a doctor or pharmacist before use if you are taking • medicine for HIV infection (such as ritonavir) • a steroid medicine for asthma, allergies or skin rash • ketoconazole pills (medicine for fungal infection) When using this product • the growth rate of some children may be slower • stinging or snezzing may occur for a few seconds right after use • do not share this bottle with anyone else as this may spread germs • remember to tell your doctor about all the medicines you take, including this one	 your symptoms do not get better within 7 days of starting use or you get new symptoms such as severe facial pain or thick nasal discharge, You may have something more than allergies, such as an infection, you get a constant whistling sound from your nose. This may be a sign of damage inside your nose, you get an allergic reaction to this product, Seek medical help right away, you get new changes to your vision that develop after starting this product you have severe or frequent nosebleeds If pregnant or breast-feeding, ask a health professional before use, 	Directions - read the Quick Start Guide for how to: - shake vigorously before each use - prime the bottle - use the spray - snift gently after each spray - clean the spray nozzle with a clean dry tissue - use this product only once a day - do not use more than directed ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER • Week 1 – use 2 sprays in each nostril once daily • Week 2 through 6 months – use 1 or 2 sprays in each nostril once daily, as	CHILDREN 2 TO 11 YEARS OF AGE • the growth rate of some children may be slower while using this product. Children should use for the shortest amount of time necessary to achieve symptom relief. Talk to your child's doctor if your child to needs to use the spray for longer than two months a year. • an adult should supervise use • use 1 spray in each nostril once daily CHILDREN UNDER 2 YEARS OF AGE • do not use Other information • you may start to feel relief the first day
Warnings Only for use in the nose. Do not spray into your eyes or mouth.				
Do not use • in children under 2 years of age • to treat asthma • if you have an injury or surgery to your nose that is not fully healed				
if you have ever had an allergic reaction to this product or any of the ingredients Ask a doctor before use if you have or	Stop use and ask a doctor if you have, or come into contact with someone who has, chicken pox, measles	Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away,	needed to treat your symptoms After 6 months of daily use – ask your doctor if you can keep using	and full effect after several days of regular, once-a-day use • store between 15° - 30°C (59° - 86°F) • do not refrigerate or freeze







Relief can be used with most non-prescription and prescription medicines. However, there are a few medicines to look out for because they may cause the level of FLONASE SENSIMIST in your body to come too high. Just to be safe, check this table to see if you're *for ages 12 and older

effect"?
No, FLONASE SENSIMIST Allergy Relief does not cause a
rebound effect. Some nasal decongestant sprays may cause your
hasal passages to swell up even more when you use them too often o
or longer than their label says you should (three days). This is
sometimes called a "rebound effect". FLONASE SENSIMIST is a
different kind of medicine and does not cause any rebound effect. You
an use FLONASE SENSIMIST daily for up to six months if you are ag
2 or older or up to two months a year if you are age 2-11 before
healing with a deater

that came in the		
		or tuberculosis or come inte
	omeone who does?	
	INASE SENSIMIST Aller	gy Relief and ask your docto
		ck nasal discharge, or think you
		ONASE SENSIMIST Allergy
		may want to consider if other
medicines are ne	eded.	,
Does FLONAS	E SENSIMIST Allergy R	elief have side effects?
through your bo can cause side e have been repor	dy. However, like all medi ffects in some people. Be	, and barely any of it travels cines, FLONASE SENSIMIST low are some side effects that E SENSIMIST. If you have any r doctor.
Side effect	What could happen	What to do
Allergic reaction to the product	Sudden swelling of face or tongue • A rash Wheezing or feeling faint	If you feel any of these symptoms, stop using FLONASE SENSIMIST and see a doctor right away.
Nose injury	Severe or frequent nosebleeds	Apply pressure to your nose. Stop using FLONASE SENSIMIST and see a doctor.
	Constant whistling sound that does not go away	This could be a sign of damage to you nose. Stop using FLONASE SENSIMI and see a doctor right away.
Eye conditions such as cataracts or glaucoma	Cloudy vision Reduced vision	Have a yearly eye exam to check for these conditions. Read the <i>Drug Facts</i> or the back of the FLONASE SENSIMIST package for more details.
Growth effects	Slower growth rate in some children while using product	Talk to your doctor if you are concerned or if a child age 2-11 need to use FLONASE SENSIMIST for longer than two months a year.
Other side effects	Headaches or sneezing Minor nosebleeds Dry or irritated nose or throat	Talk to your doctor or pharmacist if you are concerned.

 \mathcal{P}

1-844-FLONASE

(1-844-356-6273)

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD 11/15/2023 01:48:02 PM