

Syringe Assembly Directions:
See User Guide

USE ASEPTIC TECHNIQUE
Do not assemble until ready to use

1. Remove protective caps from vial and injector.
2. Align vial such that the injector needle is centered on the stopper. Thread vial into injector. 3 half turns, this will allow the needle to penetrate the center of the stopper.*



DO NOT PUSH VIAL INTO INJECTOR: THIS MAY CAUSE MISALIGNMENT.



3. Remove needle cover pulling straight up. Expel air.
- 4a) After injection, push tab forward with thumb to flip shield toward needle.
- 4b) With a FIRM, QUICK motion, press down against a flat surface until an audible "click" is heard. This sound indicates shield activation.



*CAUTION: IMPROPER ENGAGING MAY CAUSE GLASS BREAKAGE AND SUBSEQUENT INJURY.

5612400M
8-23

OPEN HERE



(01) 00376329124018

PHYTONADIONE INJECTABLE EMULSION, USP
1 mg/0.5 mL

SINGLE DOSE / NO PRESERVATIVE ADDED / PROTECT FROM LIGHT / STORE AT 20° TO 25°C (68° TO 77°F) EXCURSIONS PERMITTED TO 15° TO 30°C (59° TO 86°F) [SEE USP CONTROLLED ROOM TEMPERATURE . . / FOR INTRAVENOUS, INTRAMUSCULAR OR SUBCUTANEOUS USE (WITH CAUTION) / USUAL DOSAGE SEE INSERT

NDC 76329-1240-1 STOCK NO. 1240

▲ open

Rx Only SAF-T-JET® 27 G. X 1/2" NEEDLE



SAF-T-JET®

PHYTONADIONE INJECTABLE EMULSION USP
1 mg/0.5 mL
NEONATAL CONCENTRATION

Single Dose Discard unused portion
FOR INTRAVENOUS, INTRAMUSCULAR OR SUBCUTANEOUS USE (WITH CAUTION)

170

LOT / EXP.

NON-VARNISH AREA

Each 0.5 mL contains 1 mg phytonadione (Vitamin K1); 10 mg polysorbate 80 10.4 mg propylene glycol 0.17 mg sodium acetate anhydrous 0.00002 mL glacial acetic acid Additional acetic acid or sodium acetate anhydrous may have been added to adjust pH to meet USP limits of 3.5 to 7.0 Protect from light until ready to use Medication and fluid pathway sterile and nonpyrogenic in original, unopened package, with component caps in place Do not remove caps until ready to use



INTERNATIONAL MEDICATION SYSTEMS, LIMITED
So. El Monte, CA 91733, U.S.A.
An Amphastar Pharmaceuticals Company



(b) (4)



BARCODE 3 OF 9

6912400P

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PHYTONADIONE INJECTABLE EMULSION safely and effectively. See full prescribing information for PHYTONADIONE INJECTABLE EMULSION.

PHYTONADIONE Injection, for intravenous, intramuscular, and subcutaneous use.

Initial U.S. Approval: 1960

WARNING – HYPERSENSITIVITY REACTIONS WITH INTRAVENOUS AND INTRAMUSCULAR USE

See full prescribing information for complete boxed warning.

Fatal hypersensitivity reactions, including anaphylaxis, have occurred during and immediately after INTRAVENOUS and INTRAMUSCULAR injection of Phytonadione Injectable Emulsion. Reactions have occurred despite dilution to avoid rapid infusion and upon first and subsequent doses. Avoid the intravenous and intramuscular routes of administration unless the subcutaneous route is not feasible and the serious risk is justified (5.1)

INDICATIONS AND USAGE

Phytonadione Injectable Emulsion is a vitamin K replacement indicated for the treatment of the following coagulation disorders which are due to faulty formation of factors II, VII, IX and X when caused by vitamin K deficiency or interference with vitamin K activity.

- Anticoagulant-induced hypoprothrombinemia deficiency caused by coumarin or indanedione derivatives; (1.1)
- Hypoprothrombinemia due to antibacterial therapy; (1.1)
- Hypoprothrombinemia secondary to factors limiting absorption or synthesis of vitamin K, e.g., obstructive jaundice, biliary fistula, sprue, ulcerative colitis, celiac disease, intestinal resection, cystic fibrosis of the pancreas, and regional enteritis; (1.1)

FULL PRESCRIBING INFORMATION: CONTENTS *

WARNING: HYPERSENSITIVITY REACTIONS WITH INTRAVENOUS AND INTRAMUSCULAR USE

1 INDICATIONS & USAGE

- 1.1 Treatment of Hypoprothrombinemia Due to Vitamin K Deficiency or Interference
- 1.2 Prophylaxis and Treatment of Vitamin K-Deficiency Bleeding in Neonates

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosing Considerations
- 2.2 Recommended Dosage for Coagulation Disorders from Vitamin K Deficiency of Interference
- 2.3 Recommended Dosage for Prophylaxis and Treatment of Vitamin K Deficiency Bleeding in Neonates
- 2.4 Directions for Dilution

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Hypersensitivity Reactions
- 5.3 Cutaneous Reactions

FULL PRESCRIBING INFORMATION

WARNING — HYPERSENSITIVITY REACTIONS WITH INTRAVENOUS AND INTRAMUSCULAR USE

Fatal hypersensitivity reactions, including anaphylaxis, have occurred during and immediately after intravenous and intramuscular injection of Phytonadione Injectable Emulsion. Reactions have occurred despite dilution to avoid rapid intravenous infusion and upon first dose. Avoid the intravenous and intramuscular routes of administration unless the subcutaneous route is not feasible and the serious risk is justified [see WARNINGS AND PRECAUTIONS (5.1)].

1 INDICATIONS AND USAGE

1.1 Treatment of Hypoprothrombinemia Due to Vitamin K Deficiency or Interference

Phytonadione Injectable Emulsion is indicated for the treatment of the following coagulation disorders which are due to faulty formation of factors II, VII, IX and X when caused by vitamin K-deficiency or interference with vitamin K activity:

- anticoagulant-induced hypoprothrombinemia caused by coumarin or indanedione derivatives;
- hypoprothrombinemia due to antibacterial therapy;
- hypoprothrombinemia secondary to factors limiting absorption or synthesis of vitamin K, e.g., obstructive jaundice, biliary fistula, sprue, ulcerative colitis, celiac disease, intestinal resection, cystic fibrosis of the pancreas, and regional enteritis;
- other drug-induced hypoprothrombinemia where it is definitely shown that the result is due to interference with vitamin K metabolism, e.g., salicylates.

1.2 Prophylaxis and Treatment of Vitamin K-Deficiency Bleeding in Neonates

Phytonadione Injectable Emulsion is indicated for prophylaxis and treatment of vitamin K-deficiency bleeding in neonates.

2 DOSAGE AND ADMINISTRATION

2.1 Dosing Considerations

Whenever possible, administer Phytonadione Injectable Emulsion by the subcutaneous route [see *Boxed Warning*]. When intravenous administration is unavoidable, inject the drug very slowly, not exceeding 1 mg per minute [see WARNINGS AND PRECAUTIONS (5.1)].

Monitor international normalized ratio (INR) regularly and as clinical conditions indicate. Use the lowest effective dose of Phytonadione Injectable Emulsion.

The coagulant effects of Phytonadione Injectable Emulsion are not immediate; improvement of INR may take 1-8 hours. Interim use of whole blood or component therapy may also be necessary if bleeding is severe.

When Phytonadione Injectable Emulsion is used to correct excessive anticoagulant-induced hypoprothrombinemia, anticoagulant therapy still being indicated, the patient is again faced with the clotting hazards existing prior to starting the anticoagulant therapy. Phytonadione Injectable

- Other drug-induced hypoprothrombinemia where it is definitely shown that the result is due to interference with vitamin K metabolism, e.g., salicylates. (1.1)

Phytonadione Injectable Emulsion is indicated for prophylaxis and treatment of vitamin K-deficiency bleeding in neonates. (1.2)

DOSAGE AND ADMINISTRATION

- Administer Phytonadione Injectable Emulsion by the subcutaneous route, whenever possible. (2.1)
- When intravenous administration is unavoidable, inject the drug very slowly, not exceeding 1 mg per minute. (2.1)

DOSAGE FORMS AND STRENGTHS

Injection: 1 mg/0.5 mL single-dose vial and a SAF-T-Jet® vial injector. (3)

CONTRAINDICATIONS

Hypersensitivity to any component of this medication. (4)

WARNINGS AND PRECAUTIONS

- Cutaneous Reactions: May occur with parenteral use. Discontinue drug and manage medically. (5.3)

ADVERSE REACTIONS

Most common adverse reactions are cyanosis, diaphoresis, dizziness, dysgeusia, dyspnea, flushing, hypotension and tachycardia. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Amphastar Pharmaceuticals, Inc. at 1-800-423-4136, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Anticoagulants: May induce temporary resistance to prothrombin-depressing anticoagulants. (7)

USE IN SPECIFIC POPULATIONS

- Pregnancy: If available, use the preservative-free formulation in pregnant women. (8.1)
- Lactation: If available, use the preservative-free formulation in lactating women. (8.2)
- Pediatric Use: The safety and effectiveness of Phytonadione Injectable Emulsion in pediatric patients from 6 months to 17 years have not been established. (8.4)

Revised: 08/2023

See 17 for PATIENT COUNSELING INFORMATION

6 ADVERSE REACTIONS

6.1 Clinical Trials and Post-Marketing experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY SECTION

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed

Emulsion is not a clotting agent, but overzealous therapy with Phytonadione Injectable Emulsion may restore conditions which originally permitted thromboembolic phenomena. Dosage should be kept as low as possible, and INR should be checked regularly as clinical conditions indicate.

2.2 Recommended Dosage for Coagulation Disorders from Vitamin K Deficiency or Interference

The recommended dosage of Phytonadione Injectable Emulsion is based on whether the hypoprothrombinemia is anticoagulant-induced (e.g., due to coumarin or indanedione derivatives) or non-anticoagulant-induced (e.g., due to antibiotics; salicylates or other drugs; factors limiting absorption or synthesis) as follows:

- **Anticoagulant-Induced Hypoprothrombinemia:** Phytonadione Injectable Emulsion 2.5 mg to 10 mg or more subcutaneously, intramuscularly, or intravenously. Up to 25 mg to 50 mg may be administered as a single dose.

Repeated large doses of Phytonadione Injectable Emulsion are not warranted in liver disease if the initial response is unsatisfactory. Failure to respond to Phytonadione Injectable Emulsion may indicate that the condition being treated is inherently unresponsive to Phytonadione Injectable Emulsion.

- **Hypoprothrombinemia Due to Other Causes (Non-Anticoagulation-Induced Hypoprothrombinemia):** Phytonadione Injectable Emulsion 2.5 mg to 25 mg or more intravenously, intramuscularly, or subcutaneously. Up to 50 mg may be administered as a single dose.

Evaluate INR after 6-8 hours, and repeat dose if INR remains prolonged. Modify subsequent dosage (amount and frequency) based on the INR or clinical condition.

2.3 Recommended Dosage for Prophylaxis and Treatment of Vitamin K Deficiency Bleeding in Neonates

Prophylaxis of Vitamin K-Deficiency Bleeding in Neonates

The recommended dosage of Phytonadione Injectable Emulsion is 0.5 mg to 1 mg within one hour of birth for a single dose.

Treatment of Vitamin K-Deficiency Bleeding in Neonates

The recommended dosage of Phytonadione Injectable Emulsion is 1 mg given either subcutaneously or intramuscularly.

Consider higher doses if the mother has been receiving oral anticoagulants.

A failure to respond (shortening of the INR in 2 to 4 hours) may indicate another diagnosis or coagulation disorder.

2.4 Directions for Dilution

Dilute Phytonadione Injectable Emulsion with 0.9% Sodium Chloride Injection, 5% Dextrose Injection, or 5% Dextrose and Sodium Chloride Injection.

When diluted, start administration of Phytonadione Injectable Emulsion immediately after dilution.

Discard unused portions of diluted solution as well as unused contents of the vial.

Protect Phytonadione Injectable Emulsion from light at all times.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

3 DOSAGE FORMS AND STRENGTHS

Injection: 1 mg/0.5 mL single-dose vial and a SAF-T-Jet® vial injector.

4 CONTRAINDICATIONS

Hypersensitivity to phytonadione or any other component of this medication [see WARNINGS AND PRECAUTIONS (5.1)].

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

Fatal and severe hypersensitivity reactions, including anaphylaxis, have occurred with intravenous or intramuscular administration of Phytonadione Injectable Emulsion. Reactions have occurred despite dilution to avoid rapid intravenous infusion and upon first dose. These reactions have included shock, cardiorespiratory arrest, flushing, diaphoresis, chest pain, tachycardia, cyanosis, weakness, and dyspnea. Administer Phytonadione Injectable Emulsion subcutaneously whenever feasible. Avoid the intravenous and intramuscular routes of administration unless the subcutaneous route is not feasible and the serious risk is justified [see DOSAGE AND ADMINISTRATION (2.1)].

5.3 Cutaneous Reactions

Parenteral administration of vitamin K replacements (including Phytonadione Injectable Emulsion) may cause cutaneous reactions. Reactions have included eczematous reactions, scleroderma-like patches, urticaria, and delayed-type hypersensitivity reactions. Time of onset ranged from 1 day to a year after parenteral administration. Discontinue Phytonadione Injectable Emulsion for skin reactions and institute medical management.

6 ADVERSE REACTIONS

The following serious adverse reactions are described elsewhere in the labeling:

- Hypersensitivity Reactions [see WARNINGS AND PRECAUTIONS (5.1)]
- Cutaneous Reactions [see WARNINGS AND PRECAUTIONS (5.3)]

6.1 Clinical Trials and Post-Marketing Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The following adverse reactions have been identified during post-approval use of Phytonadione Injectable Emulsion. Because these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Cardiac Disorders: Tachycardia, hypotension.

General disorders and administration site conditions: Generalized flushing; pain, swelling, and tenderness at injection site.

Hepatobiliary Disorders: Hyperbilirubinemia

Immune System Disorders: Fatal hypersensitivity reactions, anaphylactic reactions.

Neurologic: Dysgeusia, dizziness.

Pulmonary: Dyspnea.

Skin and Subcutaneous Tissue Disorders: Erythema, pruritic plaques, scleroderma-like lesions, erythema perstans.

Vascular: Cyanosis.

7 DRUG INTERACTIONS

Anticoagulants

Phytonadione Injectable Emulsion may induce temporary resistance to prothrombin-depressing anticoagulants, especially when larger doses of Phytonadione Injectable Emulsion are used. Should this occur, higher doses of anticoagulant therapy may be needed when resuming anticoagulant therapy, or a change in therapy to a different class of anticoagulant may be necessary (i.e., heparin sodium).

Phytonadione Injectable Emulsion does not affect the anticoagulant action of heparin.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

If Phytonadione is needed during pregnancy, consider using a preservative-free formulation.

Published studies with the use of phytonadione during pregnancy have not reported a clear association with phytonadione and adverse developmental outcomes (see *Data*). There are maternal and fetal risks associated with vitamin K deficiency during pregnancy (see *Clinical Considerations*). Animal reproduction studies have not been conducted with phytonadione.

The estimated background risk for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Clinical Considerations

Disease-associated maternal and/or embryo/fetal risk

Pregnant women with vitamin K deficiency hypoprothrombinemia may be at an increased risk for bleeding diatheses during pregnancy and hemorrhagic events at delivery. Subclinical maternal vitamin K deficiency during pregnancy has been implicated in rare cases of fetal intracranial hemorrhage.

Data

Human Data

Phytonadione has been measured in cord blood of infants whose mothers were treated with phytonadione during pregnancy in concentrations lower than seen in maternal plasma. Administration of vitamin K₁ to pregnant women shortly before delivery increased both maternal and cord blood concentrations. Published data do not report a clear association with phytonadione and adverse maternal or fetal outcomes when used during pregnancy. However, these studies cannot definitively establish the absence of any risk because of methodologic limitations including small sample size and lack of blinding.

Animal Data

In pregnant rats receiving vitamin K₁ orally, fetal plasma and liver concentrations increased following administration, supporting placental transfer.

8.2 Lactation

Risk Summary

If available, preservative-free Phytonadione is recommended when Phytonadione is needed during lactation [see *USE IN SPECIFIC POPULATIONS* (8.4)].

Phytonadione is present in breastmilk. There are no data on the effects of Phytonadione Injectable Emulsion on the breastfed child or on milk production. The developmental and health benefits of breastfeeding should be considered along with the clinical need for Phytonadione Injectable Emulsion and any potential adverse effects on the breastfed child from Phytonadione Injectable Emulsion or from the underlying maternal condition.

8.4 Pediatric Use

The safety and effectiveness of Phytionadione Injectable Emulsion for prophylaxis and treatment of vitamin K deficiency have been established in neonates. Use of phytionadione injection for prophylaxis and treatment of vitamin K deficiency is based on published clinical studies.


10 OVERDOSAGE

Hemolysis, jaundice, and hyperbilirubinemia in newborns, particularly in premature infants, may result from Phytionadione Injectable Emulsion overdose.

11 DESCRIPTION

Phytionadione is a vitamin K replacement, which is a clear, yellow to amber, viscous, odorless or nearly odorless liquid. It is insoluble in water, soluble in chloroform and slightly soluble in ethanol. It has a molecular weight of 450.70.

Phytionadione is 2-methyl-3-phytyl-1, 4-naphthoquinone. Its empirical formula is $C_{31}H_{46}O_2$ and its molecular structure is:



Phytionadione Injectable Emulsion, USP injection is a yellow, sterile, aqueous colloidal solution of vitamin K_1 , with a pH of 3.5 to 7.0, available for injection by the intravenous, intramuscular, and subcutaneous routes. Phytionadione Injectable Emulsion, USP is available in 1 mg (1 mg/0.5 mL) single-dose vials. Each 0.5 mL of Phytionadione Injectable Emulsion, USP contains the following inactive ingredients: 10 mg polysorbate 80, 10.4 mg propylene glycol, 0.17 mg sodium acetate anhydrous, and 0.00002 mL glacial acetic acid. Additional glacial acetic acid or sodium acetate anhydrous may have been added to adjust pH to meet USP limits of 3.5 to 7.0.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Phytionadione Injectable Emulsion aqueous colloidal solution of vitamin K_1 for parenteral injection, possesses the same type and degree of activity as does naturally-occurring vitamin K, which is necessary for the production via the liver of active prothrombin (factor II), proconvertin (factor VII), plasma thromboplastin component (factor IX), and Stuart factor (factor X). Vitamin K is an essential cofactor for a microsomal enzyme that catalyzes the post-translational carboxylation of multiple, specific, peptide-bound glutamic acid residues in inactive hepatic precursors of factors II, VII, IX, and X. The resulting gamma-carboxy-glutamic acid residues convert the precursors into active coagulation factors that are subsequently secreted by liver cells into the blood.

In normal animals and humans, phytionadione is virtually devoid of activity. However, in animals and humans deficient in vitamin K, the pharmacological action of vitamin K is related to its normal physiological function, that is, to promote the hepatic biosynthesis of vitamin K dependent clotting factors.

12.2 Pharmacodynamics

The action of the aqueous dispersion, when administered intravenously, is generally detectable within an hour or two and hemorrhage is usually controlled within 3 to 6 hours. A normal INR may often be obtained in 12 to 14 hours.

12.3 Pharmacokinetics

Absorption:

Phytionadione is readily absorbed following intramuscular administration.

Distribution:

After absorption, phytionadione is initially concentrated in the liver, but the concentration declines rapidly. Very little vitamin K accumulates in tissues.

Elimination:

Little is known about the metabolic fate of vitamin K. Almost no free unmetabolized vitamin K appears in bile or urine.

13 NONCLINICAL TOXICOLOGY SECTION

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Studies of carcinogenicity, genotoxicity or impairment of fertility have not been conducted with phytionadione.

16 HOW SUPPLIED/STORAGE AND HANDLING

Phytionadione Injectable Emulsion, USP is a yellow, sterile, aqueous colloidal solution and is supplied in unit use packages containing one single-dose vial and a SAF-T-Jet[®] vial injector, 27 G. x 1/2" needle.

Phytionadione Injectable Emulsion USP, 1 mg in 0.5 mL

Stock No. 1240 NDC 76329-1240-1

Store at 20 to 25 C (68 to 77 F); excursions permitted to 15 to 30 C (59 to 86 F) [see USP Controlled Room Temperature].

Protect Phytionadione Injectable Emulsion, USP from light. Store container in closed original carton until contents have been used.

Syringe Assembly Directions:

See User Guide


USE ASEPTIC TECHNIQUE

Do not remove from carton or assemble until ready to use.

- 1) Remove protective caps from vial and injector.



- 3) a. Flip shield down.
b. Remove needle cover PULLING STRAIGHT UP (DO NOT TWIST). Expel air.



- 2) Align vial such that the injector needle is centered on the stopper.

Thread vial into injector 3 half turns; this will allow the needle to penetrate the stopper.*



DO NOT PUSH VIAL INTO INJECTOR; THIS MAY CAUSE MISALIGNMENT OF THE INJECTOR NEEDLE.

- 4) a. After injection, push tab forward with thumb to flip shield toward needle.

With a FIRM, QUICK motion, press down against a flat surface until an audible "click" is heard. This sound indicates shield activation. Note: Audible click may not be heard on small needle sizes: visual confirmation is required.



*CAUTION: IMPROPER ENGAGING MAY CAUSE GLASS BREAKAGE AND SUBSEQUENT INJURY.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Inform the patient of the following important risks of Phytionadione Injectable Emulsion:

Serious Hypersensitivity Reactions

Advise the patient and caregivers to immediately report signs of hypersensitivity after receiving Phytionadione Injectable Emulsion [see WARNINGS AND PRECAUTIONS (5.1)].

Cutaneous Reactions

Advise the patient and caregivers to report the occurrence of new rashes after receiving Phytionadione Injectable Emulsion. These reactions may be delayed for up to a year after treatment [see WARNINGS AND PRECAUTIONS (5.3)].

Rx Only

INTERNATIONAL MEDICATION SYSTEMS, LIMITED
So. El Monte, CA 91733 U.S.A.
An Amphastar Pharmaceuticals Company
© INTERNATIONAL MEDICATION SYSTEMS, LIMITED 2023

Rev. 8-23

(b) (4)