

NARCAN[®]

Naloxone HCl Nasal Spray 4 mg




Drug Facts (continued)

Directions




- Step 1: CHECK if you suspect an overdose:**
- CHECK** for a suspected overdose: the person will not wake up or is very sleepy or not breathing well
 - yell "Wake up!"
 - shake the person gently
 - if the person is not awake, go to Step 2



- Step 2: GIVE 1st dose in the nose**
- HOLD** the nasal spray device with your thumb on the bottom of the plunger
 - INSERT** the nozzle into either NOSTRIL
 - PRESS** the plunger firmly to give the 1st dose
 - 1 nasal spray device contains 1 dose



- Step 3: CALL**
- CALL 911** immediately after giving the 1st dose



- Step 4: WATCH & GIVE**
- WAIT** 2-3 minutes after the 1st dose to give the medicine time to work
 - if the person **wakes up**: Go to Step 5
 - if the person does **not wake up**:
 - CONTINUE TO GIVE** doses every 2-3 minutes until the person wakes up
 - it is safe to keep giving doses



- Step 5: STAY**
- STAY** until ambulance arrives: even if the person wakes up
 - GIVE** another dose if the person becomes very sleepy again
 - You may need to give all the doses in the pack

Drug Facts (continued)

Warning

When using this product some people may experience symptoms when they wake up, such as shaking, sweating, nausea, or feeling angry. This is to be expected.

Other information

- store at room temperature or refrigerated, between 2°C to 25°C (36°F to 77°F)
- do not freeze
- avoid excessive heat above 40°C (104°F)
- protect from light
- the product is packaged in individually-sealed blisters. Do not use if the blister is open or torn, or if the device appears damaged.

Inactive ingredients

benzalkonium chloride, edetate disodium, hydrochloric acid, sodium chloride, water

Questions?

call 1-844-4NARCAN (1-844-462-7226) or go to www.narcan.com

Safe to Use
Even if Opioids are Not Present

Distributed by Emergent Devices Inc.
Plymouth Meeting, PA 19462 USA

©2023 Emergent Devices Inc. Plymouth Meeting, PA.
All rights reserved.

NARCAN[®] is a registered trademark of Emergent Operations Ireland Limited.

NDC 69547-627-02



Original
Prescription
Strength

Easy to Use

Can Save a Life

NARCAN[®]

Naloxone HCl Nasal Spray 4 mg

Emergency Treatment of Opioid Overdose


Designed to
Rapidly Reverse
the Effects of a
Life-Threatening
Opioid Emergency



2 Single-Dose Nasal Spray Devices
0.003 fl oz (0.1 mL) each

For use in
nose only

For use in nose only



Do not test nasal
spray device
before use

1 nasal spray
device contains
1 dose of medicine

Each device
sprays 1 time only

Read Directions Before Use ▶

Drug Facts

Active ingredient (in each spray)	Purpose
Naloxone hydrochloride 4 mg	Emergency treatment of opioid overdose

Uses

- to "revive" someone during an overdose from many prescription pain medications or street drugs such as heroin
- this medicine can save a life

LOT XXXXXX

EXP YYYY-MM-DD



XXXXX
XXXXXX


NARCAN®

Naloxone HCl Nasal Spray 4 mg



Drug Facts (continued)

Directions



Step 1: CHECK if you suspect an overdose:

- **CHECK** for a suspected overdose: the person will not wake up or is very sleepy or not breathing well
- yell "Wake up!"
- shake the person gently
- if the person is not awake, go to Step 2

Step 2: GIVE 1st dose in the nose

- **HOLD** the nasal spray device with your thumb on the bottom of the plunger
- **INSERT** the nozzle into either NOSTRIL
- **PRESS** the plunger firmly to give the 1st dose
- 1 nasal spray device contains 1 dose

Step 3: CALL

- **CALL 911** immediately after giving the 1st dose

Step 4: WATCH & GIVE

- **WAIT** 2-3 minutes after the 1st dose to give the medicine time to work
- if the person **wakes up**: Go to Step 5
- if the person does **not wake up**:
 - **CONTINUE TO GIVE** doses every 2-3 minutes until the person wakes up
 - it is safe to keep giving doses

Step 5: STAY

- **STAY** until ambulance arrives: even if the person wakes up
- **GIVE** another dose if the person becomes very sleepy again
- You may need to give all the doses in the pack

Drug Facts (continued)

Warning

When using this product some people may experience symptoms when they wake up, such as shaking, sweating, nausea, or feeling angry. This is to be expected.

Other information

- store at room temperature or refrigerated, between 2°C to 25°C (36°F to 77°F)
- do not freeze
- avoid excessive heat above 40°C (104°F)
- protect from light
- the product is packaged in individually-sealed blisters. Do not use if the blister is open or torn, or if the device appears damaged.

Inactive ingredients

benzalkonium chloride, edetate disodium, hydrochloric acid, sodium chloride, water

Questions?

call 1-844-4NARCAN (1-844-462-7226) or go to www.narcan.com

Distributed by Emergent Devices Inc.
Plymouth Meeting, PA 19462 USA

©2023 Emergent Devices Inc. Plymouth Meeting, PA.
All rights reserved.

NARCAN® is a registered trademark of Emergent Operations Ireland Limited.

NDC 69547-627-02



NOT FOR RESALE

FOR NON-RETAIL DISTRIBUTION

NARCAN®

Naloxone HCl Nasal Spray 4 mg

Emergency Treatment of Opioid Overdose


Designed to Rapidly Reverse the Effects of a Life-Threatening Opioid Emergency



2 Single-Dose Nasal Spray Devices
0.003 fl oz (0.1 mL) each

For use in nose only

For use in nose only



Nozzle

Plunger

Do not test nasal spray device before use

1 nasal spray device contains 1 dose of medicine

Each device sprays 1 time only

Read Directions Before Use ▶

Drug Facts

Active ingredient (in each spray)	Purpose
Naloxone hydrochloride 4 mg	Emergency treatment of opioid overdose


Uses

- to "revive" someone during an overdose from many prescription pain medications or street drugs such as heroin
- this medicine can save a life

LOT XXXXXX

EXP YYYY-MM-DD

3 69547-353-025 N



XXXXX
XXXXXX

NDC 69547-627-02



NARCAN[®]
Naloxone HCl Nasal Spray 4 mg
Emergency Treatment of Opioid Overdose



N
3 69547-627-02 7

- For use in the nose only
- Do not test nasal spray device before use
- 1 nasal spray device contains 1 dose of medicine
- Each device sprays 1 time only
- Store at room temperature or refrigerated, between 2°C to 25°C (36°F to 77°F)
- Do not freeze
- Avoid excessive heat above 40°C (104°F)

LOT XXXXXX EXP YYYY-MMM-DD

Distributed by Emergent Devices Inc.
Plymouth Meeting, PA 19462 USA A1160


DIRECTIONS

NARCAN[®]
Naloxone HCl Nasal Spray 4 mg

Emergency Treatment of Opioid Overdose

Important:

- For use in the nose only
- Do not test nasal spray device before use
- 1 nasal spray device contains 1 dose of medicine
- Each device sprays 1 time only



1 CHECK




Step 1: CHECK if you suspect an overdose:

- **CHECK** for a suspected overdose: the person will not wake up or is very sleepy or not breathing well
 - yell "Wake up!"
 - shake the person gently
 - if the person is not awake, go to Step 2

**Unfold for
Directions**

2 GIVE



Step 2: Give 1st dose in the nose

- **HOLD** the nasal spray device with your thumb on the bottom of the plunger
- **INSERT** the nozzle into either NOSTRIL
- **PRESS** the plunger firmly to give the 1st dose
- 1 nasal spray device contains 1 dose

**Unfold for
Directions**


3 CALL



Step 3: Call 911

- **CALL 911** immediately after giving the 1st dose

4 WATCH/GIVE



Step 4: WATCH & GIVE

- **WAIT** 2-3 minutes after the 1st dose to give the medicine time to work
- if the person wakes up: Go to Step 5
- if the person does not wake up:
 - **CONTINUE TO GIVE** doses every 2-3 minutes until the person wakes up
 - it is safe to keep giving doses

5 STAY



Step 5: STAY

- **STAY** until ambulance arrives: even if the person wakes up
- **GIVE** another dose if the person becomes very sleepy again
- You may need to give all the doses in the pack


EMERGENT[®]

For opioid emergencies, call 911. For questions on NARCAN, call 1-844-4NARCAN (1-844-462-7226) or go to www.narcan.com.
©2023 Emergent Devices Inc. EMERGENT[®] and NARCAN[®] are registered trademarks of Emergent BioSolutions Inc, or its subsidiaries.

A1162



FRONT



BACK

