

Instructions for Use
REXTOVY (rex toe' vee)
(Naloxone Hydrochloride)
Nasal Spray

You and your family members or caregivers should read the Instructions for Use that comes with REXTOVY™ Nasal Spray before using it. Talk to your healthcare provider if you and your family members or caregivers have any questions about the use of REXTOVY™ Nasal Spray.

Use REXTOVY™ Nasal Spray for known or suspected opioid overdose in adults and children.

Important: For use in the nose only.

- Do not remove or test the REXTOVY™ Nasal Spray until ready to use.
- Each REXTOVY™ Nasal Spray has 1 dose and cannot be reused.
- You do not need to prime REXTOVY™ Nasal Spray.

How to use REXTOVY™ Nasal Spray:

1



- Lay person flat on their back.



2


- Peel the blister pack open from top left corner with the arrow.
- Remove the REXTOVY™ Nasal Spray.

Note: REXTOVY™ Nasal Spray freezes at temperatures below 5°F (-15°C). If this happens, the device will not spray. Get emergency medical help right away if this happens. Do not wait for REXTOVY™ Nasal Spray to thaw. REXTOVY™ Nasal Spray may still be used if it has been thawed after being previously frozen.



3

- Support their neck with your hand and allow the head to tilt back.
- Gently insert the nozzle **all the way** into one nostril.




4

- Push the green plunger **all the way down until it stops** to give the dose.
- Remove the REXTOVY™ Nasal Spray from nostril after dosing.



5

- Get emergency help right away or call 911.



911

6

- Watch the person for response.
- Wait 2 minutes






Wait and Watch!




7

- If no response, give a 2nd dose with a new REXTOVY™ Nasal Spray in the opposite nostril following the instructions in steps 2 to 4 above.
- If additional REXTOVY™ Nasal Sprays are available, Steps 2 to 4 may be repeated every 2 to 3 minutes, alternating nostrils, until the person responds or emergency help is received.
- If the person responds by waking up to the voice or touch or starts breathing normally, place the person on their side (recovery position).
- Watch the person until emergency help arrives.



8

- Throw away (dispose of) the used REXTOVY™ Nasal Spray in a place that is away from children.



How should I store REXTOVY™ Nasal Spray?

- Store at room temperature between 68°F to 77°F (20°C to 25°C). REXTOVY™ Nasal Spray may be stored for short periods between 39°F to 104°F (4°C to 40°C).
- Do not freeze. Do not expose to temperatures below 39°F (4°C) or above 104°F (40°C).
- Keep REXTOVY™ Nasal Spray in its box until ready to use. Protect from light.
- Replace REXTOVY™ Nasal Spray before the expiration date on the box.

Keep REXTOVY™ Nasal Spray and all medicines out of the reach of children.

Distributed by International Medication Systems, Limited, So. El Monte, CA 91733, U.S.A.
For more information, go to www.amphastar.com or call 1-800-423-4136.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Issued: 06/2023

7036690F/6-23