

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use CIBINQO safely and effectively. See full prescribing information for CIBINQO.

CIBINQO™ (abrocitinib) tablets, for oral use
Initial U.S. Approval: 2022

WARNING: SERIOUS INFECTIONS, MORTALITY, MALIGNANCY, MAJOR ADVERSE CARDIOVASCULAR EVENTS (MACE), and THROMBOSIS

See full prescribing information for complete boxed warning.

- Increased risk of serious bacterial, fungal, viral and opportunistic infections leading to hospitalization or death, including tuberculosis (TB). Discontinue treatment with CIBINQO if serious or opportunistic infection occurs. Test for latent TB before and during therapy; treat latent TB prior to use. Monitor all patients for active TB during treatment, even patients with initial negative latent TB test. (5.1)
- Higher rate of all-cause mortality, including sudden cardiovascular death, with another JAK inhibitor vs. TNF blockers in rheumatoid arthritis (RA) patients. CIBINQO is not approved for use in RA patients. (5.2)
- Malignancies have occurred with CIBINQO. Higher rate of lymphomas and lung cancers with another JAK inhibitor vs. TNF blockers in RA patients. (5.3)
- MACE has occurred with CIBINQO. Higher rate of MACE (defined as cardiovascular death, myocardial infarction, and stroke) with another JAK inhibitor vs. TNF blockers in RA patients. (5.4)
- Thrombosis has occurred with CIBINQO. Increased incidence of pulmonary embolism, venous and arterial thrombosis with another JAK inhibitor vs. TNF blockers. (5.5)

RECENT MAJOR CHANGES

Indications and Usage (1)

02/2023

INDICATIONS AND USAGE

CIBINQO is a Janus kinase (JAK) inhibitor indicated for the treatment of adults and pediatric patients 12 years of age and older with refractory, moderate-to-severe atopic dermatitis whose disease is not adequately controlled with other systemic drug products, including biologics, or when use of those therapies is inadvisable. (1)

Limitation of Use: CIBINQO is not recommended for use in combination with other JAK inhibitors, biologic immunomodulators, or with other immunosuppressants.

DOSAGE AND ADMINISTRATION

- For recommended testing, evaluations and procedures prior to CIBINQO initiation, see Full Prescribing Information. (2.1)
- Recommended dosage is 100 mg orally once daily. (2.2)
- 200 mg orally once daily is recommended for those patients who are not responding to 100 mg once daily. (2.2)
- Moderate renal impairment: 50 mg once daily or 100 mg once daily for those patients who are not responding to 50 mg once daily. (2.3)
- CYP2C19 poor metabolizer: 50 mg once daily or 100 mg once daily for those patients who are not responding to 50 mg once daily. (2.4)
- For dosage modifications for certain adverse reactions, see Full Prescribing Information. (2.6)

DOSAGE FORMS AND STRENGTHS

CIBINQO Tablets: 50 mg, 100 mg, and 200 mg (3)

CONTRAINDICATIONS

Antiplatelet therapies except for low-dose aspirin (≤ 81 mg daily), during the first 3 months of treatment. (4)

WARNINGS AND PRECAUTIONS

- Laboratory Abnormalities: Laboratory monitoring is recommended due to potential changes in platelets, lymphocytes, and lipids. (5.6)
- Immunizations: Avoid use of live vaccines prior to, during, and immediately after CIBINQO treatment. (5.7)

ADVERSE REACTIONS

Most common adverse reactions ($\geq 1\%$) in subjects receiving 100 mg and 200 mg include: nasopharyngitis, nausea, headache, herpes simplex, increased blood creatine phosphokinase, dizziness, urinary tract infection, fatigue, acne, vomiting, oropharyngeal pain, influenza, gastroenteritis. Most common adverse reactions ($\geq 1\%$) in subjects receiving either 100 mg or 200 mg also include: impetigo, hypertension, contact dermatitis, upper abdominal pain, abdominal discomfort, herpes zoster, and thrombocytopenia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Pfizer Inc. at 1-800-438-1985 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Strong inhibitors of CYP2C19: The recommended dose is 50 mg once daily or 100 mg once daily for those patients who are not responding to 50 mg once daily. (2.5, 7.1)
- Moderate to strong inhibitors of both CYP2C19 and CYP2C9, or strong CYP2C19 or CYP2C9 inducers: Avoid concomitant use. (7.1)
- P-gp substrate where small concentration changes may lead to serious or life-threatening toxicities: Monitor or titrate dosage of P-gp substrate. (7.2)

USE IN SPECIFIC POPULATIONS

- Lactation: Breastfeeding not recommended. (8.2)
- Renal Impairment: Avoid use in patients with severe renal impairment or end-stage renal disease. (8.6)
- Hepatic Impairment: Avoid use in patients with severe hepatic impairment. (8.7)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 02/2023

WARNING: SERIOUS INFECTIONS, MORTALITY, MALIGNANCY, MAJOR ADVERSE CARDIOVASCULAR EVENTS, and THROMBOSIS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Testing, Evaluations, and Procedures Prior to Treatment Initiation
- 2.2 Recommended Dosage
- 2.3 Recommended Dosage in Patients with Renal Impairment or Hepatic Impairment
- 2.4 Recommended Dosage in CYP2C19 Poor Metabolizers
- 2.5 Dosage Modifications due to Strong Inhibitors
- 2.6 Treatment Discontinuation due to Serious Infections or Hematologic Adverse Reactions
- 2.7 Administration Instructions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Serious Infections
- 5.2 Mortality
- 5.3 Malignancy and Lymphoproliferative Disorders
- 5.4 Major Adverse Cardiovascular Events
- 5.5 Thrombosis
- 5.6 Laboratory Abnormalities
- 5.7 Immunizations

FULL PRESCRIBING INFORMATION: CONTENTS*

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Effects of Other Drugs on CIBINQO
- 7.2 Effects of CIBINQO on Other Drugs

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment
- 8.8 CYP2C19 Poor Metabolizers

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.5 Pharmacogenomics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: SERIOUS INFECTIONS, MORTALITY, MALIGNANCY, MAJOR ADVERSE CARDIOVASCULAR EVENTS, and THROMBOSIS

Serious Infections

Patients treated with CIBINQO may be at increased risk for developing serious infections that may lead to hospitalization or death; The most frequent serious infections reported with CIBINQO were herpes simplex, herpes zoster, and pneumonia [see *Warnings and Precautions (5.1) and Adverse Reactions (6.1)*].

If a serious or opportunistic infection develops, discontinue CIBINQO and control the infection.

Reported infections from Janus kinase (JAK) inhibitors used to treat inflammatory conditions:

- Active tuberculosis, which may present with pulmonary or extrapulmonary disease. Test for latent TB before and during therapy; treat latent TB prior to use. Monitor all patients for active TB during treatment, even patients with initial negative, latent TB test.
- Invasive fungal infections, including cryptococcosis and pneumocystosis. Patients with invasive fungal infections may present with disseminated, rather than localized, disease.
- Bacterial, viral, including herpes zoster, and other infections due to opportunistic pathogens.

Avoid use of CIBINQO in patients with an active, serious infection including localized infections. The risks and benefits of treatment with CIBINQO should be carefully considered prior to initiating therapy in patients with chronic or recurrent infections.

Patients should be closely monitored for the development of signs and symptoms of infection during and after treatment with CIBINQO, including the possible development of tuberculosis in patients who tested negative for latent tuberculosis infection prior to initiating therapy [see *Warnings and Precautions (5.1)*].

Mortality

In a large, randomized, postmarketing safety study in rheumatoid arthritis (RA) patients 50 years of age and older with at least one cardiovascular risk factor comparing another JAK inhibitor to TNF blocker treatment, a higher rate of all-cause mortality, including sudden cardiovascular death, was observed with the JAK inhibitor. CIBINQO is not approved for use in RA patients [see *Warnings and Precautions (5.2)*].

Malignancies

Malignancies were reported in patients treated with CIBINQO. Lymphoma and other malignancies have been observed in patients receiving JAK inhibitors used to treat inflammatory conditions. In RA patients treated with another JAK inhibitor, a higher rate of malignancies (excluding non-melanoma skin cancer (NMSC)) was observed when compared with TNF blockers. Patients who are current or past smokers are at additional increased risk [see *Warnings and Precautions (5.3)*].

Major Adverse Cardiovascular Events

Major adverse cardiovascular events were reported in patients treated with CIBINQO. In RA patients 50 years of age and older with at least one cardiovascular risk factor treated with another JAK inhibitor, a higher rate of major adverse cardiovascular events (MACE) (defined as cardiovascular death, myocardial infarction, and stroke), was observed when compared with TNF blockers. Patients who are current or past smokers are at additional increased risk. Discontinue

CIBINQO in patients that have experienced a myocardial infarction or stroke [see Warnings and Precautions (5.4)].

Thrombosis

Deep venous thrombosis (DVT) and pulmonary embolism (PE) have been reported in patients treated with CIBINQO. Thrombosis, including PE, DVT, and arterial thrombosis have been reported in patients receiving JAK inhibitors used to treat inflammatory conditions. Many of these adverse reactions were serious and some resulted in death. In RA patients 50 years of age and older with at least one cardiovascular risk factor treated with another JAK inhibitor, a higher rate of thrombosis was observed when compared with TNF blockers. Avoid CIBINQO in patients at risk. If symptoms of thrombosis occur, discontinue CIBINQO and treat appropriately [see Warnings and Precautions (5.5)].

1 INDICATIONS AND USAGE

CIBINQO is indicated for the treatment of adults and pediatric patients 12 years of age and older with refractory, moderate-to-severe atopic dermatitis whose disease is not adequately controlled with other systemic drug products, including biologics, or when use of those therapies is inadvisable.

Limitations of Use

CIBINQO is not recommended for use in combination with other JAK inhibitors, biologic immunomodulators, or with other immunosuppressants.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Testing, Evaluations, and Procedures Prior to Treatment Initiation

Perform the following tests and evaluations prior to CIBINQO initiation:

- Tuberculosis (TB) infection evaluation – CIBINQO initiation is not recommended in patients with active TB. For patients with latent TB or those with a negative latent TB test who are at high risk for TB, start preventive therapy for latent TB prior to initiation of CIBINQO [see Warnings and Precautions (5.1)].
- Viral hepatitis screening in accordance with clinical guidelines – CIBINQO initiation is not recommended in patients with active hepatitis B or hepatitis C [see Warnings and Precautions (5.1)].
- A complete blood count (CBC) – CIBINQO initiation is not recommended in patients with a platelet count $<150,000/\text{mm}^3$, an absolute lymphocyte count $<500/\text{mm}^3$, an absolute neutrophil count $<1,000/\text{mm}^3$, or a hemoglobin value $<8 \text{ g/dL}$ [see Warnings and Precautions (5.6)].

Complete any necessary immunizations, including herpes zoster vaccinations, in agreement with current immunization guidelines prior to CIBINQO initiation [see Warnings and Precautions (5.7)].

2.2 Recommended Dosage

The recommended dosage of CIBINQO is 100 mg orally once daily.

If an adequate response is not achieved with CIBINQO 100 mg orally daily after 12 weeks, consider increasing dosage to 200 mg orally once daily. Discontinue therapy if inadequate response is seen after dosage increase to 200 mg once daily.

CIBINQO can be used with or without topical corticosteroids.

If a dose is missed, administer the dose as soon as possible unless it is less than 12 hours before the next dose, in which case skip the missed dose. Thereafter, resume dosing at the regular scheduled time.

2.3 Recommended Dosage in Patients with Renal Impairment or Hepatic Impairment

Renal Impairment

CIBINQO dosage recommendations for patients with renal impairment are provided in Table 1 [see *Use in Specific Populations (8.6) and Clinical Pharmacology (12.3)*]. In subjects with mild and moderate renal impairment, if an adequate response is not achieved after 12 weeks, dose of CIBINQO can be doubled [see *Dosage and Administration (2.2)*].

Table 1. Dosage Recommendations in Patients with Renal Impairment

Renal Impairment Stage	Estimated Glomerular Filtration (eGFR) ¹	Dosage
Mild	60 – 89 mL/minute	CIBINQO 100 mg once daily
Moderate	30 – 59 mL/minute	CIBINQO 50 mg once daily
Severe ²	15 – 29 mL/minute	Not recommended for use
End-Stage Renal Disease ² (ESRD)	<15 mL/minute	

¹ Glomerular filtration rate was estimated by the Modification of Diet in Renal Disease (MDRD) formula.

² Severe Renal Impairment and End-Stage Renal Disease include patients on renal replacement therapy.

Hepatic Impairment

CIBINQO is not recommended for use in patients with severe hepatic impairment [see *Use in Specific Populations (8.7) and Clinical Pharmacology (12.3)*].

2.4 Recommended Dosage in CYP2C19 Poor Metabolizers

In patients who are known or suspected to be CYP2C19 poor metabolizers, the recommended dosage of CIBINQO is 50 mg once daily [see *Use in Specific Populations (8.8) and Clinical Pharmacology (12.5)*]. If an adequate response is not achieved with CIBINQO 50 mg orally daily after 12 weeks, consider increasing dosage to 100 mg orally once daily. Discontinue therapy if inadequate response is seen after dosage increase to 100 mg once daily.

2.5 Dosage Modifications due to Strong Inhibitors

In patients taking strong inhibitors of cytochrome P450 (CYP) 2C19 reduce the dosage to 50 mg once daily [see *Drug Interactions (7.1) and Clinical Pharmacology (12.3)*]. If an adequate response is not achieved with CIBINQO 50 mg orally daily after 12 weeks, consider increasing dosage to 100 mg orally once daily. Discontinue therapy if inadequate response is seen after dosage increase to 100 mg once daily.

2.6 Treatment Discontinuation due to Serious Infections or Hematologic Adverse Reactions

Serious or Opportunistic Infections

If a patient develops a serious or opportunistic infection, discontinue CIBINQO and control the infection. The risks and benefits of treatment with CIBINQO should be carefully considered prior to reinitiating therapy with CIBINQO [see *Warnings and Precautions (5.1)*].

Hematologic Abnormalities

Recommendations for CIBINQO discontinuation for laboratory abnormalities are summarized in Table 2.

Table 2. Recommendations for CIBINQO Discontinuation for Hematologic Abnormalities

Laboratory Measure	Recommendation
Platelet Count <50,000/mm ³	Discontinue CIBINQO and follow with CBC until >100,000/mm ³
ALC <500/mm ³	Treatment should be temporarily discontinued if ALC is less than 500 cells/mm ³ and may be restarted once ALC return above this value
ANC <1,000/mm ³	Treatment should be temporarily discontinued if ANC is less than 1,000 cells/mm ³ and may be restarted once ANC return above this value
Hb value <8 g/dL	Treatment should be temporarily discontinued if Hb is less than 8 g/dL and may be restarted once Hb return above this value

Abbreviations: ALC=absolute lymphocyte count; ANC=absolute neutrophil count; CBC=complete blood count; Hb=hemoglobin

CBC evaluations are recommended at baseline, 4 weeks after treatment initiation and 4 weeks after dosing increase of CIBINQO. Laboratory evaluations may be extended for patients on chronic CIBINQO therapy who develop hematologic abnormalities [see *Warnings and Precautions (5.6)*].

2.7 Administration Instructions

Administer CIBINQO with or without food at approximately the same time each day.

Swallow CIBINQO tablets whole with water. Do not crush, split, or chew CIBINQO tablets.

3 DOSAGE FORMS AND STRENGTHS

- 50 mg: Pink, oval, film-coated tablet debossed with “PFE” on one side and “ABR 50” on the other.
- 100 mg: Pink, round, film-coated tablet debossed with “PFE” on one side and “ABR 100” on the other.
- 200 mg: Pink, oval, film-coated tablet debossed with “PFE” on one side and “ABR 200” on the other.

4 CONTRAINDICATIONS

CIBINQO is contraindicated in patients taking antiplatelet therapies, except for low-dose aspirin (≤ 81 mg daily), during the first 3 months of treatment [see *Warnings and Precautions (5.6)*, *Drug Interactions (7.2)*, and *Clinical Pharmacology (12.2)*].

5 WARNINGS AND PRECAUTIONS

5.1 Serious Infections

The most frequent serious infections reported in clinical studies with CIBINQO for atopic dermatitis were herpes simplex, herpes zoster, and pneumonia [see *Adverse Reactions (6.1)*]. Serious infections leading to hospitalization or death, including tuberculosis and bacterial, invasive fungal, viral, and other opportunistic infections, have occurred in patients receiving JAK inhibitors used to treat inflammatory conditions.

Avoid use of CIBINQO in patients with active, serious infection including localized infections.

Consider the risks and benefits of treatment prior to initiating CIBINQO in patients:

- with chronic or recurrent infection
- who have been exposed to tuberculosis

- with a history of a serious or an opportunistic infection
- who have resided or traveled in areas of endemic tuberculosis or endemic mycoses
- with underlying conditions that may predispose them to infection

Closely monitor patients for the development of signs and symptoms of infection during and after treatment with CIBINQO. If a patient develops a serious or opportunistic infection, discontinue CIBINQO. Initiate complete diagnostic testing and appropriate antimicrobial therapy. The risks and benefits of treatment with CIBINQO should be carefully considered prior to reinitiating therapy with CIBINQO.

Tuberculosis

Evaluate and test patients for TB before starting CIBINQO therapy and consider yearly screening for patients in highly endemic areas for TB. CIBINQO is not recommended for use in patients with active TB. For patients with a new diagnosis of latent TB or prior untreated latent TB, or for patients with a negative test for latent TB but who are at high risk for TB infection, start preventive therapy for latent TB prior to initiation of CIBINQO. Monitor patients for the development of signs and symptoms of TB, including patients who were tested negative for latent TB infection prior to initiating therapy.

Viral Reactivation

Viral reactivation, including herpes virus reactivation (e.g., herpes zoster, herpes simplex), was reported in clinical studies with CIBINQO [see *Adverse Reactions (6.1)*]. If a patient develops herpes zoster, consider interrupting CIBINQO until the episode resolves.

Hepatitis B virus (HBV) reactivation has been reported in patients receiving JAK inhibitors. Perform viral hepatitis screening and monitoring for reactivation in accordance with clinical guidelines before starting therapy and during therapy with CIBINQO. CIBINQO is not recommended for use in patients with active hepatitis B or hepatitis C [see *Clinical Pharmacology (12.3)*]. Monitor patients with inactive HBV for expression of HBV DNA during therapy with CIBINQO. If HBV DNA is detected during therapy with CIBINQO, consult a liver specialist.

5.2 Mortality

In a large, randomized, postmarketing safety study of another JAK inhibitor in rheumatoid arthritis (RA) patients 50 years of age and older with at least one cardiovascular risk factor, a higher rate of all-cause mortality, including sudden cardiovascular death, was observed in patients treated with the JAK inhibitor compared with TNF blockers. CIBINQO is not approved for use in RA.

Consider the benefits and risks for the individual patient prior to initiating or continuing therapy with CIBINQO.

5.3 Malignancy and Lymphoproliferative Disorders

Malignancies, including non-melanoma skin cancer (NMSC), were observed in clinical studies with CIBINQO for atopic dermatitis [see *Adverse Reactions (6.1)*].

Perform periodic skin examination for patients who are at increased risk for skin cancer. Exposure to sunlight and UV light should be limited by wearing protective clothing and using broad-spectrum sunscreen.

Malignancies, including lymphomas, have occurred in patients receiving JAK inhibitors used to treat inflammatory conditions. In a large, randomized, postmarketing safety study of another JAK inhibitor in RA patients, a higher rate of malignancies (excluding non-melanoma skin cancer (NMSC)) was observed in patients treated with the JAK inhibitor compared to those treated with TNF blockers. CIBINQO is not approved for use in RA. A higher rate of lymphomas was observed in patients treated with the JAK inhibitor compared to those treated with TNF blockers. A higher rate of lung cancers was observed in current or past smokers treated with the JAK inhibitor compared to those treated with TNF blockers. In this study, current or past smokers had an additional increased risk of overall malignancies.

Consider the benefits and risks for the individual patient prior to initiating or continuing therapy with CIBINQO, particularly in patients with a known malignancy (other than a successfully treated NMSC), patients who develop a malignancy when on treatment, and patients who are current or past smokers.

5.4 Major Adverse Cardiovascular Events

Major adverse cardiovascular events were reported in clinical studies of CIBINQO for atopic dermatitis [*see Adverse Reactions (6.1)*].

In a large, randomized, postmarketing safety study of another JAK inhibitor in RA patients 50 years of age and older with at least one cardiovascular risk factor, a higher rate of major adverse cardiovascular events (MACE) defined as cardiovascular death, non-fatal myocardial infarction (MI), and non-fatal stroke was observed with the JAK inhibitor compared to those treated with TNF blockers. CIBINQO is not approved for use in RA. Patients who are current or past smokers are at additional increased risk.

Consider the benefits and risks for the individual patient prior to initiating or continuing therapy with CIBINQO, particularly in patients who are current or past smokers and patients with other cardiovascular risk factors. Patients should be informed about the symptoms of serious cardiovascular events and the steps to take if they occur. Discontinue CIBINQO in patients that have experienced a myocardial infarction or stroke.

5.5 Thrombosis

Deep venous thrombosis (DVT) and pulmonary embolism (PE) were observed in patients receiving CIBINQO in the clinical studies for atopic dermatitis [*see Adverse Reactions (6.1)*].

Thrombosis, including DVT, PE, and arterial thrombosis have been reported in patients receiving JAK inhibitors used to treat inflammatory conditions. Many of these adverse reactions were serious and some resulted in death.

In a large, randomized, postmarketing safety study of another JAK inhibitor in RA patients 50 years of age and older with at least one cardiovascular risk factor, higher rates of overall thrombosis, DVT, and PE were observed compared to those treated with TNF blockers. CIBINQO is not approved for use in RA.

Avoid CIBINQO in patients that may be at increased risk of thrombosis. If symptoms of thrombosis occur, discontinue CIBINQO and evaluate and treat patients appropriately.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to CIBINQO during pregnancy. Pregnant women exposed to CIBINQO and health care providers are encouraged to call 1-877-311-3770.

Risk Summary

Available data from pregnancies reported in clinical trials with CIBINQO are not sufficient to establish a drug-associated risk for major birth defects, miscarriage, or other adverse maternal or fetal outcomes. In animal reproduction studies, oral administration of abrocitinib to pregnant rats and rabbits during organogenesis at exposure 11 or 4 times the maximum recommended human dose (MRHD) based on AUC comparison, respectively, resulted in maternal dystocia and skeletal variations in rats and no adverse effects in rabbits (*see Data*).

The background risks of major birth defects and miscarriage for the indicated population are unknown. All pregnancies carry some risk of birth defects, loss, or other adverse outcomes. The background risks in the U.S. general population of major birth defects and miscarriages are 2-4% and 15-20% of clinically recognized pregnancies, respectively.

Data

Animal Data

In an embryofetal development study, abrocitinib was administered orally to pregnant rats at doses of 10, 30, or 60 mg/kg/day during the period of organogenesis. No fetal malformations were observed. Abrocitinib increased the incidence of skeletal variations of short 13th ribs at 30 mg/kg/day (11 times the MRHD based on AUC comparison). Increased embryofetal lethality and additional skeletal variations (cervical arches with reduced ventral processes, thickened ribs, and unossified metatarsals) were noted at 60 mg/kg/day (17 times the MRHD based on AUC comparison).

In an embryofetal development study, abrocitinib was administered orally to pregnant rabbits at doses of 10, 30, or 75 mg/kg/day during the period of organogenesis. No abrocitinib-related maternal or developmental toxicity was noted at doses up to 75 mg/kg/day (4 times the MRHD based on AUC comparison).

In a prenatal and postnatal development study, abrocitinib was administered orally to pregnant rats at doses of 10, 30, and 60 mg/kg/day beginning on gestation day 6 and continuing through lactation day 20. Dystocia with prolonged parturition and reduced offspring body weights were noted at 30 mg/kg/day (11 times the MRHD based on AUC comparison). Postnatal survival was markedly decreased at 60 mg/kg/day (17 times the MRHD based on AUC comparison). No maternal toxicity was observed at 10 mg/kg/day (2.4 times the MRHD based on AUC comparison). No abrocitinib-related effects on postnatal developmental, neurobehavioral, or reproductive performance of offspring was noted at doses up to 30 mg/kg/day (11 times the MRHD based on AUC comparison).

8.2 Lactation

Risk Summary

There are no data on the presence of abrocitinib in human milk, the effects on the breast-fed infant, or the effects on milk production. Abrocitinib was secreted in milk of lactating rats (*see Data*). When a drug is present in animal milk, it is likely that the drug will be present in human milk. Because of the serious adverse findings in adults, including risks of serious infections, malignancy, and thrombosis, advise women not to breastfeed during treatment with CIBINQO and for one day after the last dose (approximately 5-6 elimination half-lives).

Data

Animal Data

Lactating female rats were orally administered a single dose of 10 mg/kg abrocitinib on lactation day 12. Abrocitinib AUC was approximately 5 times greater in milk than in plasma.

8.3 Females and Males of Reproductive Potential

Infertility

Females

Based on the findings in rats, oral administration of CIBINQO may impair female fertility. Impaired fertility in female rats was reversible 1 month after cessation of abrocitinib oral administration [*see Nonclinical Toxicology (13.1)*].

8.4 Pediatric Use

The safety and effectiveness of CIBINQO in pediatric patients 12 years of age and older weighing 25 kg or more with atopic dermatitis has been established. In trials Trial-AD-1 and Trial-AD-2, 124 adolescent subjects 12 to less than 18 years old with moderate-to-severe atopic dermatitis were enrolled and randomized to receive either CIBINQO 100 mg (N=51), 200 mg (N=48), or matching placebo (N=25) in monotherapy. Additional 284 adolescent subjects 12 to less than 18 years of age with moderate-to-severe atopic dermatitis, were enrolled and randomized to receive either CIBINQO 100 mg (N=95) or 200 mg (N=94) or matching placebo (N=95) in combination with topical corticosteroids in Trial-AD-4. Efficacy and adverse reaction profile were consistent between the pediatric patients and adults [*see Clinical Studies (14) and Adverse Reactions (6.1)*].

The safety and effectiveness of CIBINQO have not been established in pediatric patients below 12 years of age.

Juvenile Animal Toxicity Data

In a juvenile animal toxicity study, abrocitinib was administered orally to juvenile rats at doses of 5, 25, and 75 mg/kg/day beginning on postnatal day 10 (approximately equivalent to a human infant) and continuing through postnatal day 63 (approximately equivalent to an adolescent). Abrocitinib caused a reversible, dose-related decrease in the primary spongiosa in the metaphysis of the proximal tibia and distal femur. Abrocitinib produced adverse effects on bone development at all dose levels. Abrocitinib caused irreversible dose-related small or misshapen femoral heads at doses ≥ 5 mg/kg/day (0.8 times the MRHD based on AUC comparison). Abrocitinib also irreversibly decreased femur size and caused paw malrotation and limb impairment at doses ≥ 25 mg/kg/day (7.2 times the MRHD based on AUC comparison). At 75 mg/kg/day (27 times the MRHD based on AUC comparison), paw fractures generally corresponded to limb impairment, a fractured tibia was noted in a single female. Irreversible bone findings have not been observed in older animals.

8.5 Geriatric Use

A total of 145 (4.6%) patients 65 years of age and older, while 25 (0.8%) were 75 years of age and older, were enrolled in CIBINQO clinical trials. Clinical trials of CIBINQO did not include sufficient numbers of patients 65 years of age and older to determine whether they respond differently from younger adult patients.

A higher proportion of patients 65 years of age and older discontinued from clinical trials compared to younger patients. Among all patients exposed to CIBINQO, including the long-term extension trial, confirmed ALC $<500/\text{mm}^3$ occurred only in patients 65 years of age and older. A higher proportion of patients 65 years of age and older had platelet counts $<75,000/\text{mm}^3$. The incidence rate of herpes zoster in patients 65 years of age and older treated with CIBINQO (7.40 per 100 patient-years) was higher than that of patients 18 to less than 65 years of age (3.44 per 100 patient-years).

8.6 Renal Impairment

In patients with severe (eGFR <30 mL/min) and moderate (eGFR 30-59 mL/min) renal impairment, the combined exposure ($\text{AUC}_{\text{inf,u}}$) of abrocitinib and its two active metabolites, M1 and M2, is increased compared to patients with normal renal function (eGFR ≥ 90 mL/min) [see *Clinical Pharmacology (12.3)*]. This may increase the risk of adverse reactions such as infections.

CIBINQO is not recommended for use in patients with severe renal impairment and ESRD including those on renal replacement therapy [see *Dosage and Administration (2.3)*].

A dosage reduction in patients with moderate renal impairment is recommended. No dosage adjustment is required in patients with mild renal impairment (eGFR 60-89 mL/min) [see *Dosage and Administration (2.3)*].

CIBINQO has not been studied in patients on renal replacement therapy. In Phase 3 clinical trials, CIBINQO was not evaluated in patients with atopic dermatitis with baseline creatinine clearance values less than 40 mL/min.

8.7 Hepatic Impairment

Avoid use of CIBINQO in patients with severe (Child Pugh C) hepatic impairment.

Dosage adjustment is not required in patients with mild (Child Pugh A) or moderate (Child Pugh B) hepatic impairment based on similar combined exposure ($\text{AUC}_{\text{inf,u}}$) of abrocitinib and its two active metabolites, M1 and M2 compared to patients with normal hepatic function. In clinical trials, CIBINQO was not evaluated in patients with severe (Child Pugh C) hepatic impairment [see *Clinical Pharmacology (12.3)*].

8.8 CYP2C19 Poor Metabolizers

In patients who are CYP2C19 poor metabolizers, the AUC of abrocitinib is increased compared to CYP2C19 normal metabolizers due to reduced metabolic clearance. Dosage reduction of CIBINQO is recommended in patients who are known or suspected to be CYP2C19 poor metabolizers based on genotype or previous history/experience with other CYP2C19 substrates [see *Dosage and Administration (2.4)* and *Clinical Pharmacology (12.5)*].

10 OVERDOSAGE

There is no experience regarding human overdose with CIBINQO. There is no specific antidote for overdose with CIBINQO. In case of an overdose, call Poison Control Center at 1-800-222-1222 for latest recommendations.

11 DESCRIPTION


CIBINQO (abrocitinib) tablets contain the free base of abrocitinib, a Janus kinase (JAK) inhibitor, for oral administration.

Abrocitinib is a white to pale colored powder with the following chemical name:

N-((1*s*,3*s*)-3-(methyl(7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl)amino)cyclobutyl)propane-1-sulfonamide

The solubility of abrocitinib in water is 0.04 mg/mL at 25°C.

Abrocitinib has a molecular weight of 323.42 g/mol and a molecular formula of C₁₄H₂₁N₅O₂S. The structural formula of abrocitinib is:



Each film-coated tablet contains 50 mg or 100 mg or 200 mg of abrocitinib and the following inactive ingredients: dibasic calcium phosphate anhydrous, hypromellose, iron oxide red, lactose monohydrate, Macrogol, magnesium stearate, microcrystalline cellulose, sodium starch glycolate, titanium dioxide, and triacetin.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

CIBINQO is a Janus kinase (JAK) inhibitor.

Abrocitinib reversibly inhibits JAK1 by blocking the adenosine triphosphate (ATP) binding site. In a cell-free isolated enzyme assay, abrocitinib was selective for JAK1 over JAK2 (28-fold), JAK3 (>340-fold), and tyrosine kinase (TYK) 2 (43-fold), as well as the broader kinome. The relevance of inhibition of specific JAK enzymes to therapeutic effectiveness is not currently known. Both the parent compound and the active metabolites inhibit JAK1 activity *in vitro* with similar levels of selectivity.

12.2 Pharmacodynamics

Treatment with CIBINQO was associated with dose-dependent reduction in serum markers of inflammation, including high sensitivity C-reactive protein (hsCRP), interleukin-31 (IL-31) and thymus and activation regulated chemokine (TARC). These changes returned to near baseline within 4 weeks of drug discontinuation.

Effect on Platelet Count

Treatment with CIBINQO was also associated with a transient, dose-dependent decrease in platelet count with the nadir occurring at a median of 24 days after continuous administration of abrocitinib 200 mg once daily. The percent change from baseline of the nadir increases with decreasing baseline platelet counts (-41.2%, -33.4%, and -26.5% for baseline platelet counts of 170 , 220 , and $270 \times 10^3/\text{mm}^3$, respectively). Recovery of platelet count (~40% recovery by 12 weeks) occurred without discontinuation of the treatment.

Cardiac Electrophysiology

At a dose 3 times the maximum approved recommended dose, abrocitinib does not prolong the QT interval to any clinically relevant extent.

12.3 Pharmacokinetics

Abrocitinib plasma C_{max} and AUC increased dose proportionally up to 200 mg. Steady-state plasma concentrations of abrocitinib are achieved within 48 hours after once daily administration.

Absorption

Abrocitinib is absorbed with over 91% extent of oral absorption and absolute oral bioavailability of approximately 60%. The peak plasma concentrations of abrocitinib are reached within 1 hour.

Effect of Food

Coadministration of CIBINQO with a high-fat, high-calorie meal (total 916 calories, with approximate distribution of 55% fat, 29% carbohydrates, and 16% protein) had no clinically relevant effect on abrocitinib exposures (AUC and C_{max} of abrocitinib increased by approximately 26% and 29%, respectively, and T_{max} was prolonged by 2 hours) [see *Dosage and Administration (2.7)*].

Distribution

After intravenous administration, the volume of distribution of abrocitinib is approximately 100 L. Approximately 64%, 37% and 29% of circulating abrocitinib and its active metabolites M1 and M2, respectively, are bound to plasma proteins. Abrocitinib and its active metabolites M1 and M2 bind predominantly to albumin and distribute equally between red blood cells and plasma.

Elimination

Abrocitinib is eliminated primarily by metabolic clearance mechanisms. The mean elimination half-lives of abrocitinib and its two active metabolites, M1 and M2, range 3 to 5 hours.

Metabolism

The metabolism of abrocitinib is mediated by multiple CYP enzymes, CYP2C19 (~53%), CYP2C9 (~30%), CYP3A4 (~11%) and CYP2B6 (~6%). In a human radiolabeled study, abrocitinib was the most prevalent circulating species, with two active polar mono-hydroxylated metabolites identified as M1 (3-hydroxypropyl), and M2 (2-hydroxypropyl). Metabolite M1 is less active than abrocitinib while metabolite M2 is as active as the parent. The pharmacologic activity of abrocitinib is attributable to the unbound exposure of parent molecule (~60%) as well as M1 (~10%) and M2 (~30%) in systemic circulation. The sum of unbound exposures of

abrocitinib, M1 and M2, each expressed in molar units and adjusted for relative potencies, is referred to as the combined exposure of abrocitinib and its two active metabolites, M1 and M2.

Excretion

After a single radiolabeled abrocitinib dose, less than 1% of the dose was excreted in urine as unchanged drug. The metabolites of abrocitinib, M1 and M2 are excreted predominantly in urine, and are substrates of OAT3 transporter.

Specific Populations

Body weight, sex, race, and age did not have a clinically meaningful effect on CIBINQO exposure.

Patients with Renal Impairment

In a renal impairment study, subjects with severe (eGFR <30 mL/min as estimated by MDRD equation) and moderate (eGFR 30-59 mL/min, MDRD) renal impairment had approximately 191% and 110% increase in the combined exposure ($AUC_{inf,u}$) of abrocitinib and its active metabolites, M1 and M2, respectively, compared to subjects with normal renal function (eGFR \geq 90 mL/min, MDRD). Based on these results, a clinically significant increase in the combined exposure of abrocitinib and its active metabolites, M1 and M2, is not expected in subjects with mild renal impairment (eGFR 60 -89 mL/min, MDRD) [see *Dosage and Administration (2.3)* and *Use in Specific Population (8.6)*].

CIBINQO has not been studied in subjects on renal replacement therapy [see *Dosage and Administration (2.3)* and *Use in Specific Population (8.6)*]. In Phase 3 clinical studies, CIBINQO was not evaluated in subjects with atopic dermatitis with baseline creatinine clearance values less than 40 mL/min.

Patients with Hepatic Impairment

Subjects with mild hepatic impairment (Child Pugh A) had approximately 4% decrease in the combined exposure ($AUC_{inf,u}$) of abrocitinib and its two active metabolites, M1 and M2, compared to subjects with normal hepatic function. Subjects with moderate hepatic impairment (Child Pugh B) had approximately 15% increase in the combined exposure ($AUC_{inf,u}$) of abrocitinib and its two active metabolites, M1 and M2, compared to subjects with normal hepatic function. These changes are not clinically significant. In clinical studies, CIBINQO has not been studied in subjects with severe (Child Pugh C) hepatic impairment, or in subjects screened positive for active hepatitis B or hepatitis C [see *Use in Specific Populations (8.7)* and *Warnings and Precautions (5.1)*].

Drug Interaction Studies

Clinical Studies

The effect of coadministered drugs on the pharmacokinetics of abrocitinib is presented in Table 6.

Table 6. Change in Pharmacokinetics of the Combined Exposure of Abrocitinib and its Two Active Metabolites (M1 and M2) in the Presence of Coadministered Drugs

Coadministered Drugs	Regimen of Coadministered Drug	Dose of Abrocitinib	Ratio ^a (90% Confidence Interval)	
			C _{max,u}	AUC _{inf,u}
Strong CYP2C19 and moderate CYP3A inhibitor: Fluvoxamine [see Drug Interactions (7.1)]	50 mg once daily x 9 days	100 mg	1.33 (1.00-1.78)	1.91 (1.74-2.10)
Strong CYP2C19, moderate CYP2C9 and CYP3A inhibitor: Fluconazole [see Drug Interactions (7.1)]	400 mg on Day 1 and 200 mg on Days 2-7	100 mg	1.23 (1.08-1.42)	2.55 ^b (2.42-2.69)
Strong CYP Enzymes Inducers: Rifampin [see Drug Interactions (7.1)]	600 mg once daily x 8 days	200 mg	0.69 (0.50-0.94)	0.44 (0.41-0.47)
OAT3 inhibitor: Probenecid ^c	1,000 mg twice daily x 3 days	200 mg	1.30 (1.04-1.63)	1.66 (1.52-1.80)

^a Ratios for C_{max,u} and AUC_{inf,u} compare coadministration of the drug with abrocitinib versus administration of abrocitinib alone.

^b When coadministered with Fluconazole, the systemic exposure of abrocitinib was approximately 4.8-fold higher compared to when abrocitinib is administered alone.

^c Drug interaction with OAT3 inhibitor is not clinically significant.

The effect of abrocitinib on the pharmacokinetics of coadministered drugs is presented in Table 7.

Table 7. Change in Pharmacokinetics of Coadministered Drugs in the Presence of Abrocitinib

Coadministered Drugs or In Vivo Markers of CYP Activity	Dose Regimen of Abrocitinib	Ratio ^a (90% Confidence Interval)	
		C _{max}	AUC _{inf}
Oral contraceptive: Ethinyl estradiol (EE) and levonorgestrel (LN)	200 mg once daily x 9 days	EE: 1.07 (0.99, 1.15) LN: 0.86 (0.75, 0.97)	EE: 1.19 (1.12, 1.26) LN ^b : 0.98 (0.87, 1.10)
Sensitive CYP3A Substrate: Midazolam	200 mg once daily x 7 days	0.93 (0.84, 1.04)	0.92 (0.86, 0.99)
Sensitive P-gp substrate: Dabigatran	200 mg single dose	1.40 (0.92, 2.13)	1.53 (1.09, 2.15)
Sensitive BCRP and OAT3 substrate: Rosuvastatin	200 mg once daily x 3 days	0.99 (0.86, 1.14)	1.02 (0.93, 1.12)
Sensitive MATE1/2K substrate: Metformin	200 mg once daily x 2 days	0.88 (0.81, 0.96)	0.93 (0.85, 1.03)

^a Ratios for C_{max} and AUC_{inf} compare coadministration of abrocitinib with the drug versus administration of the drug alone.

^b AUC_{last} of levonorgestrel was reported in lieu of AUC_{inf} because the terminal phase of levonorgestrel was not well characterized.

Coadministration of dabigatran etexilate (a P-gp substrate), with a single dose of CIBINQO 200 mg increased dabigatran AUC_{inf} and C_{max} by approximately 53% and 40%, respectively, compared with administration alone. These increases in dabigatran exposure are not considered clinically significant change. However, appropriate dose titration of P-gp substrate where small concentration changes may lead to serious or life-threatening toxicities (e.g., digoxin) when coadministered with the CIBINQO would be needed.

In Vitro Studies

Cytochrome P450 (CYP) Enzymes: Abrocitinib and its metabolites M1 and M2 are not inhibitors or inducers of CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, or CYP3A4.

Uridine diphosphate (UDP)-glucuronosyl transferase (UGT) Enzymes: Abrocitinib and its metabolites M1 and M2 are not inhibitors or inducers of UGT1A1, UGT1A4, UGT1A6, UGT1A9, or UGT2B7.

Transporter Systems: Abrocitinib is an inhibitor of organic cation transporter (OCT)1 but is not an inhibitor of organic anion transporting polypeptide (OATP)1B1/1B3, bile salt export pump (BSEP), OAT1 or OCT2.

12.5 Pharmacogenomics

Patients who are CYP2C19 poor metabolizers have little to no CYP2C19 enzyme function compared to CYP2C19 normal metabolizers that have fully functional CYP2C19 enzymes.

After single doses of abrocitinib, CYP2C19 poor metabolizers demonstrated dose-normalized AUC of abrocitinib values that were 2.3-fold higher when compared to CYP2C19 normal metabolizers. Approximately 3-5% of Caucasians and Blacks and 15 to 20% of Asians are CYP2C19 poor metabolizers [see *Dosage and Administration (2.4) and Use in Specific Populations (8.8)*].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 2-year oral carcinogenicity study in rats, abrocitinib increased the incidence of benign thymomas in female rats at doses of 10 and 30 mg/kg/day (2.8 and 14 times the MRHD, respectively, based on AUC comparison). Abrocitinib was not carcinogenic in female rats at 3 mg/kg/day (0.6 times the MRHD based on AUC comparison) or male rats at doses up to 30 mg/kg/day (14 times the MRHD based on AUC comparison). Abrocitinib was not carcinogenic in Tg.rasH2 mice at oral doses up to 60 mg/kg/day in males and 75 mg/kg/day in females.

Abrocitinib was not mutagenic in the bacterial mutagenicity assay (Ames assay). Although abrocitinib was aneugenic in the in vitro TK6 micronucleus assay, abrocitinib was not aneugenic or clastogenic in an in vivo rat bone marrow micronucleus assay.

Abrocitinib did not impair male fertility at doses up to 70 mg/kg/day (26 times the MRHD based on AUC comparison) or female fertility at 10 mg/kg/day (2 times the MRHD based on AUC comparison). Abrocitinib impaired female fertility (reducing fertility index, corpora lutea, and implantation sites) at 70 mg/kg/day (29 times the MRHD based on AUC comparison). Impaired fertility in female rats reversed 1 month after cessation of abrocitinib administration.

14 CLINICAL STUDIES

The efficacy of CIBINQO as monotherapy and in combination with background topical corticosteroids was evaluated in 3 randomized, double-blind, placebo-controlled trials [Trial-AD-1 (NCT03349060), Trial-AD-2 (NCT03575871), and Trial-AD-3 (NCT03720470)] in 1615 subjects 12 years of age and older with moderate-to-severe atopic dermatitis as defined by Investigator's Global Assessment (IGA) score ≥ 3 , Eczema Area and Severity Index (EASI) score ≥ 16 , body surface area (BSA) involvement $\geq 10\%$, and Peak Pruritus Numerical Rating Scale (PP-NRS) ≥ 4 at the baseline visit prior to randomization.

The efficacy and safety of CIBINQO in combination with background topical corticosteroids were further evaluated in adolescent subjects in a randomized, double-blind, placebo-controlled trial [Trial-AD-4 (NCT03796676)]. The trial included 284 subjects who were 12 to less than 18 years of age with moderate-to-

severe atopic dermatitis as defined by IGA score ≥ 3 , EASI score ≥ 16 , BSA involvement $\geq 10\%$, and PP-NRS ≥ 4 at the baseline visit prior to randomization.

Baseline characteristics

In trials Trial-AD-1, Trial-AD-2, Trial-AD-3, 53% of subjects were male, 69% of subjects were white, 64% of subjects had a baseline IGA score of 3 (moderate AD), and 36% of subjects had a baseline IGA score of 4 (severe AD). The baseline mean EASI score was 30. The baseline mean age was 36 years old with 8% of subjects 12 to less than 18 years old and 92% of subjects 18 years of age or older. Subjects in these trials were those who had inadequate response to previous topical therapy, or were subjects for whom topical treatments were medically inadvisable, or who had received systemic therapies including dupilumab. In each of the trials, over 40% of subjects had prior exposure to systemic therapy. In Trial-AD-1 and Trial-AD-2, 6% of the subjects had received dupilumab, whereas prior use of dupilumab was not allowed in Trial-AD-3.

In Trial-AD-4, 49% of subjects were female, 56% of subjects were White, 33% of subjects were Asian and 6% of subjects were Black. The median age was 15 years and the proportion of subjects with severe atopic dermatitis (IGA of 4) was 38%.

Trial-AD-1, Trial-AD-2, Trial-AD-3, and Trial-AD-4 assessed the co-primary endpoints of IGA and EASI-75 responses at Week 12. The designs of the trials are summarized in Table 8.

Table 8. Summary of Clinical Study Designs

Study Name (regimen type)	Population (number of randomized and dosed subjects)	Treatment Arms	Co-Primary Endpoints
Trial-AD-1 (monotherapy) 12 weeks	Subjects 12 years of age or older (387)	Oral administration of: <ul style="list-style-type: none"> • CIBINQO 200 mg QD • CIBINQO 100 mg QD • Placebo 	<ul style="list-style-type: none"> • IGA response^a at Week 12 • EASI-75^b at Week 12
Trial-AD-2 (monotherapy) 12 weeks	Subjects 12 years of age or older (391)	Oral administration of: <ul style="list-style-type: none"> • CIBINQO 200 mg QD • CIBINQO 100 mg QD • Placebo 	
Trial-AD-3 (combination therapy) 16 weeks	Subjects 18 years of age or older (837)	Oral administration of: <ul style="list-style-type: none"> • CIBINQO 200 mg QD • CIBINQO 100 mg QD • Placebo Subcutaneous administration of: <ul style="list-style-type: none"> • Dupilumab 300 mg Q2W SC^c All subjects received background topical corticosteroids	
Trial-AD-4 (combination therapy) 12 weeks	Subjects 12 to less than 18 years of age (285)	Oral administration of: <ul style="list-style-type: none"> • CIBINQO 200 mg QD • CIBINQO 100 mg QD • Placebo All subjects received background topical corticosteroids	

Abbreviations: EASI=Eczema Area and Severity Index; IGA=Investigator's Global Assessment; QD=once daily; Q2W=once every 2 weeks.

a. IGA response was based on IGA score of clear (0) or almost clear (1) (on a 5-point scale) and a reduction from baseline of ≥ 2

points.

- b. EASI-75 was based on $\geq 75\%$ improvement in EASI from baseline.
- c. Dupilumab treatment in Trial-AD-3: An initial dose of 600 mg on day 1, followed by 300 mg Q2W.

Clinical Response

Monotherapy Trials

The results of the CIBINQO monotherapy trials (Trial-AD-1 and Trial-AD-2) are presented in Table 9.

Table 9. Efficacy Results of CIBINQO Monotherapy at Week 12 in Subjects with Moderate-to-Severe AD (Trial-AD-1 and Trial-AD-2)

	Trial-AD-1			Trial-AD-2		
	CIBINQO		Placebo N=77	CIBINQO		Placebo N=78
	200 mg QD N=154	100 mg QD N=156		200 mg QD N=155	100 mg QD N=158	
IGA 0 or 1 ^a	44%	24%	8%	38%	28%	9%
Difference from Placebo (95% CI)	36% (26%, 46%)	16% (7%, 25%)	-	29% (19%, 39%)	19% (9%, 29%)	-
EASI-75 ^b	62%	40%	12%	61%	44%	10%
Difference from Placebo (95% CI)	51% (40%, 61%)	28% (18%, 39%)	-	50% (40%, 61%)	33% (23%, 44%)	-

Abbreviations: CI=confidence interval; EASI=Eczema Area and Severity Index; IGA=Investigator Global Assessment; QD=once daily.

- a. IGA responders were subjects with IGA score of clear (0) or almost clear (1) (on a 5-point scale) and a reduction from baseline of ≥ 2 points.
- b. EASI -75 responders were patients with $\geq 75\%$ improvement in EASI from baseline.

The proportion of subjects achieving PP-NRS4 at Week 2 (defined as an improvement of ≥ 4 points from baseline in PP-NRS) was higher in subjects treated with CIBINQO monotherapy 200 mg once daily (28% in Trial-AD-1 and 24% in Trial-AD-2) and 100 mg once daily (11% in both trials) compared to placebo (2% in both trials).

A higher proportion of subjects in the CIBINQO monotherapy 100 mg or 200 mg once daily arm compared to placebo achieved improvement in itching at Week 12.

Combination Therapy Trial

The results of CIBINQO in combination with background topical corticosteroids (Trial-AD-3) are presented in Table 10.

Table 10. Efficacy Results of CIBINQO with Concomitant Topical Corticosteroids at Week 12 in Subjects with Moderate-to-Severe AD (Trial-AD-3)

% Responders	CIBINQO		Placebo N=131
	200 mg QD N=226	100 mg QD N=238	
IGA 0 or 1 ^a at Week 12	47%	36%	14%
Difference from Placebo (95% CI)	34% (25%, 42%)	23% (15%, 31%)	-
EASI-75 ^b at Week 12	68%	58%	27%
Difference from Placebo (95% CI)	41% (32%, 51%)	32% (22%, 41%)	-

Abbreviations: CI=confidence interval; EASI=Eczema Area and Severity Index; IGA=Investigator Global Assessment; QD=once daily.

- a. IGA responders were subjects with IGA score of clear (0) or almost clear (1) (on a 5-point scale) and a reduction from baseline of ≥ 2 points.
- b. EASI-75 responders were subjects with $\geq 75\%$ improvement in EASI, from baseline.

The proportions of subjects achieving PP-NRS4 at Week 2 was higher in subjects treated with CIBINQO 200 mg once daily (30%) and 100 mg once daily (14%) in combination with background medicated topical therapies compared to placebo (8%).

Examination of age, gender, race, weight and previous systemic AD therapy treatment did not identify differences in response to CIBINQO 100 mg or 200 mg once daily among these subgroups in Trial-AD-1, Trial-AD-2, and Trial-AD-3.

The results of CIBINQO in combination with background topical corticosteroids for adolescent subjects (Trial-AD-4) are presented in Table 11.

Table 11. Efficacy Results of CIBINQO with Concomitant Topical Corticosteroids at Week 12 in Adolescent Subjects with Moderate-to-Severe AD (Trial-AD-4)

% Responders	CIBINQO		Placebo N=95
	200 mg QD N=94	100 mg QD N=95	
IGA 0 or 1 ^a	46%	39%	24%
Difference from Placebo (95% CI)	21% (8%, 34%)	15% (2%, 28%)	-
EASI-75 ^b	71%	64%	41%
Difference from Placebo (95% CI)	29% (16%, 43%)	23% (10%, 36%)	-

Abbreviations: CI=confidence interval; EASI=Eczema Area and Severity Index; IGA=Investigator Global Assessment; N=number of patients treated; QD=once daily.

- a. IGA responders were patients with IGA score of clear (0) or almost clear (1) (on a 5-point scale) and a reduction from baseline of ≥ 2 points.
- b. EASI-75 responders were patients with $\geq 75\%$ improvement in EASI from baseline.

The proportion of adolescents achieving PP-NRS4 at Week 2 in Trial-AD-4 was higher with CIBINQO 200 mg once daily (25%) and 100 mg once daily (13%) compared to placebo (8%).

A higher proportion of subjects in the CIBINQO 200 mg once daily arm compared to placebo achieved improvement in itching at Week 12.

16 HOW SUPPLIED/STORAGE AND HANDLING

CIBINQO is supplied as:

Dosage Form	Strength	Description	Bottle Size (number of tablets)	NDC Number
Tablets	50 mg	Pink, oval tablet debossed with “PFE” on one side and “ABR 50” on the other.	30 count bottle	0069-0235-30
Tablets	100 mg	Pink, round tablet debossed with “PFE” on one side and “ABR 100” on the other.	30 count bottle	0069-0335-30
Tablets	200 mg	Pink, oval tablet debossed with “PFE” on one side and “ABR 200” on the other.	30 count bottle	0069-0435-30

Store CIBINQO at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C to 30°C (59°F to 86°F). Keep in original package. The container closure system is child resistant.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Pregnancy Registry

Advise patients to report their pregnancy to 1-877-311-3770 [see *Use in Specific Populations (8.1)*].

Serious Infections

Inform patients that they may develop infections when taking CIBINQO. Instruct patients to tell their healthcare provider if they develop any signs or symptoms of an infection [see *Warnings and Precautions (5.1)*].

Advise patients that the risk of herpes zoster is increased in patients treated with CIBINQO and some cases can be serious [see *Warnings and Precautions (5.1)*].

Malignancies

Inform patients that CIBINQO may increase their risk of certain cancers, including skin cancers. Periodic skin examinations are recommended while using CIBINQO. Advise patients that exposure to sunlight and UV light should be limited by wearing protective clothing and using a broad-spectrum sunscreen [see *Warnings and Precautions (5.3)*].

Major Adverse Cardiovascular Events

Inform patients that CIBINQO may increase their risk of major adverse cardiovascular events (MACE) including myocardial infarction, stroke, and cardiovascular death. Instruct all patients, especially current or past smokers or patients with other cardiovascular risk factors, to be alert for the development of signs and symptoms of cardiovascular events [see *Warnings and Precautions (5.4)*].

Thrombosis

Advise patients that events of DVT and PE have been reported in clinical trials with CIBINQO. Instruct patients to seek immediate medical attention if they develop any signs or symptoms of a DVT or PE [see *Warnings and Precautions (5.5)*].

Laboratory Abnormalities

Inform patients that CIBINQO may affect certain lab tests, and that blood tests are required before and during CIBINQO treatment [see *Dosage and Administration (2.1)* and *Warnings and Precautions (5.6)*].

Immunizations

Advise patients that vaccination with live vaccines is not recommended during CIBINQO treatment and immediately prior to or after CIBINQO treatment. Instruct patients to inform the healthcare practitioner that they are taking CIBINQO prior to a potential vaccination [see *Warnings and Precautions (5.7)*].

Retinal Detachment

Inform patients that retinal detachment has been reported in clinical trials for atopic dermatitis in patients who received CIBINQO. Advise patients to immediately inform their healthcare provider if they develop any sudden changes in vision while receiving CIBINQO [see *Adverse Reactions (6.1)*].

Infertility

Advise females of reproductive potential that CIBINQO may impair fertility [see *Use in Specific Populations (8.3)*].

Lactation

Advise a woman not to breastfeed during treatment with CIBINQO [see *Use in Specific Populations (8.2)*].

Administration

Advise patients not to chew, crush, or split CIBINQO tablets [see *Dosage and Administration (2.7)*].

This product's labeling may have been updated. For the most recent prescribing information, please visit www.pfizer.com.

For Medical Information about CIBINQO, please visit www.pfizermedinfo.com or call 1-800-438-1985.



Distributed by

Pfizer Labs

Division of Pfizer Inc.
New York, NY 10017

LAB-1423-1.5

Medication Guide
CIBINQO (Si BINK oh)
(abrocitinib)
tablets, for oral use

What is the most important information I should know about CIBINQO?

CIBINQO may cause serious side effects, including:

1. Serious infections

CIBINQO is a medicine that affects your immune system. CIBINQO can lower the ability of your immune system to fight infections. Some people have had serious infections while taking CIBINQO or other similar medicines, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses that can spread throughout the body. Some people have been hospitalized or died from these infections.

- Your healthcare provider should test you for TB before starting treatment with CIBINQO.
- Your healthcare provider should watch you closely for signs and symptoms of TB during treatment with CIBINQO.

You should not start taking CIBINQO if you have any kind of infection unless your healthcare provider tells you it is okay.

You may be at a higher risk of developing shingles (herpes zoster).

Before starting CIBINQO, tell your healthcare provider if you:

- are being treated for an infection
- have had an infection that does not go away or that keeps coming back
- have diabetes, chronic lung disease, HIV, or a weak immune system
- have TB or have been in close contact with someone with TB
- have had shingles (herpes zoster)
- have had hepatitis B or hepatitis C
- live or have lived or have traveled to certain parts of the country (such as the Ohio and Mississippi River valleys and the Southwest) where there is an increased chance for getting certain kinds of fungal infections. These infections may happen or become more severe if you use CIBINQO. Ask your healthcare provider if you do not know if you have lived in an area where these infections are common.
- think you have an infection or have symptoms of an infection such as:
 - fever, sweating, or chills
 - muscle aches
 - cough or shortness of breath
 - blood in your phlegm
 - weight loss
 - warm, red, or painful skin or sores on your body
 - diarrhea or stomach pain
 - burning when you urinate or urinating more often than usual
 - feeling very tired

After starting CIBINQO, call your healthcare provider right away if you have any

symptoms of an infection. CIBINQO can make you more likely to get infections or make any infections that you have worse. If you get a serious infection, your healthcare provider may stop treatment with CIBINQO until your infection is controlled.

2. Increased risk of death in people 50 years of age and older who have at least 1 heart disease (cardiovascular) risk factor and are taking a medicine in the class of medicines called Janus kinase (JAK) inhibitors. CIBINQO is a JAK inhibitor medicine.

3. Cancer and immune system problems

CIBINQO may increase your risk of certain cancers by changing the way your immune system works.

- Lymphoma and other cancers, including skin cancers, can happen in people taking CIBINQO.
- People taking a medicine in the class of medicines called Janus kinase (JAK) inhibitors have a higher risk of certain cancers including lymphoma and lung cancer, especially if you are a current or past smoker.
- Follow your healthcare provider's advice about having your skin checked for skin cancer during treatment with CIBINQO. Limit the amount of time you spend in sunlight. Avoid using tanning beds or sunlamps. Wear protective clothing when you are in the sun and use a sunscreen with a high protection factor (SPF 30 and above). This is especially important if your skin is very fair or if you have a family history of skin cancer.

Tell your healthcare provider if you have ever had any type of cancer.

4. Increased risk of major cardiovascular events such as heart attack, stroke or death in people 50 years of age and older who have at least 1 heart disease (cardiovascular) risk factor and taking a medicine in the class of medicines called JAK inhibitors, especially if you are a current or past smoker.

Some people taking CIBINQO have had major cardiovascular events.

Get emergency help right away if you develop any symptoms of a heart attack or stroke during treatment with CIBINQO, including:

- discomfort in the center of your chest that lasts for more than a few minutes, or that goes away and comes back
- severe tightness, pain, pressure, or heaviness in your chest, throat, neck, or jaw
- pain or discomfort in your arms, back, neck, jaw, or stomach
- weakness in one part or on one side of your body
- slurred speech
- shortness of breath with or without chest discomfort
- breaking out in a cold sweat
- nausea or vomiting
- feeling lightheaded

5. Blood clots

Blood clots in the veins of your legs (deep vein thrombosis, DVT) or lungs (pulmonary embolism, PE) can happen in some people taking CIBINQO. This may be life-threatening. Blood clots in the veins of the legs (deep vein thrombosis, DVT) and lungs (pulmonary embolism, PE) have happened more often in people who are 50 years of age and older and with at least 1 heart disease (cardiovascular) risk factor taking a medicine in the class of medicines called Janus kinase (JAK) inhibitors.

- Tell your healthcare provider if you have had blood clots in the veins of your legs or lungs in the past.
- Stop taking CIBINQO and get medical help right away if you have any signs and symptoms of blood clots during treatment with CIBINQO, including:
 - swelling, pain or tenderness in one or both legs
 - sudden, unexplained chest or upper back pain
 - shortness of breath or difficulty breathing

6. Changes in certain laboratory test results

Your healthcare provider should do blood tests before you start taking CIBINQO and during treatment with CIBINQO to check for the following:

- **low lymphocyte count.** Lymphocytes are white blood cells that help the body fight off infections.
- **low neutrophil count.** Neutrophils are white blood cells that help the body fight off infections.
- **low red blood cell count.** This may mean that you have anemia, which may make you feel weak and tired.
- **low platelet count.** Platelets help form clots and stop or prevent bleeding.

You should not take CIBINQO if your lymphocyte counts, neutrophil counts, red blood cell counts, or platelet counts are too low. Your healthcare provider may stop your CIBINQO treatment for a period of time if needed because of changes in these blood test results.

Increased cholesterol levels. You may also have increases in the amount of fat found in your blood. Your healthcare provider should check your cholesterol about 4 weeks after you start CIBINQO, and then as needed.

See **“What are the possible side effects of CIBINQO?”** for more information about side effects.

What is CIBINQO?

CIBINQO is a prescription medicine that is a Janus Kinase (JAK) inhibitor. CIBINQO is used to treat adults and children 12 years of age and older with moderate-to-severe atopic dermatitis (eczema) that did not respond to other treatment and is not well controlled with prescription therapies, including biologic medicines or when these medications cannot be tolerated.

It is not known if CIBINQO is safe and effective in children under 12 years of age.

During the first 3 months of treatment with CIBINQO, do not take CIBINQO with other medicines that prevent blood clots. You can take low-dose aspirin up to a dose of 81 mg each day during this time if prescribed by your healthcare provider.

Before taking CIBINQO, tell your healthcare provider about all of your medical conditions, including if you:

- See “**What is the most important information I should know about CIBINQO?**”
- have an infection
- are a current or past smoker
- have had a heart attack, other heart problems, or stroke
- have kidney problems or liver problems
- have low platelet counts or white blood cell counts
- have high levels of fat in your blood (high cholesterol)
- have any eye problems, including cataracts or retinal detachment.
- have recently received or are scheduled to receive an immunization (vaccine). People who take CIBINQO should not receive live vaccines.
- are pregnant or plan to become pregnant. It is not known if CIBINQO will harm your unborn baby.
 - **Pregnancy Exposure Registry.** Pfizer has a registry for women who take CIBINQO during pregnancy. The purpose of this registry is to check the health of you and your baby. If you are pregnant or become pregnant during treatment with CIBINQO, talk to your healthcare provider about how you can join this pregnancy registry, or you may contact the registry at 1-877-311-3770 or www.cibinqopregnancyregistry.com.
- are breastfeeding or plan to breastfeed. It is not known if CIBINQO passes into your breast milk. You and your healthcare provider should decide if you will take CIBINQO or breastfeed. You should not do both.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. CIBINQO and other medicines may affect each other causing side effects.

Especially tell your healthcare provider if you take aspirin or any antiplatelet therapies. See “**Do not take CIBINQO if you.**” Ask your healthcare provider if you are unsure.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist whenever you get a new medicine.

How should I take CIBINQO?

- Take CIBINQO exactly as your healthcare provider tells you to take it.
- Take CIBINQO 1 time each day, at about the same time each day.
- Swallow CIBINQO tablets whole with water. Do not split, crush, or chew the tablets.
- You can take CIBINQO with or without food.
- CIBINQO can be used with or without prescribed topical steroid medicines for atopic dermatitis. Prescribed topical medicine are lotions, creams, or ointments applied to your skin.
- If you miss a dose, take the dose as soon as possible. If it is less than 12 hours before the next dose, skip the dose. Take the next dose at your usually scheduled

time.

- If you take too much CIBINQO, call the Poison Control Center at 1-800-222-1222 right away.

What are the possible side effects of CIBINQO?

CIBINQO may cause serious side effects.

See **“What is the most important information I should know about CIBINQO?”**

The most common side effects of CIBINQO include:

- See **“What is the most important information I should know about CIBINQO.”**
- common cold
- nausea
- headache
- herpes simplex including cold sores
- increased blood level of creatine phosphokinase
- dizziness
- urinary tract infection
- tiredness
- acne
- vomiting
- mouth and throat pain
- flu
- stomach flu
- bacterial skin infection (impetigo)
- high blood pressure
- allergic skin rash to something you came into contact with
- stomach-area pain
- shingles
- low platelet count

Separation or tear to the lining of the back part of the eye (retinal detachment) has happened in people with atopic dermatitis treated with CIBINQO. Call your healthcare provider right away if you have any sudden changes in your vision during treatment with CIBINQO.

CIBINQO may cause fertility problems in females, which may affect your ability to get pregnant. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of CIBINQO.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

You may also report side effects to Pfizer at 1-800-438-1985.

How should I store CIBINQO?

- Store CIBINQO at room temperature between 68°F to 77°F (20°C to 25°C).
- Store CIBINQO in the original package.
- The container has a child resistant closure.

Keep CIBINQO and all medicines out of the reach of children.

General information about the safe and effective use of CIBINQO.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use CIBINQO for a condition for which it was not prescribed. Do not give CIBINQO to other people, even if they have the same symptoms you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about CIBINQO that is written for health professionals.

What are the ingredients in CIBINQO?

Active ingredient: abrocitinib

Inactive ingredients: dibasic calcium phosphate anhydrous, hypromellose, iron oxide red, lactose monohydrate, Macrogol, magnesium stearate, microcrystalline cellulose, sodium starch glycolate, titanium dioxide, and triacetin.



Distributed by

Pfizer Labs

Division of Pfizer Inc.

New York, NY 10017

LAB-1424-1.4

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Issued: 02/2023