

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VOXZOGO safely and effectively. See full prescribing information for VOXZOGO.

VOXZOGO (vosoritide) for injection, for subcutaneous use
Initial U.S. Approval: 2021

RECENT MAJOR CHANGES

Indication and Usage (1)	10/2023
Dosage and Administration (2.2)	10/2023
Dosage and Administration (2.4)	10/2023

INDICATIONS AND USAGE

VOXZOGO is a C type natriuretic peptide (CNP) analog indicated to increase linear growth in pediatric patients with achondroplasia with open epiphyses. This indication is approved under accelerated approval based on an improvement in annualized growth velocity. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s). (1)

DOSAGE AND ADMINISTRATION

- Ensure adequate food and fluid intake prior to administration. (2.1)
- Recommended dosage is based on patient's actual body weight. Administer VOXZOGO subcutaneously once daily. (2.2)
- Reconstitute prior to use. Injection volume is based on both patient's weight and concentration of reconstituted VOXZOGO. (2.2)
- Monitor growth and adjust dosage according to actual body weight. Permanently discontinue upon closure of epiphyses. (2.3)
- See full prescribing information for reconstitution, dilution, and administration instructions. (2.4)

DOSAGE FORMS AND STRENGTHS

For injection: 0.4 mg, 0.56 mg, or 1.2 mg of vosoritide as a lyophilized powder in a single-dose vial for reconstitution. (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

Risk of Low Blood Pressure: Transient decreases in blood pressure have been reported. Instruct patients to be well-hydrated and have adequate food intake prior to administration of VOXZOGO (5.1)

ADVERSE REACTIONS

Most common adverse reactions (>10%) are injection site erythema, injection site swelling, rash, vomiting, injection site urticaria, arthralgia, decreased blood pressure, and gastroenteritis. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact BioMarin Pharmaceutical Inc. at 1-866-906-6100, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

Renal Impairment: Not recommended in patients with eGFR < 60 mL/min/1.73 m². (8.6)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 10/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE	8.2	Lactation
2	DOSAGE AND ADMINISTRATION	8.4	Pediatric Use
2.1	Important Instructions Prior to Administration of VOXZOGO	8.6	Renal Impairment
2.2	Recommended Dosage and Administration	11	DESCRIPTION
2.3	Growth Monitoring	12	CLINICAL PHARMACOLOGY
2.4	Preparation and Administration	12.1	Mechanism of Action
3	DOSAGE FORMS AND STRENGTHS	12.2	Pharmacodynamics
4	CONTRAINDICATIONS	12.3	Pharmacokinetics
5	WARNINGS AND PRECAUTIONS	12.6	Immunogenicity
5.1	Risk of Low Blood Pressure	13	NONCLINICAL TOXICOLOGY
6	ADVERSE REACTIONS	13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
6.1	Clinical Trials Experience	14	CLINICAL STUDIES
8	USE IN SPECIFIC POPULATIONS	14.1	Pediatric Patients 5 Years of Age and Older
8.1	Pregnancy	16	HOW SUPPLIED/STORAGE AND HANDLING
		17	PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

VOXZOGO is indicated to increase linear growth in pediatric patients with achondroplasia with open epiphyses. This indication is approved under accelerated approval based on an improvement in annualized growth velocity [see *Clinical Studies (14)*]. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

2 DOSAGE AND ADMINISTRATION

2.1 Important Instructions Prior to Administration of VOXZOGO

To reduce the risk of low blood pressure and its associated signs and symptoms, instruct the caregiver and patient that the patient should [see *Warnings and Precautions (5.1)*]:

- Have adequate food intake prior to VOXZOGO administration.
- Drink approximately 240 to 300 mL of fluid in the hour prior to VOXZOGO administration.

2.2 Recommended Dosage and Administration

The recommended dosage of VOXZOGO is based on the patient's actual body weight (see Table 1). VOXZOGO is administered by subcutaneous injection once daily [see *Dosage and Administration (2.4)*].

Inject VOXZOGO at approximately the same time each day, if possible. The volume of VOXZOGO to be administered (injection volume) is based on the patient's actual body weight and the concentration of reconstituted VOXZOGO (0.8 mg/mL or 2 mg/mL) (see Table 1). VOXZOGO must be reconstituted prior to use [see *Dosage and Administration (2.4)*].

Table 1: Recommended VOXZOGO Daily Dosage and Injection Volume

Actual Body Weight*	Dose	Injection Volume	Vial Strength for Reconstitution**
3 kg	0.096 mg	0.12 mL	0.4 mg
4 kg	0.12 mg	0.15 mL	0.4 mg
5 kg	0.16 mg	0.2 mL	0.4 mg
6 to 7 kg	0.2 mg	0.25 mL	0.4 mg
8 to 11 kg	0.24 mg	0.3 mL	0.4 mg
12 to 16 kg	0.28 mg	0.35 mL	0.56 mg
17 to 21 kg	0.32 mg	0.4 mL	0.56 mg
22 to 32 kg	0.4 mg	0.5 mL	0.56 mg
33 to 43 kg	0.5 mg	0.25 mL	1.2 mg
44 to 59 kg	0.6 mg	0.3 mL	1.2 mg
60 to 89 kg	0.7 mg	0.35 mL	1.2 mg
≥ 90 kg	0.8 mg	0.4 mL	1.2 mg

* Intermediate body weights that fall within these weight bands should be rounded to the nearest whole number.

**The concentration of vosoritide in reconstituted 0.4 mg vial and 0.56 mg vial is 0.8 mg/mL. The concentration of vosoritide in reconstituted 1.2 mg vial is 2 mg/mL.

Missed dose

If a dose of VOXZOGO is missed, it can be administered within 12 hours of the scheduled time of administration. Beyond 12 hours, skip the missed dose and administer the next daily dose according to the usual dosing schedule.

2.3 Growth Monitoring

Monitor and assess patient body weight, growth, and physical development regularly every 3 to 6 months. Adjust the dosage according to the patient's actual body weight [see *Dosage and Administration (2.2)*].

Permanently discontinue VOXZOGO upon confirmation of no further growth potential, indicated by closure of epiphyses.

2.4 Preparation and Administration

Reconstitute VOXZOGO before administration using the provided diluent syringe containing Sterile Water for Injection, USP (see Reconstitution Instructions below).

Caregivers may inject VOXZOGO subcutaneously after proper training by a healthcare professional on the preparation and administration of VOXZOGO [see *Instructions for Use*].

Reconstitution Instructions

- Select the correct VOXZOGO vial strength (co-packaged with prefilled syringe with Sterile Water for Injection diluent) based on the patient's actual body weight [*see Dosage and Administration (2.2)*].
- Remove VOXZOGO vial and prefilled diluent syringe from the refrigerator and allow the vial and prefilled diluent syringe to reach room temperature before reconstituting VOXZOGO.
- Attach the diluent needle provided with ancillary supplies to the diluent prefilled syringe.
- Inject the entire diluent prefilled syringe volume into the vial (see Table 2).
- Gently swirl the diluent in the vial until the white powder is completely dissolved. Do not shake.
- Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit. Once reconstituted VOXZOGO is a clear, colorless to yellow liquid. The solution should not be used if discolored or cloudy, or if particles are present. The concentration of reconstituted solution is 0.8 mg/mL or 2.0 mg/mL (see Table 2).
- After reconstitution, VOXZOGO can be held in the vial at a room temperature 20°C to 25°C (68°F to 77°F) for a maximum of 3 hours.
- For administration, extract the required dose volume from the vial using the supplied administration syringe [*see Dosage and Administration (2.2)*].

Table 2: Dilution Requirements for VOXZOGO Prior to Administration

Vial Strength	Reconstitution Volume	Reconstituted Concentration
0.4 mg	0.5 mL	0.8 mg/mL
0.56 mg	0.7 mL	0.8 mg/mL
1.2 mg	0.6 mL	2 mg/mL

Discard any unused portion. Do not pool unused portions from the vials. Do not administer more than 1 dose from a vial. Do not mix with other medications.

Instructions for Subcutaneous Administration

See the Instructions for Use document for detailed, illustrated instructions.

- Ensure patients have had adequate food and fluid intake prior to VOXZOGO administration [*see Dosage and Administration (2.1)*]. Slowly withdraw the dosing volume of the reconstituted VOXZOGO solution from the single-dose vial into a syringe.
- Rotate sites for subcutaneous injections.
- The recommended injection sites for VOXZOGO are: the front middle of the thighs, the lower part of the abdomen at least 2 inches (5 centimeters) away from the navel, top of the buttocks or the back of the upper arms. The same injection area should not be used on two consecutive days. Do not inject VOXZOGO into sites that are red, swollen, or tender.

3 DOSAGE FORMS AND STRENGTHS

For injection: 0.4 mg, 0.56 mg, or 1.2 mg of vosoritide as a white to yellow lyophilized powder in a single-dose vial for reconstitution.

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Low Blood Pressure

Transient decreases in blood pressure were observed in clinical studies of VOXZOGO. Subjects with significant cardiac or vascular disease and patients on anti-hypertensive medicinal products were excluded from participation in VOXZOGO clinical trials. To reduce the risk of a decrease in blood pressure and associated symptoms (dizziness, fatigue and/or nausea), instruct patients to be well hydrated and have adequate food intake prior to administration of VOXZOGO [see *Dosage and Administration (2.1) and Adverse Reactions (6.1)*].

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Risk of Low Blood Pressure [see *Warnings and Precautions (5.1)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Pediatric Patients 5 Years of Age and Older

VOXZOGO was studied in a 52-week, randomized, double-blind, placebo-controlled trial in 121 subjects with achondroplasia (Study 1) [see *Clinical Studies (14)*].

The subjects' ages ranged from 5.1 to 14.9 years with a mean of 8.7 years. Sixty four (53%) subjects were male and 57 (47%) were female. Overall, 86 (71%) subjects were White, 23 (19%) were Asian, 5 (4%) were Black or African American, and 7 (6%) were classified as "multiple" race. The demographic and baseline characteristics were balanced between treatment groups. The subjects received either VOXZOGO 15 mcg/kg, or placebo administered subcutaneously once daily.

Table 3 shows adverse reactions that occurred in $\geq 5\%$ of patients treated with VOXZOGO and at a percentage greater than placebo.

Table 3: Adverse Reactions that Occurred in $\geq 5\%$ of Patients Treated with VOXZOGO and at a Percentage Greater than Placebo in Study 1*

Adverse Reaction	Placebo	VOXZOGO
	(N=61) n (%)	(N=60) n (%)
Injection site erythema	42 (69%)	45 (75%)
Injection site swelling	22 (36%)	37 (62%)
Vomiting	12 (20%)	16 (27%)
Injection site urticaria	6 (10%)	15 (25%)
Arthralgia	4 (7%)	9 (15%)
Decreased blood pressure	3 (5%)	8 (13%)
Gastroenteritis ^a	5 (8%)	8 (13%)
Diarrhea	2 (3%)	6 (10%)
Dizziness ^b	2 (3%)	6 (10%)
Ear pain	3 (5%)	6 (10%)
Influenza	3 (5%)	6 (10%)
Fatigue ^c	2 (3%)	5 (8%)
Seasonal allergy	1 (2%)	4 (7%)
Dry skin	0	3 (5%)

Abbreviations: N, total number of subjects in the treatment arm; n, number of subjects with the adverse reaction; %, percent of subjects with the adverse reaction.

* Includes adverse reactions occurring more frequently in the vosoritide arm and with a risk difference of $\geq 5\%$ (i.e., difference of >2 subjects) between treatment arms

^a Includes the preferred terms: gastroenteritis and gastroenteritis, viral

^b Includes the preferred terms: dizziness, presyncope, procedural dizziness, vertigo

^c Includes the preferred terms: fatigue, lethargy, malaise

Laboratory Abnormalities

Increase in Alkaline Phosphatase

More VOXZOGO-treated patients had an increase in alkaline phosphatase levels during the study compared to placebo (17% vs 7%).

Discussion of Selected Adverse Reactions

Decreased blood pressure

Eight (13%) of 60 subjects treated with VOXZOGO had a total of 11 events of transient decrease in blood pressure compared to 3 (5%) of 61 subjects on placebo, identified predominantly during periods of frequent monitoring at clinical visits after dosing over a 52-week treatment period. The median time to onset from injection was 31 (18 to 120) minutes with resolution within 31 (5 to 90) minutes in VOXZOGO-treated subjects. Two out of 60 (3%) VOXZOGO-treated subjects each had one symptomatic episode of decreased blood pressure with vomiting and/or dizziness compared to 0 of 61 (0%) subjects on placebo.

Injection site reactions

Injection site reactions occurred in 51 (85%) subjects receiving VOXZOGO and 50 (82%) subjects receiving placebo over a 52 week period of treatment. Injection site reactions included the preferred terms injection site erythema, injection site reaction, injection site swelling, injection site urticaria, injection site pain, injection site bruising, injection site pruritus, injection site hemorrhage, injection site discoloration, and injection site induration. Over a 52 week period, 51 (85%) of 60 subjects receiving VOXZOGO experienced a total of 6983 events of injection site reactions, while 50 (82%) of 61 subjects receiving placebo experienced a total of 1776 events of injections site reactions, representing 120.4 events per person/year exposure and 29.2 per person/year exposure, respectively. One injection site reaction event could have been associated with one or more injection site reaction symptoms (e.g., injection site swelling, injection site erythema, injection site urticaria, etc.). Two subjects in the VOXZOGO arm discontinued treatment due to adverse reactions of pain and anxiety with injections.

Pediatric Patients <5 Years

The safety of VOXZOGO in pediatric patients <5 years with achondroplasia was evaluated in a 52-week randomized, double blind, placebo-controlled study (Study 2). In this study, 64 patients from 4.4 months to <5 years of age were randomized to receive either a daily vosoritide dose with similar exposure to that characterized to be safe and effective in children with ACH aged ≥ 5 years old, or placebo. An additional 11 patients received open-label treatment as part of this study. Subjects received 30 mcg/kg while they were <2 years of age. The daily dose for subjects was adjusted to 15 mcg/kg immediately following their 2 year birthday. The most common adverse reactions (>10%) reported in pediatric patients <5 years were injection site reactions (86%) and rash (28%).

The overall safety profile of VOXZOGO in pediatric patients <5 years was similar to that seen in older pediatric patients.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no available data on vosoritide use in pregnant women to evaluate for a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. In animal reproduction studies, there was no evidence of embryo-fetal toxicity or congenital malformations when pregnant rats and rabbits were administered vosoritide subcutaneously at doses equivalent to 14-times and 200-times, respectively, the exposure at the maximum recommended human dose (MRHD) (*see Data*).

The estimated background risk of major birth defects for the indicated population is higher than the general population. The estimated background risk of miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Data

Animal Data

In an embryofetal developmental toxicity study in rats, vosoritide was administered at 90, 270, 540 mcg/kg once daily by subcutaneous injection during the period of major organogenesis from gestation day (GD) 6 – 17. There were no effects on maternal animals or on embryofetal development at the highest dose administered (14-times the exposure at the MRHD).

In an embryofetal developmental toxicity study in rabbits, vosoritide was administered at 45, 135, 240 mcg/kg once daily by subcutaneous injection during the period of major organogenesis (GD 7 – 19). No effects were observed in maternal animals or on embryofetal development at the highest dose administered (200-times the exposure at the MRHD).

In a pre- and postnatal toxicity study in rats, vosoritide was administered at 90, 270, and 540 mcg/kg once daily by subcutaneous injection during the period of major organogenesis and continuing to weaning (GD 6 through postpartum day 20). There were no effects on maternal animals, including maintenance of pregnancy, parturition, or care of offspring, and no effects were noted on offspring growth and development or ability to reproduce at the highest dose (14-times the exposure at the MRHD).

8.2 Lactation

Risk Summary

There is no information regarding the presence of vosoritide in human milk, the effects on the breastfed infant, or the effects on milk production. Vosoritide is present in rat milk. When a drug is present in animal milk, it is likely that the drug will be present in human milk. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for VOXZOGO and any potential adverse effects on the breastfed child from VOXZOGO or from the underlying maternal condition.

8.4 Pediatric Use

The safety and effectiveness of VOXZOGO have been established in pediatric patients for the improvement in linear growth in patients with achondroplasia with open epiphyses.

Use of VOXZOGO for this indication is supported by evidence from an adequate and well-controlled study in 121 pediatric patients aged 5 to 15 years with achondroplasia, pharmacokinetic data in pediatric patients aged 4.5 months to 15 years, and additional safety data in pediatric patients aged 4.4 months to <5 years [see *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)*, and *Clinical Studies (14)*].

8.6 Renal Impairment

The effect of renal impairment on the pharmacokinetics of VOXZOGO has not been evaluated. No dosage adjustment is needed for patients with $eGFR \geq 60$ mL/min/1.73 m². VOXZOGO is not recommended for patients with $eGFR < 60$ mL/min/1.73 m².

11 DESCRIPTION

VOXZOGO contains vosoritide, a human C type natriuretic peptide (CNP) analog. Vosoritide is a 39 amino acid peptide. Its amino acid sequence includes the 37 C terminal amino acids of the human CNP53 sequence plus Pro Gly on the N terminus to convey resistance to neutral endopeptidase (NEP) degradation. Vosoritide is manufactured from *Escherichia coli* using recombinant DNA technology. Vosoritide has a chemical formula of C₁₇₆H₂₉₀N₅₆O₅₁S₃ with a molecular weight of 4.1 kDa.

Vosoritide has the structural formula shown in Figure 1.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

In patients with achondroplasia, endochondral bone growth is negatively regulated due to a gain of function mutation in fibroblast growth factor receptor 3 (*FGFR3*). Binding of vosoritide to natriuretic peptide receptor-B (NPR-B) antagonizes *FGFR3* downstream signaling by inhibiting the extracellular signal-regulated kinases 1 and 2 (ERK1/2) in the mitogen-activated protein kinase (MAPK) pathway at the level of rapidly accelerating fibrosarcoma serine/threonine protein kinase (RAF-1). As a result, vosoritide, like CNP, acts as a positive regulator of endochondral bone growth as it promotes chondrocyte proliferation and differentiation.

In animal models with open growth plates, vosoritide administration resulted in the promotion of chondrocyte proliferation and differentiation that led to a widening of the growth plate and subsequent increase in skeletal growth. In the mouse models of *FGFR3*-related chondrodysplasia, a partial or complete normalization of the dwarfism phenotype was observed.

12.2 Pharmacodynamics

NPR-B Binding Activity Biomarker and Bone Metabolism Biomarker

An increase in urinary cyclic guanosine monophosphate (cGMP) concentrations from pre-dose baseline were observed within the first four hours post-dose, with a maximum level at 2 hours post-dose, after VOXZOGO administration to pediatric patients with achondroplasia.

Daily administration of VOXZOGO also led to the increase from baseline in serum collagen type X marker (CXM), an endochondral ossification biomarker and remains elevated beyond 24 months. In subjects aged 5 - 14 years old at screening, exposure-response analyses showed that vosoritide activity measured by urinary cGMP was near saturation at the dose of 15 mcg/kg once daily, while maximal increase in growth plate activity indicated by CXM was achieved at this dose.

Cardiac Electrophysiology

At the maximum approved recommended dose, VOXZOGO does not prolong the QT interval to any clinically relevant extent.

12.3 Pharmacokinetics

The area under the concentration-time curve (AUC) and peak concentration (C_{max}) of vosoritide increased greater than proportionally following subcutaneous administration to pediatric subjects with achondroplasia in the dose range of 7.5 to 30.0 mcg/kg. The pharmacokinetics of vosoritide were evaluated in 58 subjects aged 5 to 13 years with achondroplasia who received subcutaneous injections of vosoritide 15 mcg/kg once daily for 52 weeks. The mean (\pm SD) C_{max} and area under the concentration-time curve from time zero to the last measurable concentration (AUC_{0-t}) observed across 52 weeks of treatment ranged from 4.71 (\pm 2.32) to 7.18 (\pm 9.65) ng/mL, and 161 (\pm 98.1) to 290 (\pm 235) ng-min/mL, respectively. No drug accumulation was observed following 15 mcg/kg once daily dosing. The exposure of vosoritide increased with the duration of treatment. The mean AUC_{0-t} at week 52 increased approximately 20% compared to that at day 1.

Absorption

Absolute bioavailability for vosoritide following subcutaneous injection was not determined. Vosoritide was absorbed with a median T_{\max} of 15 minutes after dosing.

Distribution

The mean (\pm SD) apparent volume of distribution of vosoritide across 52 weeks of subcutaneous administration of VOXZOGO 15 mcg/kg once daily ranged from 2880 (\pm 2450) to 3020 (\pm 1980) mL/kg

Elimination

The mean (\pm SD) apparent clearance of vosoritide across 52 weeks of subcutaneous administration of VOXZOGO 15 mcg/kg once daily ranged from 79.4 (\pm 53.0) to 104 (\pm 98.8) mL/min/kg. The mean (\pm SD) half-life ranged from 21.0 (\pm 4.7) to 27.9 (\pm 9.9) minutes.

Metabolism

The metabolism of vosoritide is expected to occur via catabolic pathways with degradation into small peptide fragments and amino acids.

Specific Populations

No clinically significant differences in the vosoritide pharmacokinetics were observed based on age (0.4 to 15 years), sex or race. The effect of hepatic or renal impairment on the pharmacokinetics of vosoritide is unknown.

Body weight

Population pharmacokinetic analyses indicated that body weight is a significant covariate for vosoritide clearance and volume of distribution. The apparent clearance and volume of distribution of vosoritide increased with increasing body weight in patients with achondroplasia (5 to 74.5 kg).

Drug Interaction Studies

In vitro assessment of drug-drug interactions

In vitro studies showed that vosoritide, at therapeutic concentrations, does not inhibit or induce Cytochrome P450 enzymes. Based on *in vitro* studies, vosoritide is considered unlikely to inhibit the human drug uptake or efflux transporters such as OAT1, OAT3, OCT1, OCT2, OATP1B1, OATP1B3, MATE1, MATE2-K, BCRP, P-gp, and BSEP at clinically relevant concentrations and therefore, no effect of vosoritide is anticipated on concomitantly administered drugs that are substrates of these transporters.

In vivo assessment of drug-drug interactions

No clinical studies evaluating the drug-drug interaction potential of vosoritide have been conducted.

12.6 Immunogenicity

The observed incidence of anti-drug antibodies is highly dependent on the sensitivity and specificity of the assay. Differences in assay methods preclude meaningful comparisons of the incidence of anti-drug antibodies in the studies described below with the incidence of anti-drug antibodies in other studies, including those of vosoritide.

Of 131 subjects aged 5 years and older who were treated with VOXZOGO 15 mcg/kg/day and evaluable for the presence of anti-drug antibodies (ADA) for up to 240 weeks, ADA were detected in 35% (46/131). The earliest time to ADA development was day 85. All ADA-positive subjects tested negative for

anti-vosoritide neutralizing antibodies. There was no correlation between the number, duration, or severity of hypersensitivity adverse reactions or injection site reactions and ADA positivity or mean ADA titer. There was no association between ADA positivity or mean ADA titer and change from baseline in annual growth velocity (AGV) or height Z-score at month 12. There was no impact of serum ADA detected on the plasma PK measurements of vosoritide.

In subjects under 5 years of age, 33% (20/61) of vosoritide-treated subjects tested positive for ADA and all placebo-treated subjects tested negative for ADA for up to 44 months. The earliest time to ADA development was week 26. All of the ADA-positive subjects tested negative for neutralizing antibodies at all time points. There was no impact of ADA development on safety, efficacy or PK of vosoritide up to Month 12.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long term carcinogenicity studies and genotoxicity studies with vosoritide have not been performed.

In a fertility and reproductive study in male and female rats at doses up to 540 mcg/kg/day (15-times the exposure at the MRHD), vosoritide had no effect on mating performance, fertility, or litter characteristics.

14 CLINICAL STUDIES

14.1 Pediatric Patients 5 Years of Age and Older

The safety and effectiveness of VOXZOGO in patients with achondroplasia were assessed in one 52-week, multi-center, randomized, double-blind, placebo-controlled, phase 3 study - Study 1 (NCT03197766).

Study 1 was conducted in 121 subjects with genetically-confirmed achondroplasia, who were randomized to either VOXZOGO (N=60) or placebo (N=61). The dosage of VOXZOGO was 15 mcg/kg administered subcutaneously once daily. Baseline standing height, weight Z-score, body mass index (BMI) Z-score, and upper to lower body ratio were collected for at least 6 months prior to randomization. Subjects with limb-lengthening surgery in the prior 18 months or who planned to have limb-lengthening surgery during the study period were excluded. The study included a 52-week placebo-controlled treatment phase followed by an open-label treatment extension study period in which all subjects received VOXZOGO. The primary efficacy endpoint was the change from baseline in annualized growth velocity (AGV) at Week 52 compared with placebo.

The subjects' ages ranged from 5.1 to 14.9 years with a mean of 8.7 years. Sixty four (53%) subjects were male and 57 (47%) were female. Overall, 86 (71%) subjects were White, 23 (19%) were Asian, 5 (4%) were Black or African American, and 7 (6%) were classified as "multiple" race. The subjects had a mean baseline height standard deviation score (SDS) of -5.13.

Treatment with VOXZOGO for 52 weeks resulted in a treatment difference in the change from baseline in AGV of 1.57 cm/year after 52 weeks of treatment (Table 5).

Table 5: Annualized Growth Velocity (cm/year) at Week 52 in Subjects 5 Years of Age and Older with Achondroplasia - Study 1

	Placebo (N=61 ^a)	VOXZOGO 15 mcg/kg Daily (N=60 ^a)
Baseline mean (SD)^b	4.06 (1.20)	4.26 (1.53)
Change from baseline^c	-0.17	1.40
Difference in change of VOXZOGO – Placebo^c (95% CI)	1.57 (1.22, 1.93) ^d	

Abbreviations: AGV, annualized growth velocity; 95% CI, 95% confidence interval; LS, least-square; SD, standard deviation

^a All randomized subjects. Two patients in the VOXZOGO group discontinued from the study before Week 52. The values for these 2 patients were imputed assuming baseline growth rate for the period with missing data.

^b Baseline AGV was based on standing height at least 6 months prior to enrollment into the study.

^c LS means were estimated from the ANCOVA (analysis of covariance) model, which included treatment, stratum defined by sex and Tanner stage, baseline age, baseline AGV and baseline height Z-score.

^d 2-sided p-value <0.0001 for superiority.

The improvement in AGV in favor of VOXZOGO was consistent across all predefined subgroups analyzed including sex, age group, Tanner stage, baseline height Z-score, and baseline AGV.

Height Standard Deviation Score (SDS)

The LS mean change from baseline to Week 52 in height SDS was -0.02 in the placebo group and 0.26 in the VOXZOGO group. The difference in LS mean change from baseline was 0.28 (95% CI 0.17, 0.39; p<0.0001) in favor of VOXZOGO. The LS mean change from baseline to Week 52 in upper to lower body segment ratio was -0.02 in the placebo group and -0.03 in the VOXZOGO group. The difference in LS mean change from baseline was -0.01 (95% CI -0.05, 0.02; p=0.5).

Open-label extension

After the 52 week double blind, placebo-controlled, phase 3 study, Study 1, 58 subjects initially randomized to VOXZOGO enrolled into an open-label extension. Among the subjects who had 2 years of follow-up since randomization, the improvement in AGV was maintained.

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

VOXZOGO for injection is a white to yellow lyophilized powder for reconstitution and is provided as a co-pack which includes ten:

- Sterile, single-dose 2 mL glass vials containing VOXZOGO
- Diluent (Sterile Water for Injection, USP) in a single-dose prefilled syringe
- Diluent transfer needles (23 gauge)
- Single-dose administration syringes (30 gauge) both with needle retraction safety devices

Strength (mg)	Diluent (mL)	Co-pack NDC Number	Flip Cap Color
0.4	0.5	NDC 68135-082-36	White
0.56	0.7	NDC 68135-119-66	Magenta
1.2	0.6	NDC 68135-181-93	Grey

The following items to be obtained separately; alcohol aseptic wipes, gauze, bandages and sharps container.

Storage

Refrigerate VOXZOGO vials and prefilled diluent syringes at 2°C to 8°C (36°F to 46°F). Do not freeze.

VOXZOGO can be stored at room temperature 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) for 90 days. Do not return VOXZOGO to the refrigerator once stored at room temperature.

After reconstitution, VOXZOGO can be held in the vial at room temperature 20°C to 25°C (68°F to 77°F) for a maximum of 3 hours [*see Dosage and Administration (2.4)*].

Record the starting date of room-temperature storage clearly on the unopened product carton.

Do not use beyond expiration date on the label.

Store in the original package to protect from light.

Handling

Reconstituted VOXZOGO must be administered within 3 hours of reconstitution [*see Dosage and Administration (2.4)*].

17 PATIENT COUNSELING INFORMATION

Advise the patient and caregiver to read the FDA-approved patient labeling (Patient Information and Instructions for Use).

Preparation and Administration

Instruct caregivers on proper preparation and administration of VOXZOGO. Ensure caregivers have demonstrated the ability to perform a subcutaneous injection [*see Dosage and Administration (2.4)*].

Instruct caregivers in the technique of proper syringe and needle disposal, and advise them not to reuse these items. Instruct caregivers to dispose needles and syringes in a puncture-resistant container.

Risk of Low Blood Pressure

Inform caregivers and patients that VOXZOGO may lower blood pressure after administration. Instruct caregivers and patients that prior to VOXZOGO administration, the patient should have adequate food intake and within the hour prior to administration, the patient should drink approximately 8-10 ounces (240-300 mL) of fluid [*see Dosage and Administration (2.1) and Warnings and Precautions (5.1)*].

Manufactured for:
BioMarin Pharmaceutical Inc.
105 Digital Drive, Novato, CA 94949

PATIENT INFORMATION
VOXZOGO (vox zoe' goe)
(vosoritide)
for injection, for subcutaneous use

What is VOXZOGO?

VOXZOGO is a prescription medicine used to increase linear growth in children with achondroplasia with open bone growth plates (epiphyses).

Before you give your child VOXZOGO, tell your child's healthcare provider about all your child's medical conditions, including if they:

- have kidney problems.
- are pregnant or plan to become pregnant. It is not known if VOXZOGO will harm your child's unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if VOXZOGO passes into your child's breast milk. Talk to your child's healthcare provider about the best way to feed your child's baby if your child takes VOXZOGO.

Tell your child's healthcare provider about all the medicines your child takes, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Know the medicines your child takes. Keep a list of them to show your child's healthcare provider and pharmacist when your child gets a new medicine.

How should I give VOXZOGO?

- See the detailed **Instructions for Use** that comes with this Patient Information leaflet for instructions about the right way to store, prepare, and give VOXZOGO injections at home.
- VOXZOGO is given as an injection under the skin (subcutaneous or SC). Inject VOXZOGO 1 time every day, at about the same time each day.
- If your child's healthcare provider decides a caregiver can give the injections of VOXZOGO at home, your child's caregiver should receive training on the right way to prepare and inject VOXZOGO. Do not try to inject VOXZOGO until you have been shown the right way by your child's healthcare provider or nurse.
- Your child's healthcare provider will tell you how often you should give VOXZOGO. If your child misses a dose of VOXZOGO, it can be given within 12 hours of the scheduled time of injection. If more than 12 hours have passed, do not give the missed dose. Give the next daily dose according to your child's usual schedule.
- Your child should eat a meal and drink about 8 to 10 ounces of fluid within 1 hour before injection.
- In case you are not sure when to inject VOXZOGO, call your child's healthcare provider or pharmacist. Do not give VOXZOGO more often than as directed by your child's healthcare provider.
- Your child's dose of VOXZOGO depends on his or her body weight. Your child's healthcare provider will tell you which strength of VOXZOGO to use and how much to give your child.
- Your child's healthcare provider will monitor your child's growth and instruct you on when your child should stop VOXZOGO if they determine that your child is no longer able to grow. Stop giving VOXZOGO to your child if instructed by your child's healthcare provider.

What are the possible side effects of VOXZOGO?

VOXZOGO may cause serious side effects, including:

- **risk of low blood pressure.** VOXZOGO may temporarily lower blood pressure in some people. To help reduce the risk of low blood pressure and its symptoms (dizziness, feeling tired, or nausea), your child should eat a meal and drink about 8 to 10 ounces of fluid within 1 hour before receiving VOXZOGO.

The most common side effects of VOXZOGO include:

- injection site reactions (redness, itching, swelling, bruising, rash, hives, pain)
- high levels of blood alkaline phosphatase (shown in blood tests)
- vomiting
- joint pain
- decreased blood pressure

- stomach ache

These are not all the possible side effects of VOXZOGO. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store VOXZOGO?

- Store the VOXZOGO vial and prefilled diluent syringe in the refrigerator between 36°F to 46°F (2°C to 8°C).
- You may store VOXZOGO (before mixing) at room temperature between 68°F to 77°F (20°C to 25°C) for 90 days.
- Record the date you started storing VOXZOGO at room temperature on the carton to keep track of the 90 days.
- **Do not** return VOXZOGO to the refrigerator after it has been stored at room temperature. Throw VOXZOGO away if unused within 90 days of storing at room temperature.
- Do not use VOXZOGO past the expiration date.
- Do not freeze VOXZOGO.
- Store VOXZOGO out of direct sunlight.

Keep VOXZOGO and all medicines out of the reach of children.

General information about the safe and effective use of VOXZOGO.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use VOXZOGO for a condition for which it was not prescribed.

Do not give VOXZOGO to other people, even if they have the same symptoms you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about VOXZOGO that is written for health professionals.

What are the ingredients in VOXZOGO?

Active ingredient: vosoritide

Inactive ingredients: trehalose dihydrate, mannitol, sodium citrate dihydrate, methionine, citric acid monohydrate, and polysorbate 80

B:OMARIN[®]

BioMarin Pharmaceutical Inc.
Novato, CA 94949

© BioMarin Pharmaceutical Inc. All rights reserved.

VOXZOGO is a registered trademark of BioMarin Pharmaceutical Inc.

For more information, go to www.VOXZOGO.com or call 1-877-695-8826.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: mm/yyyy

INSTRUCTIONS FOR USE

VOXZOGO™ [Vox zoe' goe] (vosoritide)

for injection, for subcutaneous use Single-Use

This Instructions for Use contains information for caregivers on how to inject VOXZOGO.

Read this Instructions for Use before you start using VOXZOGO and each time you get a refill. There may be new information. This information does not take the place of talking to your child's healthcare provider about your child's medical condition and their treatment. Before you use VOXZOGO for the first time, make sure your child's healthcare provider shows you the right way to use it. Contact your child's healthcare provider if you or your child have any questions.

Important Information You Need to Know Before Injecting VOXZOGO

- **Wash your hands** with soap and water.
- **Do not** drop VOXZOGO or put opened items down on surfaces that are not clean.
- VOXZOGO is available in more than 1 strength. **Make sure the strength matches your prescription strength. Do not** open packaging until ready to use.
- Take the VOXZOGO vial and prefilled diluent syringe out of the refrigerator and allow them to reach room temperature before mixing.
- **Inspect the vial and supplies for any signs of damage or contamination. Do not** use if damaged or contaminated.
- **Check the expiration date.** The expiration date can be found on the carton, vial and prefilled diluent syringe. **Do not** use if expired.
- **Your child should eat a meal and drink a glass (about 8 to 10 ounces) of fluid (such as water, milk, or juice) within 1 hour before injection.**
- **VOXZOGO should be given at about the same time every day.**
- Do not mix VOXZOGO with other medicines.
- **After mixing the VOXZOGO, use it right away. Do not** use the mixed VOXZOGO if it has been sitting at room temperature for more than 3 hours. Throw it away (dispose of) in a sharps container. **See** step 18 and **“How to Throw Away (Dispose of) VOXZOGO”** for more information.
- **Do not reuse any of the supplies. After the injection, throw away (dispose of) the used vial even if there is VOXZOGO remaining. See** step 18 and **“How to Throw Away (Dispose of) VOXZOGO”** for more information.

How to Store VOXZOGO

- Store the VOXZOGO vial and prefilled diluent syringe in the refrigerator between 36°F to 46°F (2°C to 8°C).
- You may store VOXZOGO (before mixing) at room temperature between 68°F to 77°F (20°C to 25°C) for 90 days. Record the date you started storing VOXZOGO at room temperature on the carton to keep track of the 90 days. **Do not** return VOXZOGO to the refrigerator after it has been stored at room temperature. Throw VOXZOGO away if unused within 90 days of storing at room temperature.
- **Do not** freeze VOXZOGO.
- Store VOXZOGO out of direct sunlight.

Keep VOXZOGO and all other medicines out of the reach of children.

Supplies Needed to Inject VOXZOGO


Gather all of these supplies on a clean, flat surface before injecting.

Items supplied


VOXZOGO




Diluent Needle



Prefilled Diluent Syringe



Injection Syringe



Items not supplied

If you do not have these items, ask your pharmacist.


Alcohol Pads



Sharps Container



Gauze or Bandage



Preparing VOXZOGO for Injection


► Step 1

On a clean flat surface, flip off the vial cap and wipe the top with an alcohol pad.




► Step 2

Gently bend to snap off the cap from the prefilled diluent syringe.




► Step 3

Twist the diluent needle onto the prefilled diluent syringe until you can no longer twist it. Do not use the prefilled diluent syringe to give the injection.




► Step 4

Pull off the needle cap and insert the needle through the middle of the vial stopper. Slowly push the plunger rod down to inject all of the liquid.



► Step 5

Remove the needle from the vial, then press the blue tab for the needle to pull back (retract). Throw away the needle and syringe in a sharps container. See step 18 and “How to Throw Away (Dispose of) VOXZOGO.” Do not use the prefilled diluent syringe to give the injection.



► Step 6

Gently swirl the vial until the powder has completely dissolved and the solution is clear. Do not shake.




Make sure medicine is clear to yellow, not cloudy and does not have particles.

► Step 7

Take the injection syringe out of the carton. Pull off the needle cap from the injection syringe and insert the needle straight through the middle of the vial stopper. Be careful not to bend the needle.

► Step 8


Carefully hold the vial and syringe and turn the vial upside down with the needle still inserted. The vial should be on top. Be careful not to bend the needle.



Preparing VOXZOGO for Injection (continued)


► Step 9

Keep the needle tip in the medicine and slowly pull the plunger rod back to draw up the prescribed dose in the syringe. Check the prescription label for how much to draw up.



► Step 10

Remove large air bubbles in the syringe by gently tapping the syringe. Then push the bubbles back into the vial.



► Step 11

Repeat steps 9 and 10 until you have the correct prescribed dose in the syringe and no large bubbles.

Removing Bubbles

Remove any large bubbles



1 or 2 small bubbles are acceptable


Measuring Prescribed Dose

Make sure the dose in the syringe matches the prescribed dose



► Step 12

Make sure you have the prescribed dose in the syringe, then remove the vial and prepare to give the dose.



Select and Prepare Injection Site

► Step 13

VOXZOGO should be injected into the fatty layer under the skin (subcutaneous) only.

Do not inject into the same site 2 times in a row.




Do not inject through clothes.

Do not inject into skin that is swollen, sore, bruised, red, hard, or scarred.

The following sites are recommended for injection:


- **Thighs** or
- **Abdomen** (2 inches from belly button) or
- **Buttocks**
- Healthcare providers and caregivers may also inject VOXZOGO into the **back of the upper arms**.



► Step 14

Wipe the injection site with an alcohol pad and let the skin air dry.


Do not touch the area again before injecting.



Giving VOXZOGO Injection


► Step 15

After wiping the site with an alcohol pad, **pinch** the skin up around the selected injection site.



► Step 16

Quickly insert the needle all the way into the skin **at a 45-degree angle**.





► Step 17

Release the pinch and slowly push the plunger rod all the way down.

→ **Continue pressing the plunger rod until the needle retracts into the syringe.**

► Step 18

Throw away the used vial, syringes and needles in a sharps container. **See “How to Throw Away (Dispose of) VOXZOGO”** for more information.



How to Throw Away (Dispose of) VOXZOGO

Put your used or expired vials, needles and syringes in a FDA-cleared sharps disposal container right away after use. **Do not throw away (dispose of) the vials, loose needles and syringes in your household trash.**

If you do not have a FDA-cleared sharps disposal container, you may use a household container that:

- is made of a heavy-duty plastic,
- can be closed with a tight-fitting, puncture-resistant lid without sharps being able to come out,
- is upright and stable during use,
- is leak-resistant, and
- is properly labeled to warn of hazardous waste inside the container.

When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be local or state laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at:

<http://www.fda.gov/safesharpsdisposal>

Do not dispose of your used sharps disposal container in your household trash unless your community guidelines permit this. Do not recycle your used sharps disposal container.

After the Injection

- Inspect the injection site. If a small amount of bleeding occurs from the injection site, gently press a gauze pad on it for a few seconds or apply a bandage. **Do not** rub the injection site.
- Monitor for signs of low blood pressure, such as dizziness, tiredness, and nausea. If your child experiences these symptoms you should call your child's healthcare provider, then get your child to lay back with legs raised.

For Help or More Information

- Call your healthcare provider
- Call BioMarin at 1-800-123-4567
- Visit www.VOXZOGO.com

Manufactured for:
BioMarin Pharmaceutical Inc.,
Novato, CA 94949

REP-5233-C10

This Instructions for Use has been approved by the U.S. Food and Drug Administration.
Approved: November 2021