

## Medication Guide

### WEGOVY® (wee-GOH-vee)

(semaglutide) injection, for subcutaneous use

Read this Medication Guide and Instructions for Use before you start using WEGOVY and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

#### What is the most important information I should know about WEGOVY?

##### WEGOVY may cause serious side effects, including:

- **Possible thyroid tumors, including cancer.** Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rodents, WEGOVY and medicines that work like WEGOVY caused thyroid tumors, including thyroid cancer. It is not known if WEGOVY will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people.
- Do not use WEGOVY if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC), or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).

#### What is WEGOVY?

WEGOVY is an injectable prescription medicine that may help adults and children aged 12 years and older with obesity, or some adults with excess weight (overweight) who also have weight-related medical problems to help them lose weight and keep the weight off.

- WEGOVY should be used with a reduced calorie meal plan and increased physical activity.
- WEGOVY contains semaglutide and should not be used with other semaglutide-containing products or other GLP-1 receptor agonist medicines.
- It is not known if WEGOVY is safe and effective when taken with other prescription, over-the-counter, or herbal weight loss products.
- It is not known if WEGOVY can be used safely in people with a history of pancreatitis.
- It is not known if WEGOVY is safe and effective for use in children under 12 years of age.

#### Do not use WEGOVY if:

- you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC) or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).
- you have had a serious allergic reaction to semaglutide or any of the ingredients in WEGOVY. See the end of this Medication Guide for a complete list of ingredients in WEGOVY. Symptoms of a serious allergic reaction include:
  - swelling of your face, lips, tongue or throat
  - fainting or feeling dizzy
  - problems breathing or swallowing
  - very rapid heartbeat
  - severe rash or itching

#### Before using WEGOVY, tell your healthcare provider if you have any other medical conditions, including if you:

- have or have had problems with your pancreas or kidneys.
- have type 2 diabetes and a history of diabetic retinopathy.
- have or have had depression or suicidal thoughts, or mental health issues.
- are pregnant or plan to become pregnant. WEGOVY may harm your unborn baby. You should stop using WEGOVY 2 months before you plan to become pregnant.
  - **Pregnancy Exposure Registry:** There is a pregnancy exposure registry for women who use WEGOVY during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry or you may contact Novo Nordisk at 1-800-727-6500.
- are breastfeeding or plan to breastfeed. It is not known if WEGOVY passes into your breast milk. You should talk with your healthcare provider about the best way to feed your baby while using WEGOVY.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. WEGOVY may affect the way some medicines work and some medicines

may affect the way WEGOVY works. Tell your healthcare provider if you are taking other medicines to treat diabetes, including sulfonylureas or insulin. WEGOVY slows stomach emptying and can affect medicines that need to pass through the stomach quickly.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

### How should I use WEGOVY?

- Read the **Instructions for Use** that comes with WEGOVY.
- Use WEGOVY exactly as your healthcare provider tells you to.
- **Your healthcare provider should show you how to use WEGOVY before you use it for the first time.**
- WEGOVY is injected under the skin (subcutaneously) of your stomach (abdomen), thigh, or upper arm. **Do not** inject WEGOVY into a muscle (intramuscularly) or vein (intravenously).
- Change (rotate) your injection site with each injection. **Do not** use the same site for each injection.
- **Use WEGOVY 1 time each week, on the same day each week, at any time of the day.**
- Start WEGOVY with 0.25 mg per week in your first month. In your second month, increase your weekly dose to 0.5 mg. In the third month, increase your weekly dose to 1 mg. In the fourth month, increase your weekly dose to 1.7 mg and in the fifth month onwards, increase your weekly dose to the full dose of 2.4 mg. If you need to change the day of the week, you may do so as long as your last dose of WEGOVY was given **2** or more days before.
- If you miss a dose of WEGOVY and the next scheduled dose is more than 2 days away (48 hours), take the missed dose as soon as possible. If you miss a dose of WEGOVY and the next scheduled dose is less than 2 days away (48 hours), do not administer the dose. Take your next dose on the regularly scheduled day.
- If you miss doses of WEGOVY for more than 2 weeks, take your next dose on the regularly scheduled day or call your healthcare provider to talk about how to restart your treatment.
- You can take WEGOVY with or without food.
- If you take too much WEGOVY, you may have severe nausea, severe vomiting and severe low blood sugar. Call your healthcare provider or go to the nearest hospital emergency room right away if you experience any of these symptoms.

### What are the possible side effects of WEGOVY?

**WEGOVY may cause serious side effects, including:**

- **See “What is the most important information I should know about WEGOVY?”**
- **inflammation of your pancreas (pancreatitis).** Stop using WEGOVY and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back.
- **gallbladder problems.** WEGOVY may cause gallbladder problems including gallstones. Some gallbladder problems need surgery. Call your healthcare provider if you have any of the following symptoms:
  - pain in your upper stomach (abdomen)
  - fever
  - yellowing of skin or eyes (jaundice)
  - clay-colored stools
- **increased risk of low blood sugar (hypoglycemia) in patients with type 2 diabetes, especially those who also take medicines to treat type 2 diabetes mellitus such as sulfonylureas or insulin.** Low blood sugar in patients with type 2 diabetes who receive WEGOVY can be both a serious and common side effect. Talk to your healthcare provider about how to recognize and treat low blood sugar. You should check your blood sugar before you start taking WEGOVY and while you take WEGOVY. Signs and symptoms of low blood sugar may include:
  - dizziness or light-headedness
  - blurred vision
  - anxiety
  - irritability or mood changes
  - sweating
  - slurred speech
  - hunger
  - confusion or drowsiness
  - shakiness
  - weakness
  - headache
  - fast heartbeat
  - feeling jittery
- **kidney problems (kidney failure).** In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration) which may cause kidney problems to get worse. It is important for you to drink fluids to help reduce your chance of dehydration.

- **serious allergic reactions.** Stop using WEGOVY and get medical help right away, if you have any symptoms of a serious allergic reaction including:
  - swelling of your face, lips, tongue or throat
  - severe rash or itching
  - very rapid heartbeat
  - problems breathing or swallowing
  - fainting or feeling dizzy
- **change in vision in people with type 2 diabetes.** Tell your healthcare provider if you have changes in vision during treatment with WEGOVY.
- **increased heart rate.** WEGOVY can increase your heart rate while you are at rest. Your healthcare provider should check your heart rate while you take WEGOVY. Tell your healthcare provider if you feel your heart racing or pounding in your chest and it lasts for several minutes.
- **depression or thoughts of suicide.** You should pay attention to any mental changes, especially sudden changes in your mood, behaviors, thoughts, or feelings. Call your healthcare provider right away if you have any mental changes that are new, worse, or worry you.

**The most common side effects of WEGOVY in adults or children aged 12 years and older may include:**

- nausea
- stomach (abdomen) pain
- dizziness
- stomach flu
- diarrhea
- headache
- feeling bloated
- heartburn
- vomiting
- tiredness (fatigue)
- belching
- runny nose or sore throat
- constipation
- upset stomach
- gas

Talk to your healthcare provider about any side effect that bothers you or does not go away. These are not all the possible side effects of WEGOVY.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**General information about the safe and effective use of WEGOVY.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use WEGOVY for a condition for which it was not prescribed. Do not give WEGOVY to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about WEGOVY that is written for health professionals.

**What are the ingredients in WEGOVY?**

**Active Ingredient:** semaglutide

**Inactive Ingredients:** disodium phosphate dihydrate, 1.42 mg; sodium chloride, 8.25 mg; water for injection; and hydrochloric acid or sodium hydroxide may be added to adjust pH.

Manufactured by: Novo Nordisk A/S, DK-2880 Bagsvaerd, Denmark


WEGOVY® is a registered trademark of Novo Nordisk A/S.

PATENT Information: <http://novonordisk-us.com/products/product-patents.html>

© 2022 Novo Nordisk

For more information, go to [startWegovy.com](http://startWegovy.com) or call 1-833-Wegovy-1.

(b) (4)



30-000-22

30-000-22



Call 1-833-Wegovy-1

Or scan the QR code

Get support at [startWegovy.com](http://startWegovy.com)

Wegovy® is a registered trademark of Novo Nordisk AS. PATENT INFORMATION: <http://www.novonordisk-us.com/product/product-patent.html>



3 01694 52594 8

© 2022 Novo Nordisk A/S  
Novo Nordisk Inc.  
Plainsboro, NJ 08536  
1-833-Wegovy-1  
(See back of box for details)  
startWegovy.com  
Novo Nordisk A/S  
DK-2880 Bagsvaerd,  
Denmark

REFRIGERATE - DO NOT FREEZE  
Storage: Store in the refrigerator from 2°C to 8°C (36°F to 46°F). Do not freeze. If the original carton is damaged, the single-dose pen can be kept unrefrigerated from 8°C to 30°C (46°F to 86°F) up to 28 days.

Use Wegovy 1 time a week  
Write the day of the week you choose to inject

Each single-dose prefilled pen contains 0.25 mg semaglutide and will deliver one dose of 0.25 mg. If the seal is broken before first use, contact your pharmacist. Not a child-resistant container. Keep out of reach of children. Sodium hydroxide may be added to adjust pH. Sterile. 0.25 mg; and water for injection, hydrochloric acid or disodium phosphate dihydrate, 1.42 mg; sodium chloride semaglutide and the following inactive ingredients per 1 mL: Ingredients: Each 0.5 mL of Wegovy contains 0.25 mg of semaglutide and the following inactive ingredients per 1 mL: Information. Recommended Dosage See prescribing information. Use as directed by your healthcare professional.



EXPIRES

NDC 0169-4525-94

0.25 mg  
wegovy®  
(semaglutide) injection  
0.25 mg/0.5 mL  
4 Single-Dose Prefilled Pens

0.25 mg  
wegovy®  
(semaglutide) injection  
0.25 mg/0.5 mL

NDC 0169-4525-94 List 452594

Use Wegovy 1 time a week

4 Single-Dose Prefilled Pens  
Each pen delivers a single dose of 0.25 mg semaglutide  
For subcutaneous use only  
Single-Dose only  
Rx only. Sample. Not for Resale.  
Contains: 4 Wegovy pens, Product Literature.  
Dispense the enclosed Medication Guide to each patient.



NDC 0169-4525-94

0.25 mg  
wegovy®  
(semaglutide) injection  
0.25 mg/0.5 mL  
4 Single-Dose Prefilled Pens

0.25 mg

Get support at [startWegovy.com](http://startWegovy.com)

Push to open

NDC 0169-4525-94  
0.25 mg  
wegovy®  
(semaglutide) injection  
0.25 mg/0.5 mL  
4 Single-Dose Prefilled Pens


NDC 0169-4525-94  
0.25 mg  
wegovy®  
(semaglutide) injection  
0.25 mg/0.5 mL  
4 Single-Dose Prefilled Pens

1 852 13 050 1

1 852 13 050 1



(b) (4)



3 01697 50514 8

Use Wegovy 1 time a week  
Write the day of the week you choose to inject

Each single-dose prefilled pen contains 0.5 mg semaglutide and will deliver one dose of 0.5 mg. If the seal is broken before first use, contact your pharmacist. Not a child-resistant container. New powder in each of sodium hydroxide may be added to adjust pH. Sterile 0.5 mg; and water for injection, hydrochloric acid or disodium phosphate dihydrate, 1.42 mg; sodium chloride semaglutide and the following inactive ingredients per 1 mL: Ingredients: Each 0.5 mL of Wegovy contains 0.5 mg of Use as directed by your healthcare professional. Recommended Dosage See prescribing information. Subcutaneous use only

REFRIGERATE - DO NOT FREEZE  
Storage: Store in the refrigerator from 2°C to 8°C (36°F to 46°F) in original carton to protect from light. If needed prior to removal, each Wegovy single-dose pen can be kept unrefrigerated from 8°C to 30°C (46°F to 86°F) up to 28 days.

© 2022 Novo Nordisk A/S  
Novo Nordisk Inc.  
Plainsboro, NJ 08536  
1-833-Wegovy-1  
(See habla español)  
startWegovy.com  
Denmark  
Novo Nordisk A/S  
DK-2880 Bagsvaerd  
Manufactured by:

Or scan the QR code  
Call 1-833-Wegovy-1  
NOVO NORDISK

Get support at  
startWegovy.com  
Wegovy® is a registered trademark of Novo Nordisk A/S.  
PATENT INFORMATION: <http://www.novonordisk-us.com/product/product-patents.html>

00301694505148  
GTIN SERIAL OPERLOT

0.5 mg  
NDC 0169-4505-14

4 Single-Dose Prefilled Pens  
(semaglutide) injection  
0.5 mg/0.5 mL  
Wegovy®

NDC 0169-4505-14 List 450514

**0.5 mg**  
**wegovy®**  
(semaglutide) injection  
0.5 mg/0.5 mL

Use Wegovy 1 time a week

4 Single-Dose Prefilled Pens  
Each pen delivers a single dose of 0.5 mg semaglutide  
For subcutaneous use only  
Single-Dose only  
Rx only  
Contains: 4 Wegovy pens, Product Literature.  
Dispense the enclosed Medication Guide to each patient.



0.5 mg  
NDC 0169-4505-14

0.5 mg  
Wegovy®  
(semaglutide) injection  
0.5 mg/0.5 mL  
4 Single-Dose Prefilled Pens

0.5 mg  
Get support at [startWegovy.com](http://startWegovy.com)

Push to open

15/08/2022 5 14 24

15/08/2022 5 14 24

(b) (4)

22-004-30

301-31-11-9571-8



Call 1-833-Wegovy-1

Or scan the QR code

**Get support at**  
**startWegovy.com**

Wegovy® is a registered trademark of Novo Nordisk AS.  
PATENT INFORMATION: <http://www.novonordisk-us.com/product/product-patents.html>

3 0169450114 0

© 2022 Novo Nordisk AS  
Novo Nordisk Inc  
Plainsboro, NJ 08536  
1-833-Wegovy-1  
(See habla español)  
startWegovy.com  
Denmark  
Novo Nordisk A/S  
DK-2880 Bagsvaerd  
Manufactured by:

**REFRIGERATE - DO NOT FREEZE**

Storage: Store in the refrigerator from 2°C to 8°C (36°F to 46°F) in original carton to protect from light. If needed, prior to removal, each Wegovy single-dose pen can be kept unrefrigerated from 8°C to 30°C (46°F to 86°F) up to 28 days.

Use Wegovy 1 time a week

Write the day of the week you choose to inject

Each single-dose prefilled pen contains 1 mg semaglutide and will deliver one dose of 1 mg. If the seal is broken before first use, contact your pharmacist. Not a child-resistant container. See patient information leaflet for details. Sodium hydroxide may be added to adjust pH. Sterile and water for injection. Hydrochloride or disodium phosphate dihydrate, 1.42 mg; sodium chloride 8.5 mg; and water for injection. Active ingredients per 1 mL: semaglutide and the following inactive ingredients per 1 mL of injection: benzalkonium chloride, butylparaben, m-cresol, sodium hydroxide, and water for injection.

1 mg

Use as directed by your healthcare professional. Recommended Dosage See prescribing information.

Subcutaneous use only

4 Single-Dose Prefilled Pens

(semaglutide) injection  
1 mg/0.5 mL

**wegovy®**

NDC 0169-4501-14

00301694501140

GTIN SERIAL OPERLOT

1 mg

4 Single-Dose Prefilled Pens

(semaglutide) injection  
1 mg/0.5 mL

**wegovy®**

NDC 0169-4501-14

NDC 0169-4501-14

**1 mg**  
**wegovy®**  
(semaglutide) Injection  
1 mg/0.5 mL  
4 Single-Dose Prefilled Pens

NDC 0169-4501-14    List 450114

**1 mg**

**wegovy®**  
(semaglutide) injection  
1 mg/0.5 mL

Use Wegovy 1 time a week

4 Single-Dose Prefilled Pens

Each pen delivers a single dose of 1 mg semaglutide

For subcutaneous use only  
Single-Dose only

Rx only

Contains: 4 Wegovy pens, Product Literature.

Dispense the enclosed Medication Guide to each patient.



NDC 0169-4501-14

**1 mg**  
**wegovy®**  
(semaglutide) Injection  
1 mg/0.5 mL  
4 Single-Dose Prefilled Pens

1 mg

Get support at [startWegovy.com](http://startWegovy.com)

Push to open

1 mg

1 mg


1 mg

1 mg

1 mg/0.5 mL

1 mg/0.5 mL

(b) (4)



© 2022 Novo Nordisk A/S  
 Novo Nordisk Inc.  
 Plainsboro, NJ 08536  
 1-833-Wegovy-1  
 (See back of box for details)  
 startWegovy.com  
 Denmark

Manufactured by:  
 Novo Nordisk A/S  
 DK-2880 Bagsvaerd,  
 Denmark

Call 1-833-Wegovy-1  
 Or scan the QR code

Get support at  
 startWegovy.com

Wegovy® is a registered trademark of Novo Nordisk A/S.  
 PATENT INFORMATION: <http://www.novonordisk-us.com/product/product-patents.html>

REFRIGERATE - DO NOT FREEZE

Storage: Store in the refrigerator from 2°C to 8°C (36°F to 46°F) in original carton to protect from light. If needed prior to removal, each Wegovy single-dose pen can be kept unrefrigerated from 8°C to 30°C (46°F to 86°F) up to 28 days.

Use Wegovy 1 time a week  
 Write the day of the week you choose to inject

Each single-dose prefilled pen contains 1.7 mg semaglutide and will deliver one dose of 1.7 mg. If the seal is broken before first use, contact your pharmacist. Not a child-resistant container. Non-pourable and child-resistant. Hydrochloric acid or sodium hydroxide may be added to adjust pH. Sterile. 8.5 mg and water for injection, hydrochloric acid or sodium disodium phosphate dihydrate, 1.42 mg; sodium chloride semaglutide and the following inactive ingredients per 1 mL: Ingredients: Each 0.75 mL of Wegovy contains 1.7 mg of semaglutide and the following inactive ingredients per 1 mL: Information: Recommended Dosage See prescribing information. Subcutaneous use only.

3 016945174 1

1.7 mg  
 NDC 0169-4517-14

GTIN SERIAL OVERPRINT  
 0030169451741

4 Single-Dose Prefilled Pens  
 (semaglutide) injection  
 1.7 mg/0.75 mL  
 Wegovy®

NDC 0169-4517-14 List 451714

**1.7 mg**  
**wegovy®**  
 (semaglutide) injection  
 1.7 mg/0.75 mL

Use Wegovy 1 time a week

4 Single-Dose Prefilled Pens  
 Each pen delivers a single dose of 1.7 mg semaglutide  
 For subcutaneous use only  
 Single-Dose only  
 Rx only  
 Contains: 4 Wegovy pens, Product Literature.  
 Dispense the enclosed Medication Guide to each patient.

1.7 mg  
 wegovy®  
 (semaglutide) injection  
 1.7 mg/0.75 mL  
 4 Single-Dose Prefilled Pens

1.7 mg  
 wegovy®  
 (semaglutide) injection  
 1.7 mg/0.75 mL  
 4 Single-Dose Prefilled Pens

novo nordisk

1.7 mg

Get support at [startWegovy.com](http://startWegovy.com)

Push to open

1 852 31 301 2 ml 1

150/00/051 3 463 2





Ramesh  
Raghavachari

Digitally signed by Ramesh Raghavachari  
Date: 2/14/2023 11:38:39PM  
GUID: 502d0913000029f375128b0de8c50020