



TO OPEN PEEL FROM
ONE CORNER

Single-Dose Prefilled Pen

NDC 0597-0375-11

R_x only

CYLTEZO[®] PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

FOR SUBCUTANEOUS USE ONLY

Contains: One sterile single-dose prefilled pen with a fixed ½ inch length needle.

Contains no preservatives.

Needle Cap Contains Dry Natural Rubber.

Recommended Dosage: See Prescribing Information.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE.

If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator

___/___/___.

Do not use beyond expiration date.

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
US License No. 2006

BL7414E



Boehringer
Ingelheim



(01)00305970375111

10011164-01

LOT 1234567

EXP MMMYYYY



TO OPEN PEEL FROM
ONE CORNER

Single-Dose Prefilled Syringe

NDC 0597-0370-86

R_x only

CYLTEZO[®]
(adalimumab-adbm)
Injection

40 mg/0.8 mL

FOR SUBCUTANEOUS USE ONLY

Contains: One sterile single-dose prefilled syringe with a fixed ½ inch length needle.

Contains no preservatives.

Needle Cap Contains Dry Natural Rubber.

Recommended Dosage: See Prescribing Information.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light.

DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use.

Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator

___/___/___

Do not use beyond expiration date.

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
US License No. 2006

BL7420E



Boehringer
Ingelheim



(01)00305970370864

10011165-02

LOT 1234567

EXP MMMYYYY



TO OPEN PEEL FROM
ONE CORNER

Single-Dose Prefilled Syringe

NDC 0597-0405-06

R_x only

CYLTEZO[®]
(adalimumab-adbm)
Injection

20 mg/0.4 mL

FOR SUBCUTANEOUS USE ONLY

Contains: One sterile single-dose prefilled syringe with a fixed ½ inch length needle.

Contains no preservatives.

Needle Cap Contains Dry Natural Rubber.

Recommended Dosage: See Prescribing Information.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light.

DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use.

Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator

___/___/___.

Do not use beyond expiration date.

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
US License No. 2006

BL7976D



Boehringer
Ingelheim



10011163-02

LOT 1234567

EXP MMMYYYY



TO OPEN PEEL FROM
ONE CORNER

Single-Dose Prefilled Syringe

NDC 0597-0400-37

R_x only

CYLTEZO[®]
(adalimumab-adbm)
Injection

10 mg/0.2 mL

FOR SUBCUTANEOUS USE ONLY

Contains: One sterile single-dose prefilled syringe with a fixed ½ inch length needle.

Contains no preservatives.

Needle Cap Contains Dry Natural Rubber.

Recommended Dosage: See Prescribing Information.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light.

DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use.

Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator

___/___/___.

Do not use beyond expiration date.

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
US License No. 2006

BL10034C



Boehringer
Ingelheim



(01)00305970400370

10011316-02

LOT 1234567

EXP MMMYYYY

NDC 0597-0375-97

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Pens

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2023 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7417E

Do not use beyond the expiration date.



Needle Cap for Pen Contains Dry Natural Rubber.
Recommended Dosage: See Prescribing Information.

- This Carton Contains:
- 2 dose trays (each containing 1 single-dose prefilled pen with a fixed 1/2 inch length needle)
 - 2 alcohol preps
 - 1 Package Insert
 - 1 Medication Guide
 - 1 Instructions for Use

For Subcutaneous Use Only

Each Sterile Single-Dose Prefilled Pen Contains:

Adalimumab-adbm	40 mg
Glacial acetic acid	0.13 mg
Polysorbate 80	0.8 mg
Sodium acetate trihydrate	2.4 mg
Trehalose dihydrate	65.0 mg

Water for Injection, USP
Contains no preservatives.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator _____

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-97

NOTE TO PHARMACIST:
(Place Prescription Label Here)

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Pens

NDC 0597-0375-97

CYLTEZO® PEN
(adalimumab-adbm)
Injection

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



2 Single-Dose Prefilled Pens

For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-97



GTIN	00305970375975
SN	12345678901234
EXP	MMYYYY
LOT	1234567

10010940-02

ATTENTION PHARMACIST:
Each Patient is Required to Receive the
Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Pen Contains Dry Natural Rubber.

NDC 0597-0375-97

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

Rx only



2 Single-Dose Prefilled Pens







CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Pens

QUICKTIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection
 For complete instructions refer to the Instructions for Use in the carton.
 Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p> <ul style="list-style-type: none"> • Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button. • Choose a different injection site each time you inject, at least 1 inch away from the previous injection site. • Clean the injection site with an alcohol wipe. 	 <ul style="list-style-type: none"> • Inspect the pen, medicine, and expiration date. • Pull the cap straight off the pen. • Do Not twist the cap. 	 <p>Squeeze around injection site</p> <p>Press and hold 90°</p> <p>Tip</p> <ul style="list-style-type: none"> • Gently squeeze the skin around the injection site and hold firmly. • Hold the pen straight up from the injection site (at a 90-degree angle). • Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use. 	 <p>Press button 1 time and slowly count to 10</p> <p>Plunger</p> <ul style="list-style-type: none"> • Press the injection button 1 time. The 'click' sound indicates injection start. • Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10. • Do Not move the pen during injection. • You will see the plunger in the window at the bottom of the pen when the injection is complete.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

NDC 0597-0375-53

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

Product and trademark licensed from:
Boehringer Ingelheim
International GmbH
© 2023 Boehringer Ingelheim
International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim
Pharmaceuticals, Inc.,
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7418D

Citrate
Free

Do not use beyond the expiration date.



01)00305970375531

Needle Cap for Pen Contains Dry Natural Rubber.
Recommended Dosage: See Prescribing Information.

- 1 dose tray containing 1 single-dose prefilled pen with a fixed ½ inch length needle
- 2 alcohol preps
- 1 Package Insert
- 1 Medication Guide
- 1 Instructions for Use

This Carton Contains:

Each Sterile Single-Dose Prefilled Pen Contains:

Adalimumab-adbm	40 mg
Glacial acetic acid	0.13 mg
Polysorbate 80	0.8 mg
Sodium acetate trihydrate	2.4 mg
Trehalose dihydrate	65.0 mg

Water for Injection, USP.

Contains no preservatives.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days. Write the date removed from the refrigerator

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-53

NOTE TO PHARMACIST:
(Place Label Here)

1 Single-Dose Prefilled Pen

For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-53



GTIN 00305970375531
SN 12345678901234
EXP MMYYYY
LOT 1234567

NDC 0597-0375-53

CYLTEZO® PEN
(adalimumab-adbm)
Injection

ATTENTION PHARMACIST:
Each Patient is Required to Receive the
Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Pen Contains Dry Natural Rubber.

40 mg/0.8 mL

For Subcutaneous Use Only

NOT FOR SALE

Rx only

1 Single-Dose Prefilled Pen



CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

NDC 0597-0375-53





10010939-01

QUICK TIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection

For complete instructions refer to the Instructions for Use in the carton.

Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

For more information about CYLTEZO®, call **Boehringer Ingelheim Pharmaceuticals, Inc.** at 1-800-542-6257.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site Press and hold 90° Tip</p>	 <p>Press button 1 time and slowly count to 10 Plunger</p>
<ul style="list-style-type: none">• Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.• Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.• Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">• Inspect the pen, medicine, and expiration date.• Pull the cap straight off the pen.• Do Not twist the cap.	<ul style="list-style-type: none">• Gently squeeze the skin around the injection site and hold firmly.• Hold the pen straight up from the injection site (at a 90-degree angle).• Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use.	<ul style="list-style-type: none">• Press the injection button 1 time. The 'click' sound indicates injection start.• Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10.• Do Not move the pen during injection.• You will see the plunger in the window at the bottom of the pen when the injection is complete.

NDC 0597-0375-23

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2023 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7418D

4 Single-Dose Prefilled Pens

Starter Package for Psoriasis

Do not use beyond the expiration date.



For Subcutaneous Use Only

This Carton Contains:

- 4 dose trays (each containing 1 single-dose prefilled pen with a fixed 1/2 inch length needle)
- 4 alcohol preps
- 1 Package Insert
- 1 Medication Guide
- 1 Instructions for Use

Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.

Starter Package for Psoriasis

Each Sterile Single-Dose Prefilled Pen Contains:

Adalimumab-adbm	40 mg
Glacial acetic acid	0.13 mg
Polysorbate 80	0.8 mg
Sodium acetate trihydrate	2.4 mg
Trehalose dihydrate	65.0 mg

Water for Injection, USP.

Contains no preservatives.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use.

Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator

40 mg/0.8 mL



CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-23

NOTE TO PHARMACIST:
(Place Prescription Label Here)

4 Single-Dose Prefilled Pens

40 mg/0.8 mL

For Subcutaneous Use Only

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-23

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



4 Single-Dose Prefilled Pens

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

NDC 0597-0375-23



GTN
SN
12345678901234
EXP
MMYYYY
1234567



10-1461001

NDC 0597-0375-23

CYLTEZO® PEN
(adalimumab-adbm)
Injection



ATTENTION PHARMACIST:
Each Patient is Required to
Receive the Enclosed
Medication Guide.

The Entire Carton is to be
Dispensed as a Unit.

Needle Cap for Pen Contains
Dry Natural Rubber.

40 mg/0.8 mL

For Subcutaneous Use Only

Rx only



4 Single-Dose Prefilled Pens

Starter Package for Psoriasis



CYLTEZO® PEN
(adalimumab-adbm)
Injection





40 mg/0.8 mL

For Subcutaneous Use Only

4 Single-Dose Prefilled Pens

Starter Package for Psoriasis

QUICKTIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection
 For complete instructions refer to the Instructions for Use in the carton.
 Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p> <ul style="list-style-type: none"> • Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button. • Choose a different injection site each time you inject, at least 1 inch away from the previous injection site. • Clean the injection site with an alcohol wipe. 	 <ul style="list-style-type: none"> • Inspect the pen, medicine, and expiration date. • Pull the cap straight off the pen. • Do Not twist the cap. 	 <p>Squeeze around injection site</p> <p>Press and hold 90°</p> <p>Tip</p> <ul style="list-style-type: none"> • Gently squeeze the skin around the injection site and hold firmly. • Hold the pen straight up from the injection site (at a 90-degree angle). • Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use. 	 <p>Press button 1 time and slowly count to 10</p> <p>Plunger</p> <ul style="list-style-type: none"> • Press the injection button 1 time. The 'click' sound indicates injection start. • Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10. • Do Not move the pen during injection. • You will see the plunger in the window at the bottom of the pen when the injection is complete.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

NDC 0597-0375-16

CYLTEZO® PEN
(adalimumab-adbm)
Injection

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2023 **Boehringer Ingelheim International GmbH**
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT1419D

For Subcutaneous Use Only

6 Single-Dose Prefilled Pens

Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa

Do not use beyond the expiration date.



Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa

For Subcutaneous Use Only

This Carton Contains:
• 6 dose trays (each containing 1 single-dose prefilled pen with a fixed 1/2 inch length needle)
• 8 alcohol preps
• 1 Package Insert
• 1 Medication Guide
• 1 Instructions for Use

Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.

Each Sterile Single-Dose Prefilled Pen Contains:
Adalimumab-adbm.....40 mg
Glacial acetic acid.....0.13 mg
Polysorbate 80.....0.8 mg
Sodium acetate trihydrate.....2.4 mg
Trehalose dihydrate.....65.0 mg
Water for Injection, USP.

Contains no preservatives.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. **DO NOT FREEZE.** If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days. Write the date removed from the refrigerator ____/____/____.

40 mg/0.8 mL



CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-16

NOTE TO PHARMACIST:
(Place Prescription Label Here)

Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa

40 mg/0.8 mL
For Subcutaneous Use Only

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-16

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call **Boehringer Ingelheim Pharmaceuticals, Inc.** at 1-800-542-6257.



6 Single-Dose Prefilled Pens
For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-16



GTIN 00305970375166
SN 12345678901234
EXP MMYYYY
LOT 1234567



10-21961001

ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.

Needle Cap for Pen Contains Dry Natural Rubber.

NDC 0597-0375-16

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only
Rx only



6 Single-Dose Prefilled Pens

Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa







CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL
For Subcutaneous Use Only

6 Single-Dose Prefilled Pens

Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa

QUICKTIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection
For complete instructions refer to the Instructions for Use in the carton.
Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Press and hold 90°</p> <p>Tip</p>	 <p>Press button 1 time and slowly count to 10</p> <p>Plunger</p>
<ul style="list-style-type: none"> • Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button. • Choose a different injection site each time you inject, at least 1 inch away from the previous injection site. • Clean the injection site with an alcohol wipe. 	<ul style="list-style-type: none"> • Inspect the pen, medicine, and expiration date. • Pull the cap straight off the pen. • Do Not twist the cap. 	<ul style="list-style-type: none"> • Gently squeeze the skin around the injection site and hold firmly. • Hold the pen straight up from the injection site (at a 90-degree angle). • Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use. 	<ul style="list-style-type: none"> • Press the injection button 1 time. The 'click' sound indicates injection start. • Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10. • Do Not move the pen during injection. • You will see the plunger in the window at the bottom of the pen when the injection is complete.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

NDC 0597-0370-82

CYLTEZO[®]
(adalimumab-*adbm*)
Injection

Citrate
Free

For Subcutaneous Use Only

40 mg/0.8 mL

2 Single-Dose Prefilled Syringes

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2022 Boehringer Ingelheim
International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7422H

Do not use beyond the expiration date.



Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.

- This Carton Contains:
- 2 dose trays (each containing 1 single-dose prefilled syringe with a fixed 1/2 inch length needle)
 - 2 alcohol prep pads
 - 1 Package Insert
 - 1 Medication Guide
 - 1 Instructions for Use

For Subcutaneous Use Only

Each Sterile Single-Dose Prefilled Syringe Contains:

Adalimumab- <i>adbm</i>	40 mg
Glacial acetic acid.....	0.13 mg
Polysorbate 80.....	0.8 mg
Sodium acetate trihydrate.....	2.4 mg
Trehalose dihydrate.....	65.0 mg

Water for Injection, USP

Contains no preservatives.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator _____.

40 mg/0.8 mL

CYLTEZO[®]
(adalimumab-*adbm*)
Injection

NDC 0597-0370-82

NOTE TO PHARMACIST:
(Place Prescription Label Here)

2 Single-Dose Prefilled Syringes
For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO[®]
(adalimumab-*adbm*)
Injection

NDC 0597-0370-82

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



2 Single-Dose Prefilled Syringes
For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO[®]
(adalimumab-*adbm*)
Injection

NDC 0597-0370-82

NDC 0597-0370-82

CYLTEZO[®]
(adalimumab-*adbm*)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only
Rx only

2 Single-Dose Prefilled Syringes

ATTENTION PHARMACIST:
Each Patient is Required to Receive the
Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.







CYLTEZO[®]
(adalimumab-*adbm*)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

QUICKTIPS for USING CYLTEZO® (adalimumab-adbm) Injection
 For complete instructions refer to the Instructions for Use in the carton.
 Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Insert needle</p> <p>45°</p>	 <p>Plunger</p>
<ul style="list-style-type: none"> Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button. Choose a different injection site each time you inject, at least 1 inch away from the previous injection site. Clean the injection site with an alcohol wipe. 	<ul style="list-style-type: none"> Inspect the prefilled syringe, medicine, and expiration date. Pull the cap straight off the syringe. Do Not let the needle touch anything. Do Not recap the needle. 	<ul style="list-style-type: none"> Squeeze the skin around the injection site and hold firmly. Hold the syringe at about a 45 degree angle to the injection site. Insert the needle into the injection site with a quick, smooth motion. 	<ul style="list-style-type: none"> Slowly push the plunger down fully until the plunger reaches bottom of syringe. Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

Do not use beyond the expiration date.



NDC 0597-0405-80
CYLTEZO[®]
(adalimumab-adbm)
Injection

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2022 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7974G

20 mg/0.4 mL
For Subcutaneous Use Only
2 Single-Dose Prefilled Syringes



This Carton Contains:
• 2 dose trays (each containing 1 single-dose prefilled syringe with a fixed 1/2 inch length needle)
• 1 Package Insert
• 2 alcohol preps
• 1 Medication Guide
• 1 Instructions for Use
Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.

For Subcutaneous Use Only

Each Sterile Single-Dose Prefilled Syringe Contains:
Adalimumab-adbm.....20 mg
Glacial acetic acid.....0.06 mg
Polysorbate 80.....0.4 mg
Sodium acetate trihydrate.....1.21 mg
Trehalose dihydrate.....32.5 mg
Water for Injection, USP

Contains no preservatives.
Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. **DO NOT FREEZE.** If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.
Write the date removed from the refrigerator _____

20 mg/0.4 mL

CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0405-80

For Subcutaneous Use Only
2 Single-Dose Prefilled Syringes

NOTE TO PHARMACIST:
{Place Prescription Label Here}

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



NDC 0597-0405-80

CYLTEZO[®]
(adalimumab-adbm)
Injection

20 mg/0.4 mL
For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes



GTN
SN
EXP
LOT
00305970405801
12345678901234
MMYYYY
1234567

1001161-04

NDC 0597-0405-80

CYLTEZO[®]
(adalimumab-adbm)
Injection

20 mg/0.4 mL

For Subcutaneous Use Only
Rx only

2 Single-Dose Prefilled Syringes

ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.



Boehringer
Ingelheim





CYLTEZO[®]
(adalimumab-adbm)
Injection

20 mg/0.4 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

QUICKTIPS for USING CYLTEZO® (adalimumab-adbm) Injection
 For complete instructions refer to the Instructions for Use in the carton.
 Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Insert needle</p> <p>45°</p>	 <p>Plunger</p>
<ul style="list-style-type: none"> Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button. Choose a different injection site each time you inject, at least 1 inch away from the previous injection site. Clean the injection site with an alcohol wipe. 	<ul style="list-style-type: none"> Inspect the pre-filled syringe, medicine, and expiration date. Pull the cap straight off the syringe. Do Not let the needle touch anything. Do Not recap the needle. 	<ul style="list-style-type: none"> Squeeze the skin around the injection site and hold firmly. Hold the syringe at about a 45 degree angle to the injection site. Insert the needle into the injection site with a quick, smooth motion. 	<ul style="list-style-type: none"> Slowly push the plunger down fully until the plunger reaches bottom of syringe. Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

CUTSCORE

CUTSCORE

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

NDC 0597-0370-72

Product and trademark licensed from:
Boehringer Ingelheim
International GmbH

© 2022 Boehringer Ingelheim
International GmbH
ALL RIGHTS RESERVED

Manufactured by:
Boehringer Ingelheim
Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA

Product of the USA
US License Number 2006
CT7594H

CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Syringe

Citrate
Free



(01)00305970370727

This Carton Contains:
• 1 dose tray containing 1 single-dose prefilled syringe with a fixed 1/8 inch length needle
• 2 alcohol preps
• 1 Package Insert
• 1 Medication Guide
• 1 Instructions for Use

Recommended Dosage: See Prescribing Information.
Needle Cap for Syringe Contains Dry Natural Rubber.



10010943-04

Do not use beyond the expiration date.

Each Sterile Single-Dose Prefilled Syringe Contains:
Adalimumab-adbm.....40 mg
Glacial acetic acid.....0.13 mg
Polysorbate 80.....0.8 mg
Sodium acetate trihydrate.....2.4 mg
Trehalose dihydrate.....65.0 mg
Water for Injection, USP.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days. Write the date removed from the refrigerator _____

Contains no preservatives.

40 mg/0.8 mL

CYLTEZO®
(adalimumab-adbm)
Injection

NDC 0597-0370-72

1 Single-Dose Prefilled Syringe

For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO®
(adalimumab-adbm)
Injection

NDC 0597-0370-72



GTN
SN
EXP
LOT

00305970370727
12345678901234
MMMMYY
1234567



NOTE TO PHARMACIST:
{Place Label Here}

1 Single-Dose Prefilled Syringe

For Subcutaneous Use Only

40 mg/0.8 mL

Injection

CYLTEZO®
(adalimumab-adbm)



NDC 0597-0370-72

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



NDC 0597-0370-72

CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

NOT FOR SALE

Rx only

1 Single-Dose Prefilled Syringe



ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.

Needle Cap for Syringe Contains Dry Natural Rubber.



CYLTEZO®
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only





1 Single-Dose Prefilled Syringe

QUICKTIPS for USING CYLTEZO® (adalimumab-adbm) Injection

For complete instructions refer to the Instructions for Use in the carton. Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

CUTSCORE

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Insert needle</p> <p>45°</p>	 <p>Plunger</p>
<ul style="list-style-type: none">Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">Inspect the prefilled syringe, medicine, and expiration date.Pull the cap straight off the syringe. Do Not let the needle touch anything.Do Not recap the needle.	<ul style="list-style-type: none">Squeeze the skin around the injection site and hold firmly.Hold the syringe at about a 45 degree angle to the injection site.Insert the needle into the injection site with a quick, smooth motion.	<ul style="list-style-type: none">Slowly push the plunger down fully until the plunger reaches bottom of syringe.Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

CUTSCORE

Do not use beyond the expiration date.



NDC 0597-0400-89
CYLTEZO[®]
(adalimumab-adbm)
Injection
10 mg/0.2 mL
For Subcutaneous Use Only
2 Single-Dose Prefilled Syringes

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2022 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT9994C

For Subcutaneous Use Only
This Carton Contains:
• 2 dose trays (each containing 1 single-dose prefilled syringe with a fixed 1/2 inch length needle)
• 2 alcohol preps
• 1 Package Insert
• 1 Medication Guide
• 1 Instructions for Use
Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.



Each Sterile Single-Dose Prefilled Syringe Contains:
Adalimumab-adbm.....10 mg
Glacial acetic acid.....0.03 mg
Polysorbate 80.....0.2 mg
Sodium acetate trihydrate.....0.61 mg
Trehalose dihydrate.....16.3 mg
Water for Injection, USP
Contains no preservatives.
Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. **DO NOT FREEZE.** If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.
Write the date removed from the refrigerator _____/_____/_____

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



NDC 0597-0400-89
CYLTEZO[®]
(adalimumab-adbm)
Injection
10 mg/0.2 mL
For Subcutaneous Use Only
2 Single-Dose Prefilled Syringes



10 mg/0.2 mL
CYLTEZO[®]
(adalimumab-adbm)
Injection
NDC 0597-0400-89
NOTE TO PHARMACIST:
(Place Prescription Label Here)

NDC 0597-0400-89
CYLTEZO[®]
(adalimumab-adbm)
Injection
For Subcutaneous Use Only
2 Single-Dose Prefilled Syringes
GTN 00305970400899
SN 12345678901234
EXP MMYYY
LOT 1234567





NDC 0597-0400-89
CYLTEZO[®]
(adalimumab-adbm)
Injection
10 mg/0.2 mL
For Subcutaneous Use Only
Rx only
2 Single-Dose Prefilled Syringes
ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.
The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.



CYLTEZO[®]
(adalimumab-adbm)
Injection
10 mg/0.2 mL

For Subcutaneous Use Only 2 Single-Dose Prefilled Syringes

QUICKTIPS for USING CYLTEZO® (adalimumab-adbm) Injection
 For complete instructions refer to the Instructions for Use in the carton.
 Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Insert needle</p> <p>45°</p>	 <p>Plunger</p>
<ul style="list-style-type: none"> Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button. Choose a different injection site each time you inject, at least 1 inch away from the previous injection site. Clean the injection site with an alcohol wipe. 	<ul style="list-style-type: none"> Inspect the prefilled syringe, medicine, and expiration date. Pull the cap straight off the syringe. Do Not let the needle touch anything. Do Not recap the needle. 	<ul style="list-style-type: none"> Squeeze the skin around the injection site and hold firmly. Hold the syringe at about a 45 degree angle to the injection site. Insert the needle into the injection site with a quick, smooth motion. 	<ul style="list-style-type: none"> Slowly push the plunger down fully until the plunger reaches bottom of syringe. Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

CUTSCORE

CUTSCORE

NDC 0597-0375-48

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

Product and trademark licensed from:
Boehringer Ingelheim
International GmbH
© 2023, Boehringer Ingelheim
International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim
Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT113348

Citrate
Free

Do not use beyond the expiration date.



10011323-01

Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.

- 1 dose tray containing 1 single-dose prefilled pen with a fixed 1/8 inch length needle
- 2 alcohol preps
- 1 Package Insert
- 1 Medication Guide
- 1 Instructions for Use

For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-48

1 Single-Dose Prefilled Pen

For Subcutaneous Use Only

40 mg/0.8 mL

Injection

(adalimumab-adbm)



NDC 0597-0375-48



EXP

MMYYYY
1234567

10011323-01

NDC 0597-0375-48

CYLTEZO® PEN
(adalimumab-adbm)
Injection



Each Patient is Required to Receive the
Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.

Needle Cap for Pen Contains Dry Natural Rubber.

40 mg/0.8 mL

For Subcutaneous Use Only

PROFESSIONAL SAMPLE: NOT FOR SALE

Rx only

1 Single-Dose Prefilled Pen



CYLTEZO® PEN
(adalimumab-adbm)
Injection



40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

For more information about
CYLTEZO, including current
prescribing information and
Medication Guide, go to
www.cyltezo.com, scan the code,
or call Boehringer Ingelheim
Pharmaceuticals, Inc. at
1-800-542-6257.



CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen





NDC 0597-0375-48

QUICK TIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection

For complete instructions refer to the Instructions for Use in the carton.

Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p> <ul style="list-style-type: none"> Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button. Choose a different injection site each time you inject, at least 1 inch away from the previous injection site. Clean the injection site with an alcohol wipe. 	 <ul style="list-style-type: none"> Inspect the pen, medicine, and expiration date. Pull the cap straight off the pen. Do Not twist the cap. 	 <p>Squeeze around injection site</p> <p>Press and hold 90°</p> <p>Tip</p> <ul style="list-style-type: none"> Gently squeeze the skin around the injection site and hold firmly. Hold the pen straight up from the injection site (at a 90-degree angle). Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use. 	 <p>Press button 1 time and slowly count to 10</p> <p>Plunger</p> <ul style="list-style-type: none"> Press the injection button 1 time. The 'click' sound indicates injection start. Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10. Do Not move the pen during injection. You will see the plunger in the window at the bottom of the pen when the injection is complete.