NO TEXT AREA

NO TEXT AREA

Guide to Parts

Plunger Rod

Finger Grips -

Label And Expiration Date-

Syringe Barrel ·

Needle Safety Guard

Needle Safety Spring —

Gray Needle Cap On —

Important Information

or physical damage.

important information:

(subcutaneous injection)

• **Do not** shake the prefilled syringe.

prefilled syringe.

prefilled syringe.

Before Use

After Use

Important: The needle is covered by the gray needle cap before use.

Read the Patient Information for important information you need to know

• Store Fulphila in the refrigerator between 36°F to 46°F (2°C to 8°C).

• Avoid freezing. If Fulphila is accidentally frozen, allow the prefilled

• Throw away (dispose of) any Fulphila that has been left at room

more than 1 time. See Step 4: Disposing of used prefilled

• Keep the Fulphila prefilled syringe out of the reach of children.

Before you use a Fulphila prefilled syringe, read this

Keep the prefilled syringe in the original carton to protect from light

For questions about storage, contact your healthcare provider or pharmacist

• It is important that you do not try to give yourself the injection

unless you have received training from your healthcare provider.

cover the needle after the injection is given. The needle safety guard will

• The prefilled syringe has a needle safety guard that will be activated to

help prevent needlestick injuries to anyone who handles the prefilled

• Make sure that the name Fulphila appears on the carton and prefilled

You should not inject a dose of Fulphila to children weighing

• **Do not** use a prefilled syringe after the expiration date on the label.

• **Do not** use the prefilled syringe if the carton is open or damaged.

• **Do not** attempt to activate the prefilled syringe prior to injection.

• Do not attempt to remove the needle safety guard from the prefilled

• **Do not** attempt to remove the label from the prefilled syringe barrel

• **Do not** remove the gray needle cap from the prefilled syringe until you

less than 45 kg from a Fulphila prefilled syringe. A dose less than

• Fulphila is given as an injection into the tissue just under the skin

to reach room temperature before preparing an injection.

• Take Fulphila out of the refrigerator 30 minutes before use and allow it

• Do not use a Fulphila prefilled syringe that has been frozen more than

about Fulphila before using these Instructions for Use.

syringe to thaw in the refrigerator before injecting.

1 time. Use a new Fulphila prefilled syringe.

syringe after the injection has been given.

Storing the Fulphila prefilled syringe

-Used Syringe Barrel

-Used Needle

Used Needle

Safety Spring


Gray Needle Cap Off

NO TEXT AREA

NO TEXT AREA

NO TEXT AREA

BF1638/06



NO TEXT AREA

BF1638/06

6 mg/0.6 mL

Prefilled Syringe

NO TEXT AREA

NO TEXT AREA

Step 1: Gather supplies Instructions for Use

FULPHILA® (FULL-fil-ah) A - Find a clean, well-lit and flat work surface, such as a table. (pegfilgrastim-jmdb) **B** - Take the prefilled syringe carton out of the refrigerator and place it injection, for subcutaneous use on your clean work surface. Allow it to reach room temperature for Single-Dose Prefilled Syringe 30 minutes before giving an injection.

> **C** - Remove the prefilled syringe tray from the carton. **D** - Wash your hands thoroughly with soap and water.

- -Used Plunger Rod **E** - Gather the supplies for the injection:
 - 1 alcohol wipe
 - 1 cotton ball or gauze pad
 - 1 adhesive bandage an FDA-cleared sharps disposal container



Grab Here

_Label

-Exp Date

-Medicine

Gray

Needle Cap

Step 2: Prepare for injection

F - Open the tray by peeling away the cover. Grab the needle safety guard to remove the prefilled syringe from the tray.


For safety reasons:

- **Do not** grab the plunger rod. **Do not** grasp the gray needle cap.
- **G** Inspect the medicine and prefilled syringe. Make sure the medicine in the prefilled syringe is clear and colorless.

Do not use the prefilled syringe if:

- The medicine is cloudy or discolored, or contains flakes or particles.
- The prefilled syringe has been dropped.
- Any part appears cracked or broken.
- The gray needle cap is missing or not securely The expiration date printed on the label has passed.
- temperature, 68°F to 77°F (20°C to 25°C) for more than 72 hours or frozen In all cases, use a new prefilled syringe and call your healthcare provider.
 - **H** Prepare and clean the injection site.
 - There are 4 injection sites that you can use:

 - stomach area (abdomen), except for a 2-inch area right around the navel (belly button) • upper outer area of the buttocks (only if someone else is giving you the
 - the outer area of the upper arm (only if someone else is giving you the



0.6 mL (6 mg) cannot be accurately measured using the Fulphila Clean the injection site with an alcohol wipe. Let the skin dry.


- **Do not** touch this area again before injecting.
- **Do not** inject into areas where the skin is tender, bruised, red, or hard. Avoid injecting into areas with scars or stretch marks. • If you want to use the same injection site, make sure it is not the same
- spot on the injection site you used for a previous injection. • **Do not** use the prefilled syringe if it has been dropped on a hard surface. - Hold the prefilled syringe by the needle safety guard.
- The syringe may be broken even if you cannot see the break. Use a new When ready, carefully pull the gray needle cap straight off and away from the body.
 - **Do not** twist or bend the gray needle cap.
 - **Do not** hold the prefilled syringe by the plunger rod.
 - **Do not** put the gray needle cap back
 - onto the prefilled syringe. Dispose of (throw away) the gray needle cap in your household trash. 🛓

Step 3: Inject the dose

- **J** Pinch the cleaned injection site to create a firm surface.
- Keep skin pinched while injecting.




- **K** Hold the pinch. Insert the needle into the skin between 45 to 90 degrees.
- Do not touch the cleaned area of the skin



$oldsymbol{\mathsf{L}}$ - Using slow and constant pressure, push the plunger rod until it reaches the bottom.

The plunger must be pushed fully in order to inject the full dose.

- **M** Once the entire dose has been injected, the needle safety guard will be triggered. You can do either of the following:
- Release the plunger until the entire needle is covered and then remove the needle from the injection site.
- Gently remove the needle from the injection site and release the plunger until the entire needle is covered by the needle safety quard.



After releasing the plunger, the needle safety guard will safely cover the injection needle.

- Once the needle has been removed from the injection site, dispose of the syringe and needle in your sharps disposal container right away. See "Step 4: Disposing of used prefilled syringes".
- If the needle safety guard is not activated or only partially activated, discard the product (without replacing the needle cap). See "Step 4: Disposing of used prefilled syringes".
- If your injection is given by another person, they should also be careful when removing the needle from your skin in order to prevent accidental needlestick injury and possible infections.
- When you remove the syringe, if it looks like the medicine is still in the syringe barrel, this means you have not received the full dose. Call your healthcare provider right
- **N** Examine the injection site. If there is blood, press a cotton ball or gauze pad on the injection site. Do not rub the injection site. Apply an adhesive bandage if needed.

Step 4: Disposing of used prefilled syringes

- Put the used prefilled syringe in an FDA-cleared sharps disposal container right away after use. **Do not** throw away the syringe in the household
- If you do not have an FDA-cleared sharps disposal container, you may use a household container that
- made of heavy-duty plastic
- can be closed with a tight-fitting, punctureresistant lid without sharps being able to come
- upright and stable during use
- leak-resistant
- properly labeled to warn of hazardous waste inside the container
- When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at http://www.fda.gov/safesharpsdisposal.



Important: Keep the sharps disposal container out of the reach of children.

- **Do not** reuse the prefilled syringe.
- **Do not** recycle prefilled syringes or throw them into household waste.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.



Manufactured by: **Biocon Biologics Inc.** 245 Main St, 2nd Floor Cambridge, MA 02142, U.S.A. U.S. License No. 2324

KR/DRUGS/KTK/28D/07/2006 Copyright © 2023 Biocon Biologics Inc. All rights reserved.

Revised: 08/2023

Open Size: 390(W) x 600(H) mm; Folded Size: 100(W) x 40(H) mm; PAPER: 60gsm Maplitho Paper **BOOKLET TO BE SEALED WITH SELF ADHESIVE TAPE**

before injecting your dose of Fulphila.

Call your healthcare provider if you have any questions.

SAP CODE: 650006091





REFERENCE BLISTER LAYOUT NO. 1) PGB065191143 M1 (EUROPE PACK)

DRAWING No.


BOOKLET-1

PGB065191143 M1 (EUROPE PACK)

 \$\bigsize{\chi} Biocon Biologics \$\text{Siocon Biologics} \text{\$\text{\$\chi}\$}				Proof Date:	31.08.2023	DRAFT-3	3	
Description				No. of Colors	4	Page Count	1	
	(pegfilgrastim-jmdb) 6 mg - One 0.6 mL Single-dose Prefilled Syringe			Spot Print				
Item Code(s)		GTIN	TBD	Colors			(i	b) (4)
Component Type	IFU	Site Code	(b) (4)	Process Print Colors				
Barcode Info	TBD	Leatus Code	N/A	Non-Print				
TrackWise PR No.		Keyline/Drawing	See Below*	Colors				
Dimensions	390 mm x 600 mm	Fold Size	100 mm x 40mm	Minimum Font	Myriad Reg & Bold 10	/10 point		
Proof and Images	Please check thoroughly to ensure it meets your requirements as NO liability will be accepted for consequential loss. Barcodes provided on this proof are for purposes of encodation only and are not to be used for grading according to ANSI Standards. Print Vendor is responsible for ensuring barcode scanability in final printed product.							


1.14.1.1 DRAFT-LABELLING-CARTON-AND-CONTAINER-LABELS-CLEAN-PDF Reponse to Information Request Dated 17TH AUG 2023 (BLA 761075/S-025) FULPHILATM (Pegfilgrastim-jmdb) Injection 6 mg/0.6 mL

Carton label



1.14.1.1 DRAFT-LABELLING-CARTON-AND-CONTAINER-LABELS-CLEAN-PDF

Reponse to Information Request Dated 17TH AUG 2023 (BLA 761075/S-025)
FULPHILATM (Pegfilgrastim-jmdb) Injection
6 mg/0.6 mL



1.14.1.1 DRAFT-LABELLING-CARTON-AND-CONTAINER-LABELS-CLEAN-PDF

Reponse to Information Request Dated 17^{TH} AUG 2023 (BLA 761075/S-025)

FULPHILATM (Pegfilgrastim-jmdb) Injection 6 mg/0.6 mL

Tray label





Digitally signed by Susan Kirshner Date: 10/13/2023 08:25:01AM

GUID: 508da6db000266b77da0ba4bfa620030