

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ISOVUE-M safely and effectively. See full prescribing information for ISOVUE-M.

ISOVUE®-M (iopamidol) injection, for intrathecal use
Initial U.S. Approval: 1985

RECENT MAJOR CHANGES

Indications and Usage, CT cerebral ventriculography (1)-Removed 1/2026

INDICATIONS AND USAGE

ISOVUE-M is a radiographic contrast agent indicated for:

- Lumbar and thoracic myelography and computed tomography (CT) myelography in adults and pediatric patients aged 2 years and older
- Cervical and total columnar myelography in adults
- CT cisternography in adults (1)

Specific concentrations of ISOVUE-M are recommended for each type of imaging procedure. (2.2, 2.3)

DOSAGE AND ADMINISTRATION

- Individualize the volume and concentration of ISOVUE-M according to the specific dosing tables based on patient age, body weight, and study to be performed. (2.2, 2.3)
- See full prescribing information for important dosage and administration information. (2.1)

DOSAGE FORMS AND STRENGTHS

Injection: 200 mg Iodine/mL and 300 mg Iodine/mL in single-dose vials (3)

CONTRAINDICATIONS

None (4)

WARNINGS AND PRECAUTIONS

- Hypersensitivity Reactions: Life-threatening or fatal reactions can occur. Always have emergency resuscitation equipment and trained personnel available. (5.1)
- Acute Kidney Injury: Acute injury including renal failure can occur. Use the lowest dose and maintain adequate hydration to minimize risk. (5.2)
- Cardiovascular Adverse Reactions: Hemodynamic disturbances including shock and cardiac arrest may occur during or after ISOVUE-M administration. (5.4)
- Thyroid Dysfunction in Pediatric Patients 0 Years to 3 Years of Age: Individualize thyroid function monitoring based on risk factors such as prematurity. (5.6)

ADVERSE REACTIONS

Most common adverse reactions (incidence > 1%) are headache, nausea, vomiting, back pain, leg pain, neck pain, and hypotension. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Bracco Diagnostics Inc. at 1-800-257-5181 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

Lactation: A lactating woman may pump and discard breast milk for 10 hours after ISOVUE-M administration. (8.2)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 1/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Dosage and Administration Information
- 2.2 Recommended Dosage for Intrathecal Procedures in Adults
- 2.3 Recommended Dosage for Intrathecal Procedures in Pediatric Patients Aged 2 Years and Older

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Hypersensitivity Reactions
- 5.2 Acute Kidney Injury
- 5.3 Increased Risk of Seizures
- 5.4 Cardiovascular Adverse Reactions
- 5.5 Thyroid Storm in Patients with Hyperthyroidism
- 5.6 Thyroid Dysfunction in Pediatric Patients 0 Years to 3 Years of Age
- 5.7 Hypertensive Crisis in Patients with Pheochromocytoma
- 5.8 Sickle Cell Crisis in Patients with Sickle Cell Disease
- 5.9 Severe Cutaneous Adverse Reactions
- 5.10 Interference with Laboratory Test

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Drug-Drug Interactions
- 7.2 Drug-Laboratory Test Interactions

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

ISOVUE-M is indicated for:

- Lumbar and thoracic myelography, and computed tomography (CT) myelography in adults and pediatric patients aged 2 years and older
- Cervical and total columnar myelography and CT myelography in adults
- CT cisternography in adults

Specific concentrations of ISOVUE-M are recommended for each type of imaging procedure [*see Dosage and Administration (2.2, 2.3)*].

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Information

- ISOVUE-M is for intrathecal use only.
- Specific concentrations of ISOVUE-M are recommended for each type of imaging procedure [*see Dosage and Administration (2.2, 2.3)*].
- Individualize the volume, concentration, and injection rate of ISOVUE-M according to the dosing tables [*see Dosage and Administration (2.2, 2.3)*]. Consider factors such as age, body weight, anticipated pathology and degree and extent of opacification required, structure(s) or area to be examined, concomitant medical conditions, and imaging equipment and technique to be employed.
- Hydrate patients prior to and following ISOVUE-M administration [*see Warnings and Precautions (5.2)*].
- Use aseptic technique for all handling and administration of ISOVUE-M.
- ISOVUE-M may be administered at either body temperature (37°C, 98.6°F) or room temperature (20°C to 25°C, 68°F to 77°F).
- Visually inspect ISOVUE-M for particulate matter and discoloration prior to administration whenever the solution and container permit. Do not administer ISOVUE-M if particulate matter or discoloration are observed.
- Do not mix ISOVUE-M with other drugs.
- ISOVUE-M is packaged in a single-dose vial and intended for one procedure only. Discard any unused portion.

2.2 Recommended Dosage for Intrathecal Procedures in Adults

- The recommended doses in adults are shown in Table 1.
- Administer over 1 minute to 2 minutes.
- Allow at least 48 hours before repeat administration to avoid overdose; however, whenever possible, 5 days to 7 days is recommended.
- If CT myelography is performed, delay imaging by at least 4 hours to reduce the degree of contrast.

Table 1: Recommended Concentrations and Volumes of ISOVUE-M for Intrathecal Procedures in Adults

Imaging Procedure	Concentration (mg Iodine/mL)	Volume to Administer	Injection Type
Lumbar and thoracic myelography	200	10 mL to 15 mL	Lumbar
Cervical myelography	200	10 mL to 15 mL	Lumbar
		10 mL	Lateral cervical
	300	10 mL	Lumbar
Total columnar myelography	300	10 mL	Lumbar
CT cisternography	200	4 mL to 6 mL	Lumbar

2.3 Recommended Dosage for Intrathecal Procedures in Pediatric Patients Aged 2 Years and Older

- The recommended doses based on age for pediatric patients aged 2 years and older are shown in Table 2.
- Administer over 1 minute to 2 minutes.
- Allow at least 48 hours before repeat administration to avoid overdose; however, whenever possible, 5 days to 7 days is recommended.
- If CT myelography is performed, delay imaging by at least 4 hours to reduce the degree of contrast.

Table 2: Recommended Concentrations and Volumes of ISOVUE-M Based on Age for Intrathecal Procedures in Pediatric Patients Aged 2 Years and Older

Imaging Procedure	Age	Concentration (mg Iodine/mL)	Volume to Administer	Injection Type
Lumbar and thoracic myelography	2 years to 7 years	200	7 mL to 9 mL	Lumbar
	8 years to 12 years		8 mL to 11 mL	Lumbar
	13 years and older		10 mL to 12 mL	Lumbar

3 DOSAGE FORMS AND STRENGTHS

Injection: clear, colorless to pale yellow solution available in two concentrations of iodine:

Concentration (mg of Iodine/mL)	Package Size	Package Type
200	10 mL	Single-Dose Vial
300	15 mL	Single-Dose Vial

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

ISOVUE-M can cause life-threatening or fatal hypersensitivity reactions including anaphylaxis. Manifestations include respiratory arrest, laryngospasm, bronchospasm, angioedema, and shock [*see Adverse Reactions (6.2)*]. Most severe reactions develop shortly after the start of injection (e.g., within 1 to 3 minutes), but delayed reactions can also occur. There is increased risk of hypersensitivity reactions in patients with a history of previous reactions to contrast agents, and known allergic disorders (i.e., bronchial asthma, allergic rhinitis, and food allergies) or other hypersensitivities.

Premedication with antihistamines or corticosteroids does not prevent serious life-threatening reactions, but may reduce their incidence and severity. Obtain a history of allergy or hypersensitivity reactions to iodinated contrast agents and always have emergency resuscitation equipment and trained personnel available prior to ISOVUE-M administration. Monitor all patients for hypersensitivity reactions.

5.2 Acute Kidney Injury

Acute kidney injury, including renal failure, may occur after administration of iodinated contrast agents. Risk factors include pre-existing renal insufficiency, dehydration, diabetes mellitus, congestive heart failure, advanced vascular disease, elderly age, concomitant use of nephrotoxic or diuretic medications, multiple myeloma or other paraproteinemias, and repetitive or large doses of iodinated contrast agents.

Use the lowest dose of ISOVUE-M, especially in patients with risk factors for acute kidney injury. Adequately hydrate patients prior to and following ISOVUE-M administration.

5.3 Increased Risk of Seizures

Focal and generalized motor seizures have been reported after intrathecal use of iodinated contrast agents including ISOVUE-M. In several of the cases, higher than recommended doses were administered.

Use of medications that may lower the seizure threshold (phenothiazine derivatives, including those used for their antihistaminic properties; tricyclic antidepressants; MAO inhibitors; CNS stimulants; analeptics; antipsychotic agents) should be carefully evaluated. Consider discontinuing these agents at least 48 hours before and for at least 24 hours following intrathecal administration of ISOVUE-M.

5.4 Cardiovascular Adverse Reactions

Iodinated contrast agents increase the circulatory osmotic load and may induce acute or delayed hemodynamic disturbances in patients with congestive heart failure, severely impaired renal function, combined renal and hepatic disease, and combined renal and cardiac disease, particularly when repetitive or large doses are administered. Fatal cardiovascular reactions have occurred mostly within 10 minutes of

injection of iodinated contrast agent by an intravascular route; the main feature was cardiac arrest with cardiovascular disease as the main underlying factor. Hypotensive collapse and shock have occurred.

The administration of iodinated contrast agent may cause pulmonary edema in patients with heart failure. Based upon published reports, deaths associated with the administration of iodinated contrast agents range from 6.6 per 1 million (0.00066 percent) to 1 in 10,000 patients (0.01 percent).

Use the lowest necessary dose of ISOVUE-M in patients with congestive heart failure and always have emergency resuscitation equipment and trained personnel available. Monitor all patients for severe cardiovascular reactions.

5.5 Thyroid Storm in Patients with Hyperthyroidism

Thyroid storm has occurred after the use of iodinated contrast agents in patients with hyperthyroidism, or with an autonomously functioning thyroid nodule. Evaluate the risk in such patients before use of ISOVUE-M.

5.6 Thyroid Dysfunction in Pediatric Patients 0 Years to 3 Years of Age

Thyroid dysfunction characterized by hypothyroidism or transient thyroid suppression has been reported after both single exposure and multiple exposures to iodinated contrast agents in pediatric patients 0 years to 3 years of age.

Younger age, very low birth weight, prematurity, underlying medical conditions affecting thyroid function, admission to neonatal or pediatric intensive care units, and congenital cardiac conditions are associated with an increased risk of hypothyroidism after iodinated contrast agent exposure. Pediatric patients with congenital cardiac conditions may be at greatest risk given that they often require high doses of contrast during invasive cardiac procedures.

An underactive thyroid during early life may be harmful for cognitive and neurological development and may require thyroid hormone replacement therapy. After exposure to iodinated contrast agent, individualize thyroid function monitoring based on underlying risk factors, especially in term and preterm neonates. ISOVUE-M is not indicated for use in pediatric patients younger than 2 years of age [*see Use in Specific Populations (8.4)*].

5.7 Hypertensive Crisis in Patients with Pheochromocytoma

Hypertensive crisis in patients with pheochromocytoma has occurred with iodinated contrast agents. Closely monitor patients when administering ISOVUE-M if pheochromocytoma or catecholamine-secreting paragangliomas are suspected. Inject the minimum amount of ISOVUE-M necessary and have measures for treatment of hypertensive crisis readily available.

5.8 Sickle Cell Crisis in Patients with Sickle Cell Disease

Iodinated contrast agents may promote sickling in individuals who are homozygous for sickle cell disease. Hydrate patients prior to and following ISOVUE-M administration and use only if the necessary imaging information cannot be obtained with alternative imaging modalities.

5.9 Severe Cutaneous Adverse Reactions

Severe cutaneous adverse reactions (SCAR) may develop from 1 hour to several weeks after administration of iodinated contrast agent. These reactions include Stevens-Johnson syndrome and toxic epidermal necrolysis (SJS/TEN), acute generalized exanthematous pustulosis (AGEP), and drug reaction with eosinophilia and systemic symptoms (DRESS). Reaction severity may increase and time to onset may decrease with repeat administration of contrast agent; prophylactic medications may not prevent or mitigate severe cutaneous adverse reactions. Avoid administering ISOVUE-M to patients with a history of a severe cutaneous adverse reaction to ISOVUE-M.

5.10 Interference with Laboratory Test

ISOVUE-M can interfere with protein-bound iodine test [see *Drug Interactions* (7.2)].

6 ADVERSE REACTIONS

The following adverse reactions are described in greater detail in other sections:

- Hypersensitivity Reactions [see *Warnings and Precautions* (5.1)]
- Acute Kidney Injury [see *Warnings and Precautions* (5.2)]
- Increased Risk of Seizures [see *Warnings and Precautions* (5.3)]
- Cardiovascular Adverse Reactions [see *Warnings and Precautions* (5.4)]
- Thyroid Dysfunction in Pediatric Patients 0 to 3 Years of Age [see *Warnings and Precautions* (5.6)]
- Severe Cutaneous Adverse Reactions [see *Warnings and Precautions* (5.9)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of ISOVUE-M was evaluated in 686 adult patients receiving ISOVUE-M intrathecally in clinical studies. Table 3 shows the common adverse reactions (>1%).

These reactions usually occur 1 to 10 hours after injection, almost all occurring within 24 hours.

Table 3: Adverse Reactions Reported in >1% of Patients Receiving Intrathecal Injection of ISOVUE-M in Clinical Studies

Adverse Reaction	ISOVUE-M (N=686) %
Headache	16.4
Nausea	7.3
Vomiting	3.6
Back pain	2.2
Leg Pain	1.4
Neck Pain	1.1
Hypotension	1.1

The following additional adverse reactions occurred in $\leq 1\%$ of patients receiving intrathecal injection of ISOVUE-M:

Cardiovascular disorder: tachycardia, hypertension, chest pain

Gastrointestinal: diarrhea, heartburn

General disorders and administration site conditions: pyrexia, muscle weakness, hot flashes, malaise, fatigue, weakness, injection site pain

Musculoskeletal: leg cramps, sciatica, cervicobrachial irritation, meningeal irritation, radicular irritation lumbosacral, other musculoskeletal pain, involuntary movement, burning sensation

Nervous system: dizziness, paresthesia, confusion, hallucinations, lightheadedness, syncope, numbness, cold extremities, ataxia, irritability

Urogenital: urinary retention

Respiratory: dyspnea

Skin and subcutaneous tissues: rash

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of ISOVUE-M and other iopamidol products. Because the reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or to establish a causal relationship to drug exposure.

Blood and lymphatic system disorders: thrombocytopenia

Cardiovascular disorders: cardiopulmonary arrest, cardiac decompensation, arrhythmias, myocardial infarction, shock, electrocardiographic changes (e.g., increased QTc, increased R-R, increased T-wave amplitude), decreased systolic pressure, deep vein thrombosis, arterial spasms, transient ischemic attack, flushing, vasodilation, chest pain, pallor

Endocrine disorders: hyperthyroidism, hypothyroidism

Eye disorders: lacrimation increased, conjunctivitis, eye pruritus, transient blindness, visual disturbance, photophobia

Gastrointestinal disorders: retching, abdominal pain, salivary hypersecretion, salivary gland enlargement

General disorders and administration site conditions: chills, malaise

Immune system disorders: anaphylaxis characterized by cardiovascular, respiratory, and cutaneous manifestations (e.g., chest tightness, laryngeal edema, periorbital edema, facial edema); delayed hypersensitivity reactions including generalized maculopapular rash, erythema, pruritus, localized blistering, skin peeling

Infections and infestations: meningitis aseptic, meningitis bacterial as consequence of the procedural hazard

Musculoskeletal disorders: muscle spasm, musculoskeletal pain, muscular weakness

Nervous system: coma, seizure, tremors, syncope, cerebral edema, paralysis, depressed level of consciousness or loss of consciousness, meningism, encephalopathy

Psychiatric disorders: disorientation, agitation, restlessness, confusional state

Respiratory system disorders: respiratory arrest, respiratory failure, acute respiratory distress syndrome, respiratory distress, apnea, asthma, sneezing, choking, laryngeal edema, bronchospasm, rhinitis

Skin and subcutaneous tissue disorders: Stevens-Johnson syndrome and toxic epidermal necrolysis (SJS/TEN), acute generalized exanthematous pustulosis (AGEP), erythema multiforme and drug reaction with eosinophilia and systemic symptoms (DRESS), skin necrosis, face edema

7 DRUG INTERACTIONS

7.1 Drug-Drug Interactions

Metformin

In patients with renal impairment, metformin can cause lactic acidosis. Iodinated contrast agents appear to increase the risk of metformin-induced lactic acidosis, possibly as a result of worsening renal function. Stop metformin at the time of, or prior to, ISOVUE-M administration in patients with an eGFR between 30 and 60 mL/min/1.73 m² and in patients with a history of hepatic impairment, alcoholism, or heart

failure. Re-evaluate eGFR 48 hours after administration of ISOVUE-M and resume metformin only after renal function is stable.

Radioactive Iodine

Administration of iodinated contrast agents may interfere with thyroid uptake of radioactive iodine (I-131 and I-123) and decrease therapeutic and diagnostic efficacy. Avoid thyroid therapy or testing for up to 6 weeks post ISOVUE-M.

7.2 Drug-Laboratory Test Interactions

Protein-Bound Iodine Test

Iodinated contrast agents will temporarily increase protein-bound iodine in blood. Avoid protein-bound iodine test for at least 16 days following administration of ISOVUE-M. However, thyroid function tests that do not depend on iodine estimations, e.g., triiodothyronine (T₃) resin uptake and total or free thyroxine (T₄) assays, are not affected.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Available data from published literature and postmarketing cases from decades of use with iopamidol during pregnancy have not identified a drug-associated risk of major birth defects, miscarriage, or other adverse maternal or fetal outcomes. Iopamidol crosses the placenta and reaches fetal tissues in small amounts (*see Data*). In animal reproduction studies, no adverse developmental outcomes were observed with intravenous administration of iopamidol to pregnant rats and rabbits during organogenesis at doses up to 2.7 and 1.4 times, respectively, the maximum recommended human dose (*see Data*).

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Human Data

Literature reports show that intravenously administered iopamidol crosses the placenta and is visualized in the digestive tract of exposed infants after birth.

Animal Data

Iopamidol did not affect fetal development and did not induce teratogenic changes in the offspring in either rats or rabbits at the following dose levels tested: 600 mg, 1,500 mg, or 4,000 mg iodine/kg in rats, administered intravenously once a day during days 6 through 15 of pregnancy; 300 mg, 800 mg, or 2,000 mg iodine/kg in rabbits, administered intravenously once a day during days 6 through 18 of pregnancy.

8.2 Lactation

Risk Summary

There are no data on the presence of iopamidol in human milk, the effects on the breastfed infant, or the effects on milk production. Iodinated contrast agents are present unchanged in human milk in very low amounts, with poor absorption from the gastrointestinal tract of a breastfed infant. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ISOVUE-M and any potential adverse effects on the breastfed infant from ISOVUE-M or from the underlying maternal condition.

Clinical Considerations

Interruption of breastfeeding after exposure to iodinated contrast agents is not necessary because the potential exposure of the breastfed infant to iodine is small. However, a lactating woman may consider interrupting breastfeeding and pumping and discarding breast milk for 10 hours (approximately 5 half-lives) after ISOVUE-M administration in order to minimize drug exposure to a breastfed infant.

8.4 Pediatric Use

The safety and effectiveness of ISOVUE-M for lumbar and thoracic myelography and CT myelography have been established in pediatric patients aged 2 years and older.

Pediatric patients at higher risk of experiencing adverse reactions during and after any iodinated contrast agent administration may include those having asthma, sensitivity to medication or allergens, cyanotic heart disease, congestive heart failure, or serum creatinine greater than 1.5 mg/dL.

Thyroid function tests indicative of thyroid dysfunction, characterized by hypothyroidism or transient thyroid suppression have been reported following iodinated contrast agent administration in pediatric patients, including term and preterm neonates; Some patients were treated for hypothyroidism. After exposure to iodinated contrast agent, individualize thyroid function monitoring in pediatric patients 0 to 3 years of age based on underlying risk factors, especially in term and preterm neonates [see *Warnings and Precautions* (5.6) and *Adverse Reactions* (6.2)].

The safety and effectiveness of ISOVUE-M for lumbar and thoracic myelography and CT myelography have not been established in pediatric patients younger than 2 years of age.

The safety and effectiveness of ISOVUE-M for cervical and total columnar myelography and CT myelography as well as for CT cisternography have not been established in pediatric patients of any age.

8.5 Geriatric Use

Iopamidol is excreted by the kidney, and the risk of adverse reactions to ISOVUE-M may be greater in patients with renal impairment. Because patients 65 years of age and older are more likely to have renal impairment, care should be taken in dose selection, and it may be useful to monitor renal function [see *Warnings and Precautions* (5.2) and *Use in Specific Populations* (8.6)].

8.6 Renal Impairment

The clearance of iopamidol decreases with increasing degree of renal impairment. In addition, preexisting renal impairment increases the risk for acute kidney injury [see *Warnings and Precautions* (5.2)].

10 OVERDOSAGE


Doses above 3,000 mg iodine in adults and 2,400 mg iodine in pediatric patients aged 2 years and older may result in an increased frequency and severity of adverse reactions including seizures.

Treatment of an overdose is directed toward the support of all vital functions and prompt institution of symptomatic therapy. Iopamidol can be removed by dialysis.

11 DESCRIPTION

ISOVUE-M (iopamidol) injection is a radiographic contrast agent for intrathecal use.

Iopamidol is designated chemically as (S)-N,N'-bis[2-hydroxy-1-(hydroxymethyl)-ethyl]-2,4,6-triiodo-5-lactamidoisophthalamide with a molecular weight of 777.09, an empirical formula of $C_{17}H_{22}I_3N_3O_8$, and the following structural formula:



ISOVUE-M is a sterile, clear, colorless to pale yellow solution available in two concentrations of iodine:

- ISOVUE-M 200 mg iodine/mL: Each mL contains 408 mg iopamidol (providing 200 mg organically bound iodine) and the following inactive ingredients: 0.26 mg edetate calcium disodium (providing 0.029 mg sodium) and 1 mg tromethamine.
- ISOVUE-M 300 mg iodine/mL: Each mL contains 612 mg iopamidol (providing 300 mg organically bound iodine) and the following inactive ingredients: 0.39 mg edetate calcium disodium (providing 0.043 mg sodium) and 1 mg tromethamine.

The pH of ISOVUE-M has been adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.

Physicochemical characteristics are shown in Table 4. ISOVUE-M is hypertonic as compared to plasma and cerebrospinal fluid (approximately 285 and 301 mOsm/kg water, respectively).

Table 4: Physicochemical Characteristics of ISOVUE-M

Concentration (mg Iodine/mL)	200	300
Osmolality @ 37°C (mOsm/kg water)	413	616
Viscosity (cP) @ 37°C	2.0	4.7
Viscosity (cP) @ 20°C	3.3	8.8
Specific Gravity @ 37°C	1.227	1.339

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Intrathecal administration of iopamidol opacifies the body structures where the contrast agent is present, permitting their radiographic visualization through attenuation of photons.

12.2 Pharmacodynamics

Following intrathecal injection, iopamidol provides diagnostic contrast for at least 30 minutes for conventional myelography. At about 1 hour, contrast will no longer be sufficient for conventional myelography. However, diagnostic contrast for CT myelography remains at least 6 hours after administration. The exposure-response relationships and time course of pharmacodynamic response of iopamidol have not been fully characterized.

12.3 Pharmacokinetics

Absorption

Iopamidol is absorbed into the bloodstream from cerebrospinal fluid (CSF); following intrathecal administration, iopamidol appears in plasma within 1 hour and virtually all of the drug reaches the systemic circulation within 24 hours.

Distribution

Iopamidol did not bind to serum or plasma proteins at 1 hour after administration.

Elimination

The plasma half-life is approximately 2 hours; the half-life is not dose dependent.

Metabolism

Iopamidol does not undergo significant metabolism, deiodination, or biotransformation.

Excretion

Iopamidol is excreted mainly through the kidneys following intrathecal administration, and the drug is essentially undetectable in the plasma 48 hours later.

In patients with normal renal function, the cumulative urinary excretion for iopamidol, expressed as a percentage of administered intravenous dose is approximately 35% to 40% at 60 minutes, 80% to 90% at 8 hours, and 90% or more in the 72- to 96-hour period after administration. In patients with normal renal function, approximately 1% or less of the administered dose appears in cumulative 72- to 96-hour fecal specimens.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed with iopamidol to evaluate carcinogenic potential. No evidence of genetic toxicity was obtained in *in vitro* tests. In animal reproduction studies performed on rats, intravenously administered iopamidol did not induce adverse effects on fertility or general reproductive performance.

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

ISOVUE-M (iopamidol) injection is a clear, colorless to pale yellow solution available in the following presentations:

Concentration (mg Iodine/mL)	Package Size	Package Type	Sale Unit	NDC
200	10 mL	Single-Dose Vial	Carton of 10	0270-1411-11
300	15 mL	Single-Dose Vial	Carton of 10	0270-1412-15

Storage and Handling

Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature]. Protect from light.

17 PATIENT COUNSELING INFORMATION

Hypersensitivity Reactions

Advise the patient concerning the risk of hypersensitivity reactions that can occur both during and after ISOVUE-M administration. Advise the patient to report any signs or symptoms of hypersensitivity reactions during the procedure and to seek immediate medical attention for any signs or symptoms experienced after discharge [see *Warnings and Precautions* (5.1)].

Advise patients to inform their physician if they develop a rash after receiving ISOVUE-M [see *Warnings and Precautions* (5.9)].

Acute Kidney Injury

Advise the patient concerning appropriate hydration to decrease the risk of contrast induced kidney injury [see *Warnings and Precautions* (5.2)].

Thyroid Dysfunction

Advise parents/caregivers about the risk of developing thyroid dysfunction after ISOVUE-M administration. Advise parents/caregivers about when to seek medical care for their child to monitor for thyroid function [see *Warnings and Precautions* (5.6)].

Lactation

Advise lactating women that interruption of breast feeding is not necessary, however, to avoid any exposure a lactating woman may consider pumping and discarding breast milk for 10 hours after ISOVUE-M administration to minimize drug exposure to a breastfed infant [see *Use in Specific Populations* (8.2)].

Manufactured for:
Bracco Diagnostic Inc.
Princeton, NJ 08540

Manufactured by:
BIPSO GmbH
78224 Singen (Germany)

ISOVUE-M is a registered trademark of Bracco Diagnostics Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ISOVUE Imaging Bulk Package safely and effectively. See full prescribing information for ISOVUE Imaging Bulk Package.

ISOVUE® (iopamidol) injection, for intra-arterial or intravenous use
Initial U.S. Approval: 1985

WARNING: RISKS ASSOCIATED WITH INTRATHECAL ADMINISTRATION

Intrathecal administration of ISOVUE, even if inadvertent, can cause death, convulsions, cerebral hemorrhage, coma, paralysis, arachnoiditis, acute renal failure, cardiac arrest, seizures, rhabdomyolysis, hyperthermia, and brain edema. ISOVUE Imaging Bulk Package is for intra-arterial or intravenous use only. (5.1)

INDICATIONS AND USAGE

ISOVUE is a radiographic contrast agent indicated for:

Intra-arterial Procedures[†] (1.1)

- Cerebral arteriography in adults
- Peripheral arteriography in adults
- Selective visceral arteriography and aortography in adults
- Coronary arteriography and cardiac ventriculography in adults
- Angiocardiology in pediatric patients

Intravenous Procedures[†] (1.2)

- Excretory urography in adults and pediatric patients
- Computed tomography (CT) of head and body in adults and pediatric patients
- Peripheral venography in adults

[†]Specific concentrations are recommended for each type of imaging procedure. (2.2, 2.3, 2.4)

DOSAGE AND ADMINISTRATION

- Individualize the volume and concentration according to the specific dosing tables accounting for factors such as age, body weight, size of the vessel, and the rate of blood flow within the vessel. (2.2, 2.3, 2.4)
- See full prescribing information for important dosage and administration information and directions for use of imaging bulk packages. (2.1, 2.5)

DOSAGE FORMS AND STRENGTHS

Injection: 300 mg Iodine/mL and 370 mg Iodine/mL in imaging bulk packages for use only with an automated contrast injection system, contrast management system, or contrast media transfer set approved or cleared for use with these imaging bulk packages (2.5, 3)

CONTRAINDICATIONS

None (4)

WARNINGS AND PRECAUTIONS

- Hypersensitivity Reactions: Life-threatening or fatal reactions can occur. Always have emergency resuscitation equipment and trained personnel available. (5.2)
- Acute Kidney Injury: Acute injury including renal failure can occur. Use the lowest dose and maintain adequate hydration to minimize risk. (5.3)
- Cardiovascular Adverse Reactions: Hemodynamic disturbances including shock and cardiac arrest may occur during or after ISOVUE administration. (5.4)
- Thyroid Dysfunction in Pediatric Patients 0 Years to 3 Years of Age: Individualize thyroid function monitoring based on risk factors such as prematurity. (5.8)

ADVERSE REACTIONS

Most common adverse reactions (incidence >1%) from intra-arterial or intravenous use are pain, hot flashes, burning sensation, nausea, and warmth. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Bracco Diagnostics Inc. at 1-800-257-5181 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

Lactation: A lactating woman may pump and discard breast milk for 10 hours after ISOVUE administration. (8.2)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 1/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: RISKS ASSOCIATED WITH INTRATHECAL ADMINISTRATION

1 INDICATIONS AND USAGE

- 1.1 Intra-arterial Procedures
- 1.2 Intravenous Procedures

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Dosing and Administration Information
- 2.2 Recommended Dosage for Intra-arterial Procedures in Adults
- 2.3 Recommended Dosage for Intravenous Procedures in Adults
- 2.4 Recommended Dosage for Intra-arterial and Intravenous Procedures in Pediatric Patients
- 2.5 Directions for Use of Imaging Bulk Package

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Risks Associated with Intrathecal Administration
- 5.2 Hypersensitivity Reactions
- 5.3 Acute Kidney Injury
- 5.4 Cardiovascular Adverse Reactions
- 5.5 Thromboembolic Events
- 5.6 Extravasation and Injection Site Reactions
- 5.7 Thyroid Storm in Patients with Hyperthyroidism
- 5.8 Thyroid Dysfunction in Pediatric Patients 0 Years to 3 Years of Age
- 5.9 Hypertensive Crisis in Patients with Pheochromocytoma
- 5.10 Sickle Cell Crisis in Patients with Sickle Cell Disease
- 5.11 Severe Cutaneous Adverse Reactions
- 5.12 Interference with Laboratory Tests

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Drug-Drug Interactions
- 7.2 Drug-Laboratory Test Interactions

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

WARNING: RISKS ASSOCIATED WITH INTRATHECAL ADMINISTRATION

Intrathecal administration of ISOVUE, even if inadvertent, can cause death, convulsions, cerebral hemorrhage, coma, paralysis, arachnoiditis, acute renal failure, cardiac arrest, seizures, rhabdomyolysis, hyperthermia, and brain edema [see Warnings and Precautions (5.1)]. ISOVUE Imaging Bulk Package is for intra-arterial or intravenous use only [see Dosage and Administration (2.2, 2.3, 2.4)].

1 INDICATIONS AND USAGE

1.1 Intra-arterial Procedures[†]

ISOVUE is indicated for:

- Cerebral arteriography in adults
- Peripheral arteriography in adults
- Selective visceral arteriography and aortography in adults
- Coronary arteriography and cardiac ventriculography in adults
- Angiocardiology in pediatric patients

1.2 Intravenous Procedures[†]

ISOVUE is indicated for:

- Excretory urography in adults and pediatric patients
- Computerized tomography (CT) of the head and body in adults and pediatric patients
- Peripheral venography in adults

[†]Specific concentrations of ISOVUE are recommended for each type of imaging procedure [see Dosage and Administration (2.2, 2.3, 2.4)].

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosing and Administration Information

- ISOVUE Imaging Bulk Package (IBP) is for intra-arterial or intravenous use only. Do not administer intrathecally [see Warnings and Precautions (5.1)].
- Specific concentrations of ISOVUE are recommended for each type of imaging procedure [see Dosage and Administration (2.2, 2.3, 2.4)].
- Individualize the volume, concentration, and rate of administration of ISOVUE according to the specific dosing tables [see Dosage and Administration (2.2, 2.3, 2.4)]. Consider factors such as: age, body weight, blood vessel size and blood flow rate, anticipated pathology and degree and extent of opacification required, structures or area to be examined, concomitant medical conditions, imaging equipment, and technique to be employed.
- Hydrate patients before and after ISOVUE administration [see Warnings and Precautions (5.3)].
- ISOVUE may be administered at either body temperature (37°C, 98.6°F) or room temperature (20°C to 25°C, 68°F to 77°F).
- Visually inspect ISOVUE for particulate matter or discoloration before administration. Do not administer ISOVUE if particulate matter or discolorations are observed.

- Do not mix ISOVUE with other drugs or inject in intravenous lines containing other drugs or total nutritional admixtures.

2.2 Recommended Dosage for Intra-arterial Procedures in Adults

The recommended doses for intra-arterial procedures in adults are shown in Table 1.

Table 1: Recommended Concentrations and Volumes of ISOVUE for Intra-arterial Procedures in Adults

Imaging Procedure	Concentration (mg Iodine/mL)	Volume to Administer per Single Injection for Selected Injection Sites	Maximum Cumulative Total Dose
Cerebral Arteriography	300	8 mL to 12 mL by carotid puncture or transfemoral catheterization	90 mL
Peripheral Arteriography	300	5 mL to 40 mL into the femoral artery or subclavian artery 25 mL to 50 mL into the aorta for a distal runoff	250 mL
Selective Visceral Arteriography and Aortography	370	- Up to 10 mL for the renal arteries - Up to 50 mL into the larger vessels such as the aorta or celiac artery	225 mL
Coronary Arteriography and Cardiac Ventriculography	370	2 mL to 10 mL for selective coronary artery injection 25 mL to 50 mL for cardiac ventriculography or for nonselective opacification of multiple coronary arteries following injection at the aortic root	200 mL

2.3 Recommended Dosage for Intravenous Procedures in Adults

The recommended doses for intravenous procedures in adults are shown in Table 2.

Table 2: Recommended Concentrations and Volumes of ISOVUE for Intravenous Procedures in Adults

Imaging Procedure	Concentration (mg Iodine/mL)	Volume to Administer
Excretory Urography	300	50 mL to 100 mL by rapid injection
	370	50 mL by rapid injection
CT of the Head	300	100 mL to 200 mL
CT of the Body	300	100 mL to 200 mL by rapid infusion or bolus injection

	370	80 mL to 160 mL by rapid infusion or bolus injection
Peripheral Venography	300	15 mL to 100 mL per lower extremity; the maximum total dose is 230 mL

2.4 Recommended Dosage for Intra-arterial and Intravenous Procedures in Pediatric Patients

The recommended doses for intra-arterial and intravenous procedures in pediatric patients are shown in Table 3.

Table 3: Recommended Concentrations and Volumes per Body Weight of ISOVUE for Intra-arterial and Intravenous Procedures in Pediatric Patients

Imaging Procedure	Concentration (mg Iodine/mL)	Volume per Body Weight to Administer	Maximum Dose
<i>Intra-arterial Procedures</i>			
Angiocardiology	370	0.5 mL/kg to 2 mL/kg per single injection	<u>Maximum Cumulative Dose by Weight</u> • Neonates: 5 mL/kg • Aged 4 weeks and older: 8 mL/kg • Do not exceed maximum cumulative doses by age below. <u>Maximum Cumulative Dose by Age</u> • <2 years 40 mL • 2 years to 4 years 50 mL • 5 years to 9 years 100 mL • 10 years to 18 years 125 mL
<i>Intravenous Procedures</i>			
Excretory Urography	300	1 mL/kg to 3 mL/kg	100 mL
CT of the Head and Body	300	1 mL/kg to 3 mL/kg	100 mL

2.5 Directions for Use of Imaging Bulk Package

- The ISOVUE Imaging Bulk Package (IBP) is for use with an automated contrast injection system, contrast management system, or contrast media transfer set approved or cleared for use with this contrast agent in this IBP.
- See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use.
- The IBP is to be used only in a room designated for radiological procedures that involve intravascular administration of a contrast agent.
- Use aseptic technique for all handling and administration of the IBP.

- Prior to penetrating the container closure, swab the face of the container stopper with 70% isopropyl alcohol.
- Penetrate the container closure only once with a suitable sterile component of the automated contrast injection system, contrast management system, or a contrast media transfer set that is approved or cleared for use with this IBP.
- Once the IBP is punctured, do not remove it from the work area during the entire period of use. Maintain the container in an inverted position such that container contents are in continuous contact with the dispensing set.
- A maximum use time for the ISOVUE IBP is 10 hours from initial puncture. Discard any unused portion remaining in the container.
- Storage temperature of the ISOVUE IBP after the closure has been entered should not exceed 25° C (77° F); however, its content may be warmed to body temperature prior to injection.
- After the container closure is punctured, if the integrity of the IBP and the delivery system cannot be assured through direct continuous supervision, the IBP and all associated disposables for the automated contrast injection system, contrast management system, or contrast media transfer set should be discarded.

3 DOSAGE FORMS AND STRENGTHS

Injection: Clear, colorless to pale yellow solution available in the following concentrations of iodine:

Concentration (mg Iodine/mL)	Package Size	Package Type
300	200 mL	Imaging Bulk Package
	500 mL	Imaging Bulk Package
370	200 mL	Imaging Bulk Package
	500 mL	Imaging Bulk Package

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Risks Associated with Intrathecal Administration

Intrathecal administration of ISOVUE, even if inadvertent, can cause death, convulsions, cerebral hemorrhage, coma, paralysis, arachnoiditis, acute renal failure, cardiac arrest, seizures, rhabdomyolysis, hyperthermia, and brain edema. ISOVUE Imaging Bulk Package is for intra-arterial or intravenous use only [see *Dosage and Administration* (2.2, 2.3, 2.4)].

5.2 Hypersensitivity Reactions

ISOVUE can cause life-threatening or fatal hypersensitivity reactions including anaphylaxis. Manifestations include respiratory arrest, laryngospasm, bronchospasm, angioedema, and shock [see *Adverse Reactions* (6.2)]. Most severe reactions develop shortly after the start of administration (e.g.,

within 1 to 3 minutes), but delayed reactions can also occur. There is increased risk of hypersensitivity reactions in patients with a history of previous reactions to contrast agents, and allergic disorders (i.e., bronchial asthma, allergic rhinitis, and food allergies) or other hypersensitivities.

Premedication with antihistamines or corticosteroids to avoid or minimize possible allergic reactions does not prevent serious life-threatening reactions but may reduce both their incidence and severity. Obtain a history of allergy, hypersensitivity, or hypersensitivity reactions to iodinated contrast agents and always have emergency resuscitation equipment and trained personnel available prior to ISOVUE administration. Monitor all patients for hypersensitivity reactions.

5.3 Acute Kidney Injury

Acute kidney injury, including renal failure, may occur after intravascular administration of ISOVUE. Risk factors include: pre-existing renal insufficiency, dehydration, diabetes mellitus, congestive heart failure, advanced vascular disease, elderly age, concomitant use of nephrotoxic or diuretic medications, multiple myeloma or other paraproteinemias, and repetitive or large doses of ISOVUE.

Use the lowest necessary dose of ISOVUE in patients with renal impairment. Adequately hydrate patients prior to and following ISOVUE administration. Do not use laxatives, diuretics, or preparatory dehydration prior to ISOVUE administration.

5.4 Cardiovascular Adverse Reactions

Intravascular administration of ISOVUE increases the circulatory osmotic load and may induce acute or delayed hemodynamic disturbances in patients with congestive heart failure, severely impaired renal function, combined renal and hepatic disease, and combined renal and cardiac disease, particularly when repetitive or large doses are administered. Fatal cardiovascular reactions have occurred mostly within 10 minutes of ISOVUE injection; the main feature was cardiac arrest with cardiovascular disease as the main underlying factor. Hypotensive collapse and shock have occurred. Cardiac decompensation, serious arrhythmias, and myocardial ischemia or infarction can occur during coronary arteriography and ventriculography.

The administration of ISOVUE may cause pulmonary edema in patients with heart failure. Based upon published reports, deaths associated with the administration of iodinated contrast agents range from 6.6 per 1 million (0.00066 percent) to 1 in 10,000 patients (0.01 percent). Use the lowest necessary dose of ISOVUE in patients with congestive heart failure and always have emergency resuscitation equipment and trained personnel available. Monitor all patients for severe cardiovascular reactions.

5.5 Thromboembolic Events

Serious, in some cases fatal, thromboembolic events, including myocardial infarction and stroke, can occur during angiographic procedures. During these procedures, increased thrombosis and activation of the complement system occurs. Risk factors for developing thromboembolic events include: length of procedure, catheter and syringe material, underlying disease state, and concomitant medications.

To minimize thromboembolic events, use meticulous angiographic techniques and minimize the length of the procedure. Avoid blood remaining in contact with syringes containing iodinated contrast agents, which increases risk of clotting. Avoid angiocardiology in patients with homocystinuria because of the risk of inducing thrombosis and embolism.

5.6 Extravasation and Injection Site Reactions

Extravasation can occur with intravascular ISOVUE administration, particularly in patients with severe arterial or venous disease. Inflammation, blistering, skin necrosis, and compartment syndrome have been reported following extravasation. In addition, injection site reactions such as pain and swelling at the injection site can also occur [*see Adverse Reactions (6.2)*]. Ensure intravascular placement of catheters

prior to injection. Monitor patients for extravasation and advise patients to seek medical care for progression of symptoms.

5.7 Thyroid Storm in Patients with Hyperthyroidism

Thyroid storm has occurred after the intravascular use of iodinated contrast agents in patients with hyperthyroidism or with an autonomously functioning thyroid nodule. Evaluate the risk in such patients before use of ISOVUE.

5.8 Thyroid Dysfunction in Pediatric Patients 0 Years to 3 Years of Age

Thyroid dysfunction characterized by hypothyroidism or transient thyroid suppression has been reported after both single exposure and multiple exposures to iodinated contrast agents in pediatric patients 0 years to 3 years of age.

Younger age, very low birth weight, prematurity, underlying medical conditions affecting thyroid function, admission to neonatal or pediatric intensive care units, and congenital cardiac conditions are associated with an increased risk of hypothyroidism after iodinated contrast agent exposure. Pediatric patients with congenital cardiac conditions may be at greatest risk given that they often require high doses of contrast during invasive cardiac procedures.

An underactive thyroid during early life may be harmful for cognitive and neurological development and may require thyroid hormone replacement therapy. After exposure to ISOVUE, individualize thyroid function monitoring based on underlying risk factors, especially in term and preterm neonates.

5.9 Hypertensive Crisis in Patients with Pheochromocytoma

Hypertensive crisis in patients with pheochromocytoma has occurred with iodinated contrast agents. Closely monitor patients when administering ISOVUE intravascularly if pheochromocytoma or catecholamine-secreting paragangliomas are suspected. Inject the minimum amount of ISOVUE necessary and have measures for treatment of hypertensive crisis readily available.

5.10 Sickle Cell Crisis in Patients with Sickle Cell Disease

ISOVUE administered intravascularly can promote sickling in individuals who are homozygous for sickle cell disease. Hydrate patients prior to and following ISOVUE administration and use only if the necessary imaging information cannot be obtained with alternative imaging modalities.

5.11 Severe Cutaneous Adverse Reactions

Severe cutaneous adverse reactions (SCAR) may develop from 1 hour to several weeks after intravascular ISOVUE administration. These reactions include Stevens-Johnson syndrome and toxic epidermal necrolysis (SJS/TEN), acute generalized exanthematous pustulosis (AGEP), and drug reaction with eosinophilia and systemic symptoms (DRESS). Reaction severity may increase and time to onset may decrease with repeat administration of a contrast agent; prophylactic medications may not prevent or mitigate severe cutaneous adverse reactions. Avoid administering ISOVUE to patients with a history of a severe cutaneous adverse reaction to ISOVUE.

5.12 Interference with Laboratory Tests

ISOVUE can interfere with protein-bound iodine test [*see Drug Interactions (7.2)*].

6 ADVERSE REACTIONS

The following adverse reactions are described in greater detail in elsewhere in the labeling:

- Risks Associated with Intrathecal Administration [*see Warnings and Precautions (5.1)*]
- Hypersensitivity Reactions [*see Warnings and Precautions (5.2)*]

- Acute Kidney Injury [see Warnings and Precautions (5.3)]
- Cardiovascular Adverse Reactions [see Warnings and Precautions (5.4)]
- Thromboembolic Events [see Warnings and Precautions (5.5)]
- Extravasation and Injection Site Reactions [see Warnings and Precautions (5.6)]
- Thyroid Dysfunction in Pediatric Patients 0 to 3 Years of Age [see Warnings and Precautions (5.8)]
- Severe Cutaneous Adverse Reactions [see Warnings and Precautions (5.11)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse Reactions from Intra-arterial or Intravenous Use in Adults

The safety of ISOVUE was evaluated in 2,246 adult patients receiving ISOVUE by intra-arterial or intravenous route in clinical studies. Table 4 shows the common adverse reactions (>1%).

Table 4: Adverse Reactions Reported in >1% of Patients Receiving Intra-arterial or Intravenous Injection of ISOVUE in Clinical Studies

Adverse Reaction	ISOVUE (N=2,246) %
Pain	2.8
Hot flashes	1.5
Burning sensation	1.4
Nausea	1.2
Warmth	1.1

The following adverse reactions occurred in $\leq 1\%$ of patients receiving intra-arterial or intravenous injection of ISOVUE:

Cardiovascular disorders: tachycardia, hypotension, hypertension, myocardial ischemia, circulatory collapse, S-T segment depression, bigeminy, extrasystoles, ventricular fibrillation, angina pectoris, bradycardia, transient ischemic attack, thrombophlebitis

Gastrointestinal disorders: vomiting, anorexia

General disorders: headache, fever, chills, excessive sweating, back spasm

Nervous system disorders: vasovagal reaction, tingling in arms, grimace, faintness

Renal and urinary disorders: urinary retention

Respiratory: throat constriction, dyspnea, pulmonary edema

Skin and subcutaneous tissues: rash, urticaria, pruritus, flushing

Special senses: taste alterations, nasal congestion, visual disturbances

Adverse Reactions from Intra-arterial Use in Pediatric Patients

In a clinical trial with 76 pediatric patients undergoing angiocardiology, two adverse reactions (2.6%) were reported: worsening cyanosis and worsening peripheral perfusion.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of ISOVUE by intra-arterial or intravenous administration. Because the reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or to establish a causal relationship to drug exposure.

Blood and lymphatic system disorders: thrombocytopenia

Cardiovascular disorders: cardiopulmonary arrest, cardiac decompensation, arrhythmias, myocardial infarction, shock, electrocardiographic changes (e.g., increased QTc, increased R-R, increased T-wave amplitude), decreased systolic pressure, deep vein thrombosis, arterial spasms, vasodilation, chest pain, pallor

Endocrine disorders: hyperthyroidism, hypothyroidism

Eye disorders: lacrimation increased, conjunctivitis, eye pruritus, transient blindness, visual disturbance, photophobia

Gastrointestinal disorders: retching, abdominal pain, salivary hypersecretion, salivary gland enlargement

General disorders and administration site conditions: injection site pain, malaise

Immune system disorders: anaphylaxis characterized by cardiovascular, respiratory, and cutaneous manifestations (e.g., chest tightness, laryngeal edema, periorbital edema, facial edema); delayed hypersensitivity reactions including generalized maculopapular rash, erythema, pruritus, localized blistering, skin peeling

Musculoskeletal disorders: compartment syndrome following extravasation, muscle spasm, musculoskeletal pain, muscular weakness

Nervous system disorders: coma, seizure, tremors, syncope, depressed level of consciousness or loss of consciousness, encephalopathy

Psychiatric disorders: confusional state

Respiratory system disorders: respiratory arrest, respiratory failure, acute respiratory distress syndrome, respiratory distress, apnea, asthma, sneezing, choking, laryngeal edema, bronchospasm, rhinitis

Skin and subcutaneous tissue disorders: Stevens-Johnson syndrome and toxic epidermal necrolysis (SJS/TEN), acute generalized exanthematous pustulosis (AGEP), erythema multiforme and drug reaction with eosinophilia and systemic symptoms (DRESS), skin necrosis, face edema

7 DRUG INTERACTIONS

7.1 Drug-Drug Interactions

Metformin

In patients with renal impairment, metformin can cause lactic acidosis. Iodinated contrast agents appear to increase the risk of metformin-induced lactic acidosis, possibly as a result of worsening renal function. Stop metformin at the time of, or prior to, ISOVUE administration in patients with an eGFR between 30 and 60 mL/min/1.73 m²; in patients with a history of hepatic impairment, alcoholism, or heart failure; or in patients who will be administered intra-arterial iodinated contrast agents. Re-evaluate eGFR 48 hours after the imaging procedure and reinstitute metformin use only after renal function is stable.

Radioactive Iodine

Administration of ISOVUE may interfere with thyroid uptake of radioactive iodine (I-131 and I-123) and decrease therapeutic and diagnostic efficacy. Avoid thyroid therapy or testing for up to 6 weeks post ISOVUE.

7.2 Drug-Laboratory Test Interactions

Protein-Bound Iodine Test

Iodinated contrast agents, including ISOVUE, will temporarily increase protein-bound iodine in blood. Avoid protein-bound iodine test for at least 16 days following administration of ISOVUE. However, thyroid function tests that do not depend on iodine estimations, e.g., triiodothyronine (T₃) resin uptake and total or free thyroxine (T₄) assays, are not affected.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Available data from published literature and postmarketing cases from decades of use with iopamidol during pregnancy have not identified a drug-associated risk of major birth defects, miscarriage, or other adverse maternal or fetal outcomes. Iopamidol crosses the placenta and reaches fetal tissues in small amounts (*see Data*). In animal reproduction studies, no adverse developmental outcomes were observed with intravenous administration of iopamidol to pregnant rats and rabbits during organogenesis at doses up to 2.7 and 1.4 times, respectively, the maximum recommended human dose (*see Data*).

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Human Data

Literature reports show that intravenously administered iopamidol crosses the placenta and is visualized in the digestive tract of exposed infants after birth.

Animal Data

Iopamidol did not affect fetal development and did not induce teratogenic changes in the offspring in either rats or rabbits at the following dose levels tested: 600 mg, 1,500 mg, or 4,000 mg iodine/kg in rats, administered intravenously once a day during days 6 through 15 of pregnancy; 300 mg, 800 mg, or 2,000 mg iodine/kg in rabbits, administered intravenously once a day during days 6 through 18 of pregnancy.

8.2 Lactation

Risk Summary

There are no data on the presence of iopamidol in human milk, the effects on the breastfed infant, or the effects on milk production. Iodinated contrast agents are present unchanged in human milk in very low amounts, with poor absorption from the gastrointestinal tract of a breastfed infant. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ISOVUE and any potential adverse effects on the breastfed infant from ISOVUE or from the underlying maternal condition.

Clinical Considerations

Interruption of breastfeeding after exposure to iodinated contrast agents is not necessary because the potential exposure of the breastfed infant to iodine is small. However, a lactating woman may consider interrupting breastfeeding and pumping and discarding breast milk for 10 hours (approximately 5 half-lives) after ISOVUE administration in order to minimize drug exposure to a breastfed infant.

8.4 Pediatric Use

The safety and effectiveness of ISOVUE have been established in pediatric patients for intra-arterial administration for angiocardiology and for intravenous administration for excretory urography and contrast computed tomography (head and body).

Pediatric patients at higher risk of experiencing adverse reactions during and after ISOVUE administration may include those having asthma, sensitivity to medication or allergens, cyanotic heart disease, congestive heart failure, or serum creatinine greater than 1.5 mg/dL, or those less than 12 months of age.

Thyroid function tests indicative of thyroid dysfunction, characterized by hypothyroidism or transient thyroid suppression have been reported following iodinated contrast agent administration in pediatric patients, including term and preterm neonates; some patients were treated for hypothyroidism. After exposure to ISOVUE, individualize thyroid function monitoring in pediatric patients 0 years to 3 years of age based on underlying risk factors, especially in term and preterm neonates [see *Warning and Precautions* (5.8) and *Adverse Reactions* (6.2)].

The safety and effectiveness of ISOVUE for cerebral, peripheral, and selective visceral arteriography, aortography, coronary arteriography, cardiac ventriculography, and peripheral venography have not been established in pediatric patients.

8.5 Geriatric Use

Iopamidol is excreted by the kidney, and the risk of adverse reactions to ISOVUE may be greater in patients with renal impairment. Because patients 65 years of age and older are more likely to have renal impairment, care should be taken in dose selection, and it may be useful to monitor renal function [see *Warnings and Precautions* (5.3) and *Use in Specific Populations* (8.6)].

8.6 Renal Impairment

The clearance of iopamidol decreases with increasing degree of renal impairment and results in delayed opacification of the urinary system. In addition, preexisting renal impairment increases the risk for acute kidney injury [see *Warnings and Precautions* (5.3)]. Iopamidol can be removed by dialysis.


10 OVERDOSAGE

The manifestations of overdosage are life-threatening and affect mainly the pulmonary and cardiovascular systems. Treatment of an overdose is directed toward support of all vital functions and the prompt institution of symptomatic therapy. Iopamidol can be removed by dialysis.

11 DESCRIPTION

ISOVUE (iopamidol) injection is a radiographic contrast agent for intra-arterial or intravenous use.

Iopamidol is designated chemically as (S)-N,N'-bis[2-hydroxy-1-(hydroxymethyl)-ethyl]-2,4,6-triiodo-5-lactamidoisophthalamide with a molecular weight of 777.09, an empirical formula of C₁₇H₂₂I₃N₃O₈, and the following structural formula:



ISOVUE is a sterile, clear, colorless to pale yellow solution. The ISOVUE Imaging Bulk Package is available in two concentrations of iodine:

- ISOVUE 300 mg iodine/mL: Each mL contains 612 mg iopamidol (providing 300 mg bound iodine) and the following inactive ingredients: 0.39 mg edetate calcium disodium (providing 0.043 mg sodium) and 1 mg tromethamine.
- ISOVUE 370 mg iodine/mL: Each mL contains 755 mg iopamidol (providing 370 mg bound iodine) and the following inactive ingredients: 0.48 mg edetate calcium disodium (providing 0.053 mg sodium) and 1 mg tromethamine.

The pH of ISOVUE has been adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.

Physicochemical characteristics are shown in Table 5. ISOVUE is hypertonic as compared to plasma and cerebrospinal fluid (approximately 285 and 301 mOsm/kg water, respectively).

Table 5: Physicochemical Characteristics of ISOVUE

Concentration (mg Iodine/mL)	300	370
Osmolality @ 37°C (mOsm/kg water)	616	796
Viscosity (cP) @ 37°C	4.7	9.4
Viscosity (cP) @ 20°C	8.8	20.9
Specific Gravity @ 37°C	1.339	1.405

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The iodine atoms in iopamidol provide attenuation of X-rays in direct proportion to the concentration of iopamidol. Since concentration changes over time, iopamidol provides time-dependent image contrast, which may assist in visualizing body structures.

In imaging of the body, iopamidol diffuses from the vessels into the extravascular space. In normal brain with an intact blood-brain barrier, iopamidol does not diffuse into the extravascular space. In patients with a disrupted blood-brain barrier, iopamidol accumulates in the extravascular space in the region of disruption.

12.2 Pharmacodynamics

Following administration of ISOVUE, the degree of enhancement is related to the iodine concentration in the tissue of interest. However, the exposure-response relationships and time course of pharmacodynamic response of iopamidol have not been fully characterized.

12.3 Pharmacokinetics

Distribution

After intravascular administration, plasma concentrations of iodine fall within 5 to 10 minutes due to distribution into the vascular and extracellular fluid compartments. Equilibration with the extracellular compartments is reached in about 10 minutes.

The apparent volume of distribution suggests that iopamidol is distributed evenly between blood and extracellular fluid. Iopamidol may be visualized in the renal parenchyma within 30 to 60 seconds following rapid intravenous administration. Iopamidol did not bind to serum or plasma proteins at 1 hour after administration.

Elimination

The plasma half-life is approximately 2 hours after intravascular administration; the half-life is not dose dependent.

Metabolism

Iopamidol does not undergo significant metabolism, deiodination, or biotransformation.

Excretion

After intravascular administration, iopamidol is excreted primarily through the kidneys. In patients with normal renal function, the cumulative urinary excretion for iopamidol, expressed as a percentage of administered intravenous dose, is approximately 35% to 40% at 60 minutes, 80% to 90% at 8 hours, and 90% or greater in the 72- to 96-hour period after administration. In patients with normal renal function, approximately 1% or less of the administered dose appears in cumulative 72- to 96-hour fecal samples.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed with iopamidol to evaluate carcinogenic potential. No evidence of genetic toxicity was obtained in *in vitro* tests. In animal reproduction studies performed on rats, intravenously administered iopamidol did not induce adverse effects on fertility or general reproductive performance.

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

ISOVUE (iopamidol) injection is a clear, colorless to pale yellow solution. The ISOVUE Imaging Bulk Package is available in the following presentations:

Concentration (mg Iodine/mL)	Package Size	Package Type	Sale Unit	NDC
300	200 mL	Imaging Bulk Package	Carton of 10	0270-1315-45
	500 mL	Imaging Bulk Package	Carton of 6	0270-1315-95
370	200 mL	Imaging Bulk Package	Carton of 10	0270-1316-45
	500mL	Imaging Bulk Package	Carton of 6	0270-1316-95

Storage and Handling

Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature]. Protect from light.

17 PATIENT COUNSELING INFORMATION

Hypersensitivity Reactions

Advise the patient concerning the risk of hypersensitivity reactions that can occur both during and after ISOVUE administration. Advise the patient to report any signs or symptoms of hypersensitivity reactions during the procedure and to seek immediate medical attention for any signs or symptoms experienced after discharge [see *Warnings and Precautions* (5.2)].

Advise patients to inform their physician if they develop a rash after receiving ISOVUE [see *Warnings and Precautions* (5.11)].

Acute Kidney Injury

Advise the patient concerning appropriate hydration to decrease the risk of contrast induced kidney injury [see *Warnings and Precautions* (5.3)].

Extravasation

If extravasation occurs during injection, advise patients to seek medical care for progression of symptoms [see *Warnings and Precautions* (5.6)].

Thyroid Dysfunction

Advise parents/caregivers about the risk of developing thyroid dysfunction after ISOVUE administration. Advise parents/caregivers about when to seek medical care for their child to monitor for thyroid function [see *Warnings and Precautions* (5.8)].

Lactation

Advise a lactating woman that interruption of breastfeeding is not necessary, however, to minimize exposure to a breastfed infant, a lactating woman may consider pumping and discarding breast milk for 10 hours after ISOVUE administration [see *Use in Specific Populations* (8.2)].

Manufactured for:
Bracco Diagnostic Inc.
Princeton, NJ 08540

Manufactured by:
BIPSO GmbH
78224 Singen (Germany)

ISOVUE is a registered trademark of Bracco Diagnostics Inc.

200 mL

NDC 0270-1315-45

ISOVUE®
(iopamidol) injection

300 mg Iodine/mL

Imaging Bulk Package
For use only with an automated contrast injection system or contrast management system approved or cleared for use with this contrast agent in this Imaging Bulk Package

For intra-arterial or intravenous use only

ISOVUE 300 mg Iodine/mL



(01) 10302701315450

Manufactured for Bracco Diagnostics Inc., Princeton, NJ 08540
by BIPSO GmbH
78224 Singen (Germany)

Not For Intrathecal Use • Recommended Dosage: See Prescribing Information • See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use. • Each mL contains 612 mg iopamidol (providing 300 mg bound iodine), 0.48 mg edetate calcium disodium (providing 0.053 mg sodium), and 1 mg tromethamine; pH adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.
Provided the container is held under the labeled storage conditions the maximum use time is 10 hours from the initial entry.
Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature] • Protect from light
Rx only
Discard after ____/____/____ **at** ____:____

CBIC30X-US-F-173009778

ISOVUE® 300 mg Iodine/mL
LOT:

ISOVUE® 300 mg Iodine/mL
LOT:

LOT:
EXP:

125

150

175

200

APPROXIMATE

Reference ID: 5726847

500 mL

NDC 0270-1315-95

ISOVUE® (iopamidol) injection

300 mg Iodine/mL

Imaging Bulk Package

For use only with an automated contrast injection system or contrast management system approved or cleared for use with this contrast agent in this Imaging Bulk Package

**For intra-arterial or
intravenous use only**

ISOVUE® 300 mg Iodine/mL

Not For Intrathecal Use

Recommended Dosage: See Prescribing Information

See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use.

Each mL contains 612 mg iopamidol (providing 300 mg bound iodine), 0.48 mg edetate calcium disodium (providing 0.053 mg sodium), and 1 mg tromethamine; pH adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.

Provided the container is held under the labeled storage conditions the maximum use time is 10 hours from the initial entry.

Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature]
Protect from light

Rx only



(01) 10302701315955

Discard after ____/____/____ at ____:



Manufactured for Bracco Diagnostics Inc., Princeton, NJ 08540
by BIPSO GmbH
78224 Singen (Germany)

02701315955

ISOVUE® 300 mg Iodine/mL

LOT:

ISOVUE® 300 mg Iodine/mL

LOT:

LOT:

EXP.:

APPROXIMATE



10 Imaging Bulk Packages
200 mL each NDC 0270-1315-45

ISOVUE[®]

(iopamidol) injection

300 mg Iodine/mL

Imaging Bulk Package

For use only with an automated contrast injection system or contrast management system approved or cleared for use with this contrast agent in this Imaging Bulk Package

For intra-arterial or intravenous use only



(01)30302701315454(30)1

Manufactured for Bracco Diagnostics Inc.
Princeton, NJ 08540
by BIPSO GmbH, 78224 Singen (Germany)



CZ7380X - US F.1/3100054

Not For Intrathecal Use

Recommended Dosage: See Prescribing Information

See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use. Each mL contains 612 mg iopamidol (providing 300 mg bound iodine), 0.48 mg edetate calcium disodium (providing 0.053 mg sodium), and 1 mg tromethamine; pH adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.

Provided the container is held under the labeled storage conditions the maximum use time is 10 hours from the initial entry.

KEEP BOTTLES IN BOX WITH COVER CLOSED TO PROTECT FROM LIGHT

Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature]

Rx only

LOT:

EXP.:



6 Imaging Bulk Packages
500 mL each NDC 0270-1315-95

ISOVUE[®]

(iopamidol) injection

300 mg Iodine/mL

Imaging Bulk Package

For use only with an automated contrast injection system or contrast management system approved or cleared for use with this contrast agent in this Imaging Bulk Package

For intra-arterial or intravenous use only



(01)30302701315959(30)1

Manufactured for Bracco Diagnostics Inc.
Princeton, NJ 08540
by BIPSO GmbH, 78224 Singen (Germany)



CE7ZE0X - US F: 13100056

Not For Intrathecal Use

Recommended Dosage: See Prescribing Information

See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use. Each mL contains 612 mg iopamidol (providing 300 mg bound iodine), 0.48 mg edetate calcium disodium (providing 0.053 mg sodium), and 1 mg tromethamine; pH adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.

Provided the container is held under the labeled storage conditions the maximum use time is 10 hours from the initial entry.

KEEP BOTTLES IN BOX WITH COVER CLOSED TO PROTECT FROM LIGHT

Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature]

Rx only

LOT:

EXP.:



200 mL

NDC 0270-1316-45

ISOVUE® (iopamidol) injection

370 mg Iodine/mL

Imaging Bulk Package

For use only with an automated contrast injection system or contrast management system approved or cleared for use with this contrast agent in this Imaging Bulk Package

For intra-arterial or intravenous use only

ISOVUE® 370 mg Iodine/mL



Not For Intrathecal Use • Recommended Dosage: See Prescribing Information • See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use. • Each mL contains 755 mg iopamidol (providing 370 mg bound iodine), 0.48 mg edetate calcium disodium (providing 0.053 mg sodium), and 1 mg tromethamine; pH adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.

Provided the container is held under the labeled storage conditions the maximum use time is 10 hours from the initial entry.

Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature] • Protect from light

Rx only

Discard after __/__/__ at __: __

Manufactured for Bracco Diagnostics Inc., Princeton, NJ 08540
by BIPSO GmbH
78224 Singen (Germany)

C19C50X - US F 173009779

ISOVUE® 370 mg Iodine/mL

LOT:

ISOVUE® 370 mg Iodine/mL

LOT:

LOT:

EXP:

125

150

175

200

APPROXIMATE

500 mL

NDC 0270-1316-95

ISOVUE® (iopamidol) injection

370 mg Iodine/mL

Imaging Bulk Package

For use only with an automated contrast injection system or contrast management system approved or cleared for use with this contrast agent in this Imaging Bulk Package

**For intra-arterial or
intravenous use only**

ISOVUE® 370 mg Iodine/mL

Not For Intrathecal Use

Recommended Dosage: See Prescribing Information

See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use.

Each mL contains 755 mg iopamidol (providing 370 mg bound iodine), 0.48 mg edetate calcium disodium (providing 0.053 mg sodium), and 1 mg tromethamine; pH adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.

Provided the container is held under the labeled storage conditions the maximum use time is 10 hours from the initial entry.

Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature]

Protect from light

Rx only



Discard after ___/___/___ at ___:___



Manufactured for Bracco Diagnostics Inc., Princeton, NJ 08540
by BIPSO GmbH
78224 Singen (Germany)

09C70X - US F:12009781

ISOVUE® 370 mg Iodine/mL

LOT:

ISOVUE® 370 mg Iodine/mL

LOT:

LOT:

EXP.:

APPROXIMATE

200

250

300

350

400

450

500

10 Imaging Bulk Packages
200 mL each NDC 0270-1316-45

ISOVUE[®]

(iopamidol) injection

370 mg Iodine/mL

Imaging Bulk Package

For use only with an automated contrast injection system or contrast management system approved or cleared for use with this contrast agent in this Imaging Bulk Package

For intra-arterial or intravenous use only



(01)30302701316451(30)1

Manufactured for Bracco Diagnostics Inc.
Princeton, NJ 08540
by BIPSO GmbH, 78224 Singen (Germany)



CE7320X - US F. 1/3100055

Not For Intrathecal Use

See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use. Each mL contains 755 mg iopamidol (providing 370 mg bound iodine), 0.48 mg edetate calcium disodium (providing 0.053 mg sodium), and 1 mg tromethamine; pH adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.

Provided the container is held under the labeled storage conditions the maximum use time is 10 hours from the initial entry.

KEEP BOTTLES IN BOX WITH COVER CLOSED TO PROTECT FROM LIGHT

Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature]

Rx only

LOT:

EXP.:



6 Imaging Bulk Packages
500 mL each NDC 0270-1316-95

ISOVUE[®]

(iopamidol) injection

370 mg Iodine/mL

Imaging Bulk Package

For use only with an automated contrast injection system or contrast management system approved or cleared for use with this contrast agent in this Imaging Bulk Package

For intra-arterial or intravenous use only



(01)30302701316956(30)1

Manufactured for Bracco Diagnostics Inc.
Princeton, NJ 08540
by BIPSO GmbH, 78224 Singen (Germany)



CE7240X - US F.1/3100057

Not For Intrathecal Use

Recommended Dosage: See Prescribing Information

See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use. Each mL contains 755 mg iopamidol (providing 370 mg bound iodine), 0.48 mg edetate calcium disodium (providing 0.053 mg sodium), and 1 mg tromethamine; pH adjusted to 6.5 to 7.5 with hydrochloric acid and/or sodium hydroxide.

Provided the container is held under the labeled storage conditions the maximum use time is 10 hours from the initial entry.

KEEP BOTTLES IN BOX WITH COVER CLOSED TO PROTECT FROM LIGHT

Store at 20°C to 25°C (68°F to 77°F) [See USP controlled room temperature]

Rx only

LOT:

EXP.:



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

AUGUST A HOFLING
01/13/2026 08:59:42 AM