

NDC 0116-1021-04

DYNA-HEX 2[®]

Chlorhexidine Gluconate 2% Solution

Antiseptic

Non-sterile Solution

For Single-Use
Net Content: 4 fl oz (118 mL)

Drug Facts

Active ingredient	Purpose
Chlorhexidine gluconate 2%	Antiseptic

Uses

■ healthcare personnel handwash:
helps reduce bacteria that potentially can cause disease

■ patient preoperative skin preparation:
preparation of the patient's skin prior to surgery

■ skin wound and general skin cleansing

Drug Facts (continued)

Warnings

For external use only

Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

■ if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product

■ in contact with meninges

■ in the genital area

■ as a preoperative skin preparation of the head or face

When using this product

■ keep out of the eyes, ears and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums.

Drug Facts (continued)

■ if solution should contact these areas, rinse out promptly and thoroughly with water

■ wounds which involve more than the superficial layers of the skin should not be routinely treated

■ repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Healthcare personnel handwash:

■ wet hands with water

Drug Facts (continued)

■ dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds ■ rinse and dry thoroughly

Patient preoperative skin preparation:

■ apply product liberally to surgical site and swab for at least 2 minutes and dry with a sterile towel

■ repeat procedure for an additional 2 minutes and dry with a sterile towel

Skin wound and general skin cleaning:

■ thoroughly rinse the area to be cleaned with water

■ apply the minimum amount of product necessary to cover the skin or wound area and wash gently ■ rinse again thoroughly

Other information

■ store at 20-25°C (68-77°F)

■ avoid excessive heat above 40°C (104°F)

Inactive ingredients

citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

011610210406

2BLDY04BTLBLF

© 2016 Xiumin Laboratories, Inc. Mount Prospect, IL 60056

Reference ID: 5840774

Drug Facts (continued)**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with 5 mL of solution (4 full strokes) and a wet brush, paying close attention to the nails, cuticles and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of solution (4 full strokes) and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense 5 mL of solution (4 full strokes) into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

Questions or comments?

call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundry/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

2DYN32SQLBL

NDC 0116-1022-32

DYNA-HEX 2[®]

Chlorhexidine Gluconate 2% Solution

Antiseptic

Contains: 2% Chlorhexidine Gluconate

Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056



CHLORHEXIDINE GLUCONATE

Net Contents: 32 fl oz (1 qt) (946 mL)

Drug Facts**Active ingredient**

Chlorhexidine gluconate 2% solution Antiseptic

Purpose**Uses**

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care

Drug Facts (continued)

- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease

Warnings**For external use only****Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient are allergic to chlorhexidine gluconate or any other ingredient in this product

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.