







**Active ingredient (in each tablet)**

Cetirizine HCl 5 mg

**Purpose**

Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat.

**Warnings: Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose. **Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. **When using this product**

■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:**

■ if breast-feeding, not recommended ■ if pregnant, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control

Center right away. (1-800-222-1222) **Directions** **Adults and children**

**6 years and over:** 1 tablet (5 mg) or 2 tablets (10 mg) once daily

depending upon severity of symptoms; do not take more than 2 tablets

(10mg) in 24 hours. **Adults 15 years and over:** 1 tablet (5 mg) once daily;

do not take more than 1 tablet (5 mg) in 24 hours. **Children under 6 years of age:**

ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information**

■ store between 20° to 25°C (68° to 77°F) ■ do not use if foil inner seal imprinted with

"Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2. **Inactive**

**ingredients:** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose

monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol,

titanium dioxide. **Questions?** call 1-800-343-7065 (toll-free) or 215-273-8755 (collect)

The trade dress of this ZYRTEC® package is subject to trademark protection.

Active ingredient made in Switzerland. Distributed by: **JOHNSON & JOHNSON**

**CONSUMER INC.** McKel Consumer Healthcare Division Fort Washington, PA 19034 USA

Pat. www.jjcpats.com ©J&J 2003 **zyrtec.com**



To Close: Turn until Click

NDC 50580-754-35

**ZYRTEC**  
Cetirizine HCl tablets **ALLERGY**  
5 mg/antihistamine

**35**  
Tablets  
5 mg each

LOT: XXXXXXXX  
EXP: YYYYMMMM

**GO PACKS®**  
3 Portable Pouches  
One 10 mg tablet per pouch



Original Prescription Strength

**ZYRTEC®**  
NDC 50580-726-93  
Cetirizine HCl tablets **ALLERGY**  
10 mg antihistamine

Indoor &  
Outdoor  
Allergies  
**10** mg Strength  
Tablets

**24**  
Hour

relief of



Sneezing



Runny Nose



Itchy, Watery  
Eyes



Itchy Throat  
or Nose

**3** Tablets  
10 mg each



Actual Size

**3** TABLETS



89865005



30059868



3 00452042 3

Active ingredient made in Switzerland  
Distributed by:  
Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands 2025  
[www.zyrtec.com](http://www.zyrtec.com)  
Pat. [www.kenvuepats.com](http://www.kenvuepats.com)  
The trade dress of this ZYRTEC® package  
is subject to trademark protection.  
3005986888



LOT: XXXXXXX  
EXP: YYYY/MM

**Drug Facts**

**Active ingredient (in each tablet)** Cetirizine HCl 10 mg  
**Purpose** Antihistamine

**Uses** Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings**  
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.  
**Ask a doctor before use if you have liver or kidney disease.** Your doctor should determine if you need a different dose.  
**Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.**  
**When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens.  
**Stop use and ask a doctor if an allergic reaction to this product occurs.**  
Seek medical help right away:  
**If pregnant or breast-feeding:** ■ If breast-feeding: not recommended ■ If pregnant: ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**  
adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours.  
ask a doctor if symptoms are less severe.  
adults 65 years and over: ask a doctor.  
children under 6 years of age: ask a doctor.  
consumers with liver or kidney disease: ask a doctor.

**Other information** ■ store between 20° to 25°C (68° to 77°F) ■ do not use if pouch is torn or damaged ■ meets USP Dissolution Test 2

**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

**Questions?** call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)

**GO PACKS®**  
5 Portable Pouches One 10 mg tablet per pouch



**Original Prescription Strength**

**ZYRTEC®**  
Cetirizine HCl tablets **ALLERGY**  
10 mg /antihistamine

NDC 5565-725-36  
© Karmun Brands 2025  
www.zyrtec.com  
Pat. www.karmunpats.com

**24**  
Hour relief of

Sneezing  
Runny Nose

**5**  
Tablets  
10 mg each

Actual Size

**5**  
TABLETS

Itchy, Watery Eyes

Itchy Throat or Nose

Active Ingredient made in Switzerland  
Distributed by: Karmun Brands LLC, Summit, NJ 07901  
© Karmun Brands 2025  
www.zyrtec.com  
Pat. www.karmunpats.com  
NDC 5565-725-36  
3 12547 20430 9

**ZYRTEC®**  
ALLERGY

**Drug Facts**  
**(in each tablet)**  
Cetirizine HCl 10 mg  
Antihistamine

**Warnings**  
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have liver or kidney disease.** Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.**

**When using this product:**  
■ drowsiness may occur  
■ avoid alcoholic drinks  
■ alcohol, sedatives, and tranquilizers may increase drowsiness

■ Do not drive or operate machinery or use heavy machinery until you know how you react to this product.  
■ In rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens.

**Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away if you experience:  
■ hives  
■ difficulty breathing  
■ swelling of the face, lips, tongue, or throat

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**  
Adults and children 6 years and over:  
one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.

Children under 6 years of age:  
ask a doctor

Adults 65 years and over:  
ask a doctor

Children under 6 years of age:  
ask a doctor

Common side effects include:  
■ drowsiness

**Other information**  
■ Store between 20° to 25°C (68° to 77°F).  
■ Do not use if pouch is torn or damaged. ■ Needs USP Dissolution Test 2

**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide, triethylamine, xanthan gum

**Questions?** Call 1-800-343-1789 (toll-free) or 215-273-8733 (local).

**Uses**  
Importantly relieves these symptoms due to hay fever or other upper respiratory allergies:  
■ Runny nose ■ Sneezing ■ Itchy, watery eyes ■ Itching of the nose or throat

LOT: XXXXXX  
EXP: YYYY/MM/II





**Active ingredient in each tablet**  
Cetirizine HCl 10 mg

**Use:** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings:** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product:** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breastfeeding:** ■ if breastfeeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Purpose**  
Antihistamine

**Directions:** Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor. **Other information:** ■ store between 20° to 25° C (68° to 77° F) ■ do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions?** call 1-800-342-7665 (call-free) or 215-273-8756 (collect). The trade dress of the Zyrtec® package is subject to trademark protection. Active ingredient made in Switzerland. Distributed by: **KENVUE BRANDS LLC** Summit, NJ 07901 © Kenvue Brands LLC 2025 Pat. www.kenvuebrands.com

NDC 50580-726-71

**ZYRTEC®**

Cetirizine HCl ALLERGY  
antihistamine  
10 mg tablets

**30**  
TABLETS  
10 mg each

LOT: XXXXXX  
EXP: YYMM



**Active ingredient (in each tablet)**

Cetirizine HCl 10 mg  
**Use** ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat  
**Warnings** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breastfeeding: ■ if breastfeeding: not recommended ■ if pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical

**Purpose**

Antihistamine  
help or contact a Poison Control Center right away. (1-800-222-1222)  
**Directions** Adults and children 6 years and over: one 10 mg tablet once daily do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.  
Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor.  
Consumers with liver or kidney disease: ask a doctor. Other information ■ store between 20° to 25°C (68° to 77°F) ■ do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2.  
**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions** Call 1-800-343-7865 (toll-free) or 215-273-8755 (collect). The trade dress of this ZYRTEC® package is subject to trademark protection.  
Active ingredient made in Switzerland.  
Dist. by: KENVUE BRANDS LLC Summit, NJ 07801 © Kenvue Brands LLC 2025  
Pat. www.kenvuebrands.com



**40**  
TABLETS  
10 mg each

LOT: XXXXXX  
EXP: YYYYMM



Original Prescription Strength

Indoor & Outdoor Allergies

10 mg Strength  
4.5 mg Tablets

**ZYRTEC**  
Cetirizine HCl Tablets **ALLERGY**  
10 mg antihistamine

24 Hour relief of

- Sneezing
- Itchy Throat or Nose
- Itchy, Watery Eyes
- Runny Nose



4.5 TABLETS



Actual Size

LOT: XXXXXX  
EXP: YYYYMM

**Active ingredient (in each tablet)**

Cetirizine HCl 10 mg

**Use** ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat. **Warnings Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use** if you have liver or kidney disease. Your doctor should determine if you need a different dose. **Ask a doctor or pharmacist before use** if you are taking tranquilizers or sedatives. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. **Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breastfeeding** ■ if breastfeeding not recommended ■ if pregnant, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help

**Purpose**

Antihistamine

or contact a Poison Control Center right away. (1-800-222-1222)

**Directions Adults and children 6 years and over:** one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. **Adults 65 years and over:** ask a doctor. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information** ■ store between 20° to 25° C (68° to 77° F) ■ do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2. **Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions?** call 1-800-343-7865 (toll-free) or 215-273-0755 (collect). The trade dress of this Zyrtec® package is subject to trademark protection. **Active ingredient made in Switzerland.** Distributed by: Kenvue Brands LLC, Summit, NJ 07901 © Kenvue Brands LLC 2025 Pat. www.kenvuebrands.com



68865003

zyrtec.com

NDC 50580-726-45  
**ZYRTEC®**  
Cetirizine HCl ALLERGY  
antihistamine  
10 mg tablets

**45**  
TABLETS  
10 mg each

LOT: XXXXXX  
EXP: YYMM

**Active ingredient (in each tablet)**

Cetirizine HCl 10 mg

**Uses** ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat. **Warnings** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breastfeeding. ■ if breastfeeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a

**Purpose**

Antihistamine

Poison Control Center right away, (1-800-222-1222) **Directions** Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor. **Other information** ■ store between 20° to 25°C (68° to 77°F) ■ do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2. **Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions?** call 1-800-343-7895 (toll-free) or 215-273-8755 (collect). The trade dress of this Zyrtec® package is subject to trademark protection. Active made in Switzerland. Dist. by: KENVUE BRANDS LLC Summit, NJ 07801 © Kenvue Brands LLC 2025 Pat. www.kenvuepat.com



683960DC

NDC 50580-726-50  
**ZYRTEC®**  
Cetirizine HCl ALLERGY  
antihistamine  
10 mg tablets

**50**  
TABLETS  
10 mg each

LOT: XXXXXX  
EXP: YYYYMM

The trade dress of this ZYRTEC® package is subject to trademark protection.

10 mg antihistamine  
Cetirizine HCl Tablets  
**ZYRTEC**  
ALLERGY

Indoor & Outdoor Allergies

Original Prescription Strength

NDC 5050-730-73

**ZYRTEC**  
Cetirizine HCl Tablets ALLERGY  
10 mg/antihistamine

Indoor & Outdoor Allergies  
10 mg Strength Tablets

24 Hour relief of

Sneezing

Runny Nose

Itchy, Watery Eyes

Itchy Throat or Nose

60 Tablets 10 mg each

Actual Size

60 TABLETS

LOT: XXXXXX  
EXP: YYYY/MM

Active ingredient made in Switzerland.  
Distributed by KENVUE BRANDS LLC Summit, NJ 07901  
© Kenvue Brands LLC 2025  
Pat. www.kenvuepats.com  
zyrtec.com

30059891

3 00450 20666 2

**Drug Facts**

**Important:** Read all product information before using. Keep this box for important information.

**Directions**

**Warnings**

**Other information**

**Contains 60 Tablets**

**Actual Size**

**Active ingredient (in each tablet)**  
Cetirizine HCl 10 mg

**Use:** ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat.

**Warnings:** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you have liver or kidney disease.** Your doctor should determine if you need a different dose. **Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.**

**When using this product:** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. **Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breastfeeding:** ■ if breastfeeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or

**Purpose**  
Antihistamine

**Directions:** Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. **Adults 65 years and over:** ask a doctor. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information:** ■ store between 20° to 25°C (68° to 77°F) ■ do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2.

**Inactive ingredients:** colloid silicon dioxide, croscarmellose sodium, hydroxypropylcellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions?** call 1-800-343-7865 (toll-free) or 215-273-4755 (collect). The trade dress of this Zyrtec® package is subject to trademark protection. Active made in Switzerland.

Dist. by: **KENVUE BRANDS LLC** Summit, NJ 07901  
©Kenvue Brands LLC 2025 Pat. www.kenvuebrands.com **zyrtec.com**

891652015

NDC 50580-726-73

**ZYRTEC®**

Cetirizine HCl **ALLERGY**  
antihistamine  
10 mg tablets

**60**  
TABLETS  
10 mg each

LOT: XXXXXX  
EXP: YYYY/MM

**Active ingredient (in each tablet)**

Cetirizine HCl 10 mg

**Warnings:** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product, drowsiness may occur. Avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. In rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away if pregnant or breastfeeding. If breastfeeding, not recommended. If pregnant, ask a health professional before use.

**Purpose**

Antihistamine

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: runny nose, sneezing, itchy, watery eyes, itching of the nose or throat.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions:** Adults and children 6 years and over: one 10 mg tablet once daily, do not take more than one 10 mg tablet in 24 hours.

**Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information:** store between 20° to 25°C (68° to 77°F).

**Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.** Meets USP Dissolution Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.

**Questions?** call 1-800-343-7005 toll-free or 215-273-8755 (collect). The trade dress of this Zyrtec package is subject to trademark protection. Active made in Switzerland. Dist. by: KENVUE BRANDS LLC Summit, NJ 07901 © Kenvue Brands LLC 2025 Pat. www.kenvuepat.com zyrtec.com



30059040

NDC 50580-726-70  
**ZYRTEC**  
Cetirizine HCl ALLERGY  
antihistamine  
10 mg tablets

**70**  
TABLETS  
10 mg each

LOT: XXXXXX  
EXP: YYYY/MM



**Active ingredient (in each tablet)**

Cetirizine HCl 10 mg

**Use:** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: **runny nose** ■ **sneezing** ■ **itchy, watery eyes** ■ **itching of the nose or throat**  
**Warnings:** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product, drowsiness may occur. Avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. In rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away if pregnant or breastfeeding. If breastfeeding: not recommended. If pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison

**Purpose**

Antihistamine

Control Center right away. (1-800-222-1222) **Directions:** Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor. **Other information:** store between 20° to 25° C (68° to 77° F) ■ do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide. **Questions?** Call 1-800-342-7665 (toll-free) or 215-275-8755 (collect). The trade dress of the ZYRTEC® package is subject to trademark protection. Active ingredient made in Switzerland. Distributed by: Kenvue Brands LLC, Summit, NJ 07901 © Kenvue Brands LLC 2025 Pat. www.kenvuebrands.com



8886500C

zyrtec.com

NDC 50580-726-75  
**ZYRTEC®**  
Cetirizine HCl ALLERGY  
antihistamine  
10 mg tablets

**75**  
TABLETS  
10 mg each

LOT: XXXXXX  
EXP: YYMM

The trade dress of the ZYRTEC package is subject to trademark protection.

Indoor & Outdoor Allergies

10 mg Strength Tablets

Original Prescription Strength

**ZYRTEC**  
Cetirizine HCl tablets, ALLERGY  
10 mg/antihistamine

Indoor & Outdoor Allergies

10 mg Strength Tablets

24 Hour relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat of Nose

90 Tablets  
10 mg strength

Actual Size



90 TABLETS

Active ingredient made in Switzerland.  
Distributed by: KENVUE BRANDS LLC Summit, NJ 07901  
© Kenvue Brands LLC 2005  
Pat. www.kenvue.com  
zyrtec.com



30059502

**Keep this and product information before using. Keep this for important information.**

**Drug Facts (continued)**

**Warnings:** Tell your doctor if you are taking any other medicines, especially other antihistamines, sedatives, or tranquilizers. Tell your doctor if you are pregnant, planning to get pregnant, or breastfeeding. Tell your doctor if you have any other medical conditions, especially liver or kidney disease, or if you have ever had an allergic reaction to any medicine. Tell your doctor if you are taking any other medicines, especially other antihistamines, sedatives, or tranquilizers. Tell your doctor if you are pregnant, planning to get pregnant, or breastfeeding. Tell your doctor if you have any other medical conditions, especially liver or kidney disease, or if you have ever had an allergic reaction to any medicine.

**Directions:** See important information about this medicine on page 2 of the patient information. This medicine is to be used as directed. Do not take more than the recommended dose. Do not take this medicine if you are allergic to cetirizine or any of the ingredients. Do not take this medicine if you are taking any other medicines, especially other antihistamines, sedatives, or tranquilizers. Do not take this medicine if you are pregnant, planning to get pregnant, or breastfeeding. Do not take this medicine if you have any other medical conditions, especially liver or kidney disease, or if you have ever had an allergic reaction to any medicine.

**Other information:** This medicine may cause drowsiness. Do not drive or operate machinery until you know how this medicine affects you. Do not drink alcohol while taking this medicine. Do not take this medicine if you are taking any other medicines, especially other antihistamines, sedatives, or tranquilizers. Do not take this medicine if you are pregnant, planning to get pregnant, or breastfeeding. Do not take this medicine if you have any other medical conditions, especially liver or kidney disease, or if you have ever had an allergic reaction to any medicine.

LOT: XXXXXXX  
EXP: YYMMMM



Actual Size

**Active ingredient (in each tablet)**

Cetirizine HCl 10 mg

**Uses:** ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings:** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away, if pregnant or breastfeeding: ■ if breastfeeding: not recommended ■ if pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical

**Purpose**

Antihistamine

help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions:** Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.

**Adults 65 years and over:** ask a doctor. **Children under 6 years of age:** ask a doctor.

**Consumers with liver or kidney disease:** ask a doctor. **Other information:** ■ store between 20° to 25°C (68° to 77°F) ■ do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2. **Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.

**Questions?** call 1-800-343-7895 (toll-free) or 215-273-6755 (collect). The trade dress of the Zyrtec® package is subject to trademark protection. Active made in Switzerland. Dist. by KENVUE BRANDS LLC Summit, NJ 07801 © Kanvue Brands LLC 2025 Pat. www.kanvuetpets.com



3J0894Z

zyrtec.com

NDC 50580-726-74  
**ZYRTEC®**  
Cetirizine HCl ALLERGY  
antihistamine  
10 mg tablets

90  
TABLETS  
10 mg each

LOT: XXXXXXXX  
EXP: YYYY/MMMM

**Keep out of reach of children.** See important information about this product on page 2 of this package insert. See the back of this package for more information.

**Warnings:** Do not take if you are allergic to cetirizine or any of the ingredients. Tell your doctor if you are taking any other medicines, especially other antihistamines, sedatives, or alcohol. Tell your doctor if you are pregnant, planning to get pregnant, or breastfeeding. Tell your doctor if you have any other medical conditions, especially liver or kidney disease, or if you are taking any other medicines. Tell your doctor if you are taking any other medicines, especially other antihistamines, sedatives, or alcohol. Tell your doctor if you are pregnant, planning to get pregnant, or breastfeeding. Tell your doctor if you have any other medical conditions, especially liver or kidney disease, or if you are taking any other medicines.

**Directions:** Adults and children 12 years of age and older: Take 10 mg (1 tablet) once daily. Children 6 to 11 years of age: Take 5 mg (1/2 tablet) once daily. Children 2 to 5 years of age: Take 2.5 mg (1/4 tablet) once daily. Children 6 months to 1 year of age: Take 1.25 mg (1/8 tablet) once daily. Do not take more than 10 mg (1 tablet) in 24 hours. Do not take more than 10 mg (1 tablet) in 24 hours.

**Side Effects:** The most common side effects of Zyrtec are drowsiness, dry mouth, sore throat, and fatigue. Other side effects include headache, dizziness, and constipation. Tell your doctor if you experience any of these side effects. Tell your doctor if you experience any of these side effects.

**Drug Interactions:** Zyrtec may interact with other medicines, especially other antihistamines, sedatives, or alcohol. Tell your doctor if you are taking any other medicines, especially other antihistamines, sedatives, or alcohol. Tell your doctor if you are pregnant, planning to get pregnant, or breastfeeding. Tell your doctor if you have any other medical conditions, especially liver or kidney disease, or if you are taking any other medicines.

**Use in Children:** Zyrtec is safe and effective for the treatment of allergic rhinitis and allergic conjunctivitis in children 6 months of age and older. Tell your doctor if you are taking any other medicines, especially other antihistamines, sedatives, or alcohol. Tell your doctor if you are pregnant, planning to get pregnant, or breastfeeding. Tell your doctor if you have any other medical conditions, especially liver or kidney disease, or if you are taking any other medicines.

**How to Use:** Zyrtec is taken orally with or without food. Do not crush or chew the tablets. Swallow the tablets whole. Do not take more than 10 mg (1 tablet) in 24 hours. Do not take more than 10 mg (1 tablet) in 24 hours.

**Storage:** Store Zyrtec at room temperature (20° to 25°C). Excursions permitted to 15° to 30°C. Do not store in the bathroom. Do not store in the bathroom. Do not store in the bathroom.

**Keep out of reach of children.** See important information about this product on page 2 of this package insert. See the back of this package for more information.

**ZYRTEC**  
Cetirizine HCl Tablets **ALLERGY**  
Indoor & Outdoor  
10 mg

Original Prescription Strength

**ZYRTEC**  
Cetirizine HCl tablets **ALLERGY**  
10 mg antihistamine

Indoor & Outdoor Allergies  
10 mg Strength Tablets

24 Hour relief of

Actual Size

Sneezing Runny Nose Itchy, Watery Eyes Itchy Throat or Nose



120 Total Tablets

70ct + 50ct = 120ct TABLETS

Active ingredient made in Switzerland.  
Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025  
Pat. www.kenvuebrands.com  
zyrtec.com



30059338

LOT: XXXXXXXX  
EXP: YYMM



The trade dress of this Zyrtec® package is subject to trademark protection.



Original Prescription Strength

Original Prescription Strength

Drug Facts									
<b>Active ingredient (in each tablet)</b> Cetirizine HCl 10 mg									
<b>Purpose</b> Antihistamine									
<b>Uses</b> Temporarily relieves these symptoms due to hay fever or other upper respiratory allergens: <ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ sneezing</li> <li>■ itchy, watery eyes</li> <li>■ itching of the nose or throat</li> </ul>									
<b>Warnings</b> Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.									
<b>When using this product</b> <ul style="list-style-type: none"> <li>■ drowsiness may occur</li> <li>■ avoid alcoholic drinks</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> <li>■ in rare cases, you may experience itching after stopping cetirizine.</li> </ul>									
<b>Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.</b> <ul style="list-style-type: none"> <li>■ if pregnant or breast-feeding, not recommended</li> <li>■ if pregnant, ask a health professional before use.</li> </ul>									
<b>Directions</b> <table border="1"> <tr> <td>adults and children 6 years and over</td> <td>one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours.</td> </tr> <tr> <td>adults 65 years and over</td> <td>A 5 mg product may be appropriate for less severe symptoms.</td> </tr> <tr> <td>children under 6 years of age</td> <td>ask a doctor</td> </tr> <tr> <td>consumers with liver or kidney disease</td> <td>ask a doctor</td> </tr> </table>		adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours.	adults 65 years and over	A 5 mg product may be appropriate for less severe symptoms.	children under 6 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor
adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours.								
adults 65 years and over	A 5 mg product may be appropriate for less severe symptoms.								
children under 6 years of age	ask a doctor								
consumers with liver or kidney disease	ask a doctor								
<b>Other information</b> <ul style="list-style-type: none"> <li>■ store between 20° to 25°C (68° to 77°F)</li> <li>■ do not use if pouch is torn or damaged</li> <li>■ meets USP Dissolution Test 2</li> </ul>									
<b>Inactive ingredients</b> colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide									
<b>Questions?</b> call 1-800-343-7808 (ask three) or 215-273-8750 (collect)									

Active ingredient made in Switzerland.  
Distributed by: Keniva Brands LLC, Summit, NJ 07901  
© Keniva Brands LLC 2025 www.zyrtec.com  
Pat. www.kenivabrands.com  
The trade dress of this Zyrtec® package is subject to trademark protection.

3 00450 20450 9

LOT: XXXXXX  
EXP: YYYYMM

50

Pouches of 1 Tablet Per Pouch

50

Pouches of 1 Tablet Per Pouch

50

Pouches of 1 Tablet Per Pouch

NDC 50580-726-05

Zyrtec®

Cetirizine HCl tablets ALLERGY  
10 mg/antihistamine

Indoor & Outdoor Allergies

10 mg Tablets

24

Hour relief of

Sneezing

Runny Nose

50

Pouches 10 mg each

Actual Size

Itchy, Watery Eyes

Itchy Throat or Nose



**Drug Facts**

Active ingredient (in each tablet)	Purpose
Cetirizine HCl 10 mg	Antihistamine

**Uses:** temporarily relieve these symptoms due to hay fever or other upper respiratory allergen:  
 ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings**  
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.  
 Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.  
 Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product:** ■ drowsiness may occur ■ avoid alcoholic drinks  
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness  
 ■ be careful when driving a motor vehicle or operating machinery  
 ■ in rare cases, you may experience itching after stopping cetirizine.  
 Consult your healthcare provider if this happens.

**Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.**  
 If pregnant or breast-feeding: ■ if breast-feeding, not recommended ■ if pregnant, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-252-1522).

**Directions**

adults and children 6 years and over	one 10 mg tablet once daily. Do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over	ask a doctor
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information** ■ store between 20° to 25°C (68° to 77°F) ■ do not use if pouch is torn or damaged  
 ■ meets USP Dissolution Test 2

**Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, hydroxypropylcellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

**Questions?** call 1-800-343-7805 (toll-free) or 215-273-8755 (outside)

Active ingredient made in Switzerland.  
 Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
 © Kenvue Brands LLC 2025 www.zyrtec.com  
 Pat. www.kenvue.com  
 The trade dress of this ZYRTEC® package is subject to trademark protection.

3 00450 20450 9

LOT: XXXXXX  
 BP: YYYYMM

Original Prescription Strength

**ZYRTEC**  
Cetirizine HCl tablets **ALLERGY**  
10 mg/antihistamine

Indoor & Outdoor Allergies  
10 mg Strength

FREE SAMPLES  
Not For Retail Sale

**50** Pouches of 1 Tablet Per Pouch

Original Prescription Strength

NDC 50580-726-92

**ZYRTEC**  
Cetirizine HCl tablets **ALLERGY**  
10 mg/antihistamine

Indoor & Outdoor Allergies  
10 mg Strength

**24** Hour relief of  
 Sneezing  
 Runny Nose  
 Itchy, Watery Eyes  
 Itchy Throat or Nose

**50** Pouches of 10 mg each  
 Actual Size

**50** Pouches of 1 Tablet Per Pouch

Original Prescription Strength

**ZYRTEC**  
Cetirizine HCl tablets **ALLERGY**  
10 mg/antihistamine

Indoor & Outdoor Allergies  
10 mg Strength

**50** Pouches of 1 Tablet Per Pouch







This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine**. If your doctor has prescribed an epinephrine injector for "bug bites" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times. **Do not use to prevent hives** from any known cause such as: **■** insect stings **■** medications **■** latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. There can sometimes be a return. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. **■** If you have ever had an allergic reaction to this product or any of its ingredients or to a medicine containing hydroxyzine. **Ask a doctor before use if you have** **■** liver or kidney disease. Your doctor should determine if you need a different dose. **■** Hives that are an unusual color, look broken or blistered **■** Hives that do not stop. **Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product if drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■** In rare cases, you may experience itching after stopping treatment. Consult your healthcare provider if this happens. **Stop use and ask a doctor if ■** an allergic reaction to this product occurs. Seek medical help right away. **■** symptoms do not improve after 3 days of treatment **■** the hives have lasted more than 6 weeks **■** pregnant or breast-feeding **■** if breast-feeding not recommended **■** if pregnant, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-322-1222).

**Active ingredient (in each tablet)** **■** **Hydroxyzine** (HCl) 10 mg  
**Purpose**  
 Zyrtec Hives 10 mg tablets reduce hives and relieve itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.  
**Warnings: Severe Allergy Warning:** Get emergency help **immediately** if you have hives along with any of the following symptoms:  
 ■ trouble breathing ■ dizziness or loss of consciousness ■ swelling of tongue ■ swelling in or around mouth ■ trouble speaking ■ wheezing ■ wheezing or problems breathing. These symptoms may be signs of anaphylactic shock.  
**PEEL HERE**



LOT: XXXXXX  
 EXP: YYYYMM

**Directions:** Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. **Adults 18 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor.**  
**Other information:** ■ Store between 20° to 25° (68° to 77°) ■ Do not use if full liner seal imprinted with "Sealed For Your Safety" is broken or missing ■ meets USP Dissolution Test 2  
**Inactive ingredients:** croscarmellose sodium, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide  
**Questions?** call 1-800-343-7000 (ask free) or 215-273-8755 (collect).

The trade dress of this ZYRTEC® package is subject to trademark protection.  
 Active ingredient made in Switzerland.  
 Distributed by: **KENVUE BRANDS LLC**, Summit, NJ 07901  
 © Kenvue Brands LLC 2023. [www.kenvuebrands.com](http://www.kenvuebrands.com)

[zyrtec.com](http://zyrtec.com)

SPR6900C