

Vivelle®
estradiol transdermal system

Continuous delivery for twice-weekly application

Rx only

Prescribing Information

WARNING: ENDOMETRIAL CANCER WITH UNOPPOSED ESTROGEN IN WOMEN WITH A UTERUS

- There is an increased risk of endometrial cancer in a woman with a uterus who uses unopposed estrogens. Adding a progestogen to estrogen-only therapy has been shown to reduce the risk of endometrial hyperplasia, which may be a precursor to endometrial cancer.
- Perform adequate diagnostic measures, including directed or random endometrial sampling when indicated, to rule out malignancy in menopausal women abnormal genital bleeding of unknown etiology (see WARNINGS)


DESCRIPTION

The Vivelle® (estradiol transdermal system) contains estradiol in a multipolymeric adhesive. The system is designed to release estradiol continuously upon application to intact skin.

Five systems are available to provide nominal *in vivo* delivery of 0.025, 0.0375, 0.05, 0.075, or 0.1 mg of estradiol per day via skin of average permeability. Each corresponding system having an active surface area of 7.25, 11.0, 14.5, 22.0, or 29.0 cm² contains 2.17, 3.28, 4.33, 6.57, or 8.66 mg of estradiol USP, respectively. The composition of the systems per unit area is identical.


Estradiol USP is a white, crystalline powder, chemically described as estra-1,3,5(10)-triene-3,17β-diol.

The structural formula is



The molecular formula of estradiol is C₁₈H₂₄O₂. The molecular weight is 272.39.

The Vivelle system comprises three layers. Proceeding from the visible surface toward the surface attached to the skin, these layers are (1) a translucent flexible film consisting of an ethylene vinyl alcohol copolymer film, a polyurethane film, urethane polymer and epoxy resin, (2) an adhesive formulation containing estradiol USP, acrylic adhesive, polyisobutylene, ethylene vinyl acetate copolymer, 1,3 butylene glycol, styrene-butadiene rubber, oleic acid NF, lecithin, propylene glycol, bentonite NF, mineral oil USP, and dipropylene glycol, and (3) a polyester release liner that is attached to the adhesive surface and must be removed before the system can be used.



The active component of the system is estradiol. The remaining components of the system are pharmacologically inactive.

CLINICAL PHARMACOLOGY

Endogenous estrogens are largely responsible for the development and maintenance of the female reproductive system and secondary sexual characteristics. Although circulating estrogens exist in a dynamic equilibrium of metabolic interconversions, estradiol is the principal intracellular human estrogen and is substantially more potent than its metabolites, estrone and estriol, at the receptor level.

The primary source of estrogen in normally cycling adult women is the ovarian follicle, which secretes 70 to 500 mcg of estradiol daily, depending on the phase of the menstrual cycle. After menopause, most endogenous estrogen is produced by conversion of androstenedione, secreted by the adrenal cortex, to estrone by peripheral tissues. Thus, estrone and the sulfate conjugated form, estrone sulfate, are the most abundant circulating estrogens in postmenopausal women.

Estrogens act through binding to nuclear receptors in estrogen-responsive tissues. To date, two estrogen receptors have been identified. These vary in proportion from tissue to tissue.

Circulating estrogens modulate the pituitary secretion of the gonadotropins, luteinizing hormone (LH) and follicle stimulating hormone (FSH) through a negative feedback mechanism. Estrogens act to reduce the elevated levels of these hormones seen in postmenopausal women.

Pharmacokinetics


Absorption

In a multiple-dose study consisting of three consecutive patch applications of the Vivelle system, which was conducted in 17 healthy, postmenopausal women, blood levels of estradiol and estrone were compared following application of these units to sites on the abdomen and buttocks in a crossover fashion. Patches that deliver nominal estradiol doses of approximately

0.0375 mg/day and 0.1 mg/day were applied to abdominal application sites while the 0.1 mg/day doses were also applied to sites on the buttocks. These systems increased estradiol levels above baseline within four hours and maintained respective mean levels of 25 and 79 pg/mL above baseline following application to the abdomen; slightly higher mean levels of 88 pg/mL above baseline were observed following application to the buttocks. At the same time, increases in estrone plasma concentrations averaged about 12 and 50 pg/mL, respectively, following application to the abdomen and 61 pg/mL for the buttocks. While plasma concentrations of estradiol and estrone remained slightly above baseline at 12 hours following removal of the patches in this study, results from another study show these levels to return to baseline values within 24 hours following removal of the patches.

The figure (see Figure 1) illustrates the mean plasma concentrations of estradiol at steady state during application of these patches at four different dosages.

Figure 1
Steady-State Estradiol Plasma Concentrations
for Systems Applied to the Abdomen
Nonbaseline-Corrected Levels



The corresponding pharmacokinetic parameters are summarized in Table 1 below.

Table 1
Steady-State Estradiol Pharmacokinetic Parameters
for Systems Applied to the Abdomen
(mean $\pm$ standard deviation)
Nonbaseline-Corrected Data*

Dosage (mg/day)	C _{max} [†] (pg/mL)	C _{avg} [†] (pg/mL)	C _{min} (84 hr) [§] (pg/mL)
0.0375	46 $\pm$ 16	34 $\pm$ 10	30 $\pm$ 10
0.05	83 $\pm$ 41	57 $\pm$ 23 [#]	41 $\pm$ 11 [#]
0.075	99 $\pm$ 35	72 $\pm$ 24	60 $\pm$ 24
0.1	133 $\pm$ 51	89 $\pm$ 38	90 $\pm$ 44
0.1	145 $\pm$ 71	104 $\pm$ 52	85 $\pm$ 47

*Mean baseline estradiol concentration = 11.7 pg/mL

[†]Peak plasma concentration

‡Average plasma concentration

§Minimum plasma concentration at 84 hr

#Measured over 80 hr

¶Applied to the buttocks

Distribution

The distribution of exogenous estrogens is similar to that of endogenous estrogens. Estrogens are widely distributed in the body and are generally found in higher concentrations in the sex hormone target organs. Estrogens circulate in the blood largely bound to sex hormone binding globulin (SHBG) and albumin.

Metabolism

Exogenous estrogens are metabolized in the same manner as endogenous estrogens. Circulating estrogens exist in a dynamic equilibrium of metabolic interconversions. These transformations take place mainly in the liver. Estradiol is converted reversibly to estrone, and both can be converted to estriol, which is the major urinary metabolite. Estrogens also undergo enterohepatic recirculation via sulfate and glucuronide conjugation in the liver, biliary secretion of conjugates into the intestine, and hydrolysis in the gut followed by reabsorption. In postmenopausal women, a significant proportion of the circulating estrogens exist as sulfate conjugates, especially estrone sulfate, which serves as a circulating reservoir for the formation of more active estrogens.

Excretion

Estradiol, estrone, and estriol are excreted in the urine along with glucuronide and sulfate conjugates. Studies conducted with the Vivelle system show the drug has an apparent mean half-life of 4.4 ± 2.3 hours. After removal of the transdermal systems, serum concentrations of estradiol and estrone returned to baseline levels within 24 hours.

Special Populations

Vivelle was only investigated in postmenopausal women.

Drug Interactions

In vitro and in vivo studies have shown that estrogens are metabolized partially by cytochrome P450 3A4 (CYP3A4). Therefore, inducers or inhibitors of CYP3A4 may affect estrogen drug metabolism. Inducers of CYP3A4 such as St. John's Wort preparations (*Hypericum perforatum*), phenobarbital, carbamazepine and rifampin may reduce plasma concentrations of estrogens, possibly resulting in a decrease in therapeutic effects and/or changes in the uterine bleeding profile. Inhibitors of CYP3A4 such as erythromycin, clarithromycin, ketoconazole, itraconazole, ritonavir and grapefruit juice may increase plasma concentrations of estrogens and may result in side effects.


Adhesion

Data showing the number of systems in controlled studies that required replacement due to inadequate adhesion is not available.

CLINICAL STUDIES

In two controlled clinical trials of 356 subjects, the 0.075 and 0.1 mg doses were superior to placebo in relieving vasomotor symptoms at Week 4 and maintained efficacy through Weeks 8 and 12 of treatment. In this original study, the 0.0375 and 0.05 mg doses, however, did not differ from placebo until approximately Week 6; therefore, an additional 12-week placebo-controlled study in 255 patients was performed to establish the efficacy of the lowest dose of 0.0375 mg. The baseline mean daily number of hot flushes in these 255 patients was 11.5. Results at Weeks 4, 8, and 12 of treatment are shown in the figure below (see Figure 2).

Figure 2
Mean (SD) Change from Baseline in Mean Daily Number of Flushes
for Vivelle 0.0375 mg Versus Placebo in a 12-Week Trial




The 0.0375-mg dose was superior to placebo in reducing both the frequency and severity of vasomotor symptoms at Week 4 and maintained efficacy through Weeks 8 and 12 of treatment. The following doses of Vivelle: 0.0375 mg, 0.05 mg, 0.075 mg, and 0.1 mg, are effective for the control of vasomotor symptoms.

Efficacy and safety of the Vivelle system in the prevention of postmenopausal osteoporosis have been studied in a 2-year double-blind, randomized, placebo-controlled, parallel group study. A total of 261 hysterectomized (161) and non-hysterectomized (100), surgically or naturally menopausal women (within five years of menopause), with no evidence of osteoporosis (lumbar spine bone mineral density within two standard deviations of average peak bone mass, i.e., $=0.827$ g/cm²) were enrolled in this study; 194 patients were randomized to one of the four doses of Vivelle (0.1, 0.05, 0.0375, or 0.025 mg/day) and 67 patients to placebo. Over two years, study systems were applied to the buttock or the abdomen twice a week. Non-hysterectomized women received oral medroxyprogesterone acetate (2.5 mg/day) throughout the study.

The study population comprised naturally (82%) or surgically (18%) menopausal, hysterectomized (61%) or non-hysterectomized (39%) women with a mean age of 52.0 years (range 27 to 62 years); the mean duration of menopause was 31.7 months (range 2 to 72 months). Two hundred thirty-two (89%) of randomized subjects (173 on active drug, 59 on placebo) contributed data to the analysis of percent change from baseline in bone mineral density (BMD) of the AP lumbar spine, the primary efficacy variable. Patients were given supplemental dietary calcium (1000 mg elemental calcium/day) but no supplemental vitamin D. There was an increase in BMD of the AP lumbar spine in all Vivelle dose groups; in


contrast to this, a decrease in AP lumbar spine BMD was observed in placebo patients. All Vivelle doses were significantly superior to placebo ($p < 0.05$) at all time points with the exception of Vivelle 0.05 mg/day at six months. The highest dose of Vivelle was superior to the three lower doses. There were no statistically significant differences in pairwise comparisons among the three lower doses (see Figure 3).

Figure 3
Bone Mineral Density - AP Lumbar Spine
Least Squares Means of Percentage Change from Baseline
All Randomized Patients with at Least One Post-Baseline Assessment Available with
Last Post-Baseline Observation Carried Forward



Analysis of percent change from baseline in femoral neck BMD, a secondary efficacy outcome variable, showed qualitatively similar results; all doses of Vivelle were significantly superior to placebo ($p < 0.05$) at 24 months. The highest Vivelle dose was superior to placebo at all time points. A mixture of significant and non-significant results were obtained for the lower dose groups at earlier time points. The highest Vivelle dose was superior to the three lower doses, and there were no significant differences among the three lower doses at this skeletal site (see Figure 4).

Figure 4
Bone Mineral Density - Femoral Neck
Least Squares Means of Percentage Change from Baseline
All Randomized Patients with at Least One Post-Baseline Assessment Available with
Last Post-Baseline Observation Carried Forward



The mean serum osteocalcin (a marker of bone formation) and urinary excretion of cross-link N-telopeptides of type 1 collagen (a marker of bone resorption) decreased numerically in most of the active treatment groups relative to baseline. However, the

decreases in both markers were inconsistent across treatment groups and the differences between active treatment groups and placebo were not statistically significant.

Women's Health Initiative Estrogen-Alone Trial

The WHI estrogen-alone trial enrolled predominantly healthy postmenopausal women in trial to assess the risks and benefits of daily oral CE (0.625 mg)-alone compared to placebo in the prevention of certain chronic diseases. The primary endpoint was the incidence of CHD (defined as nonfatal MI, silent MI and CHD death), with invasive breast cancer as the primary adverse outcome. A “global index” included the earliest occurrence of CHD, invasive breast cancer, stroke, PE, colorectal cancer, hip fracture, or death due to other cause. This trial did not evaluate the effects of CE-alone on menopausal symptoms.

The WHI estrogen-alone trial was stopped early because an increased risk of stroke was observed, and it was deemed that no further information would be obtained regarding the risks and benefits of estrogen-alone in predetermined primary endpoints.

Centrally adjudicated results for stroke events, after an average follow-up of 7.1 years, reported estrogen-alone increased the risk for ischemic stroke compared to placebo, and this excess risk was present in all subgroups of women examined.

No overall difference for primary CHD events (nonfatal MI, silent MI and CHD death) and invasive breast cancer incidence in women receiving CE-alone compared to placebo was reported in final centrally adjudicated results from the estrogen-alone trial, after an average follow-up of 7.1 years.^{2,3}

Results of the estrogen-alone trial, which included 10,739 women (average age of 63 years, range 50 to 79 years; 75.3% White, 15.1% Black, 6.1% Hispanic, 3.6% Other), after an average follow-up of 7.1 years are presented in Table A.

Table A: Relative and Risk Difference Observed in the WHI Estrogen-Alone Trial at an Average of 7.1 Years of Follow-up^a

Event	Relative Ratio (95% CI) ^c	Risk Difference (CE vs placebo/10,000 WYs)
CHD events	0.94 (0.78-1.14)	-3 (55 vs 58)
Non-fatal MI	0.97 (0.79-1.21)	-1 (44 vs 45)
CHD death	1.00 (0.77-1.31)	0 (29 vs 29)
All Strokes	1.35 (1.07-1.70)	11 (45 vs 34)
Deep vein thrombosis ^d	1.48 (1.06-2.07)	7 (23 vs 15)
Pulmonary embolism	1.35 (0.89-2.05)	4 (14 vs 10)
Invasive breast cancer	0.79 (0.61-1.02)	-7 (28 vs 35)
Colorectal cancer	1.15 (0.81-1.64)	2 (17 vs 15)
Hip fracture	0.67 (0.46-0.96)	-6 (13 vs 19)
Vertebral fractures ^d	0.64 (0.44-0.93)	-6 (12 vs 18)
Total fractures ^d	0.72 (0.64-0.80)	-61 (153 vs 214)
Overall mortality ^{c,f}	1.03 (0.88-1.21)	3 (80 vs 77)
Global Index ^g	1.03 (0.93-1.13)	4 (208 vs 204)

^a Adapted from 2013 WHI trial (CE n=5,310, placebo n=5,429). WHI publications can be viewed at www.nhlbi.nih.gov/whi.

^b Results are based on centrally adjudicated data.

^c In the WHI studies, hazard ratios were estimated using Cox proportional hazards models comparing treatment to placebo; however, they are described here as relative risks. Nominal confidence intervals unadjusted for multiple looks and multiple comparisons.

^d Not included in “global index.”

^e Includes metastatic and non-metastatic breast cancer with the exception of in situ cancer.

^f All deaths, except from breast or colorectal cancer, definite or probable CHD, PE or cerebrovascular disease.

^g A subset of the events was combined in a “global index,” defined as the earliest occurrence of CHD events, invasive breast cancer, stroke, PE, colorectal cancer, hip fracture, or death due to other causes.

Timing of the initiation of estrogen-alone therapy relative to the start of menopause may affect the overall risk benefit profile. The study results for women 50-59 years old in the WHI estrogen-alone trial are shown in Table B.

Table B. Relative Risk and Risk Difference Observed Among Women 50-59 Years of Age in the WHI Estrogen – Alone Trial at an Average of 7.1 Years^{a,b}

Event	Relative Ratio (95% CI) ^c	Risk Difference (CE vs placebo/10,000 WYs)
CHD events	0.60 (0.35-1.04)	-11 (17 vs 28)
Non-fatal MI	0.55 (0.31-1.00)	-11 (14 vs 25)
CHD death	0.80 (0.32-2.04)	-1 (7 vs 8)
All Strokes	0.99 (0.53-1.85)	-1 (16 vs 17)
Deep vein thrombosis ^d	1.66 (0.75-3.67)	5 (13 vs 8)
Pulmonary embolism	1.53 (0.63-3.75)	4 (10 vs 6)
Invasive breast cancer ^e	0.82 (0.50-1.34)	-5 (24 vs 29)
Colorectal cancer	0.71 (0.30-1.67)	-3 (7 vs 10)
Hip fracture	5.01 (0.59-42.91)	3 (1 vs 3)
Vertebral fractures ^d	0.50 (0.17-1.47)	-4 (4 vs 8)
Total fractures ^d	0.90 (0.72-1.11)	-16 (133 vs 149)
Overall mortality ^{c,f}	0.70 (0.46-1.09)	-11 (29 vs 40)
Global Index ^g	0.84 (0.66-1.07)	-19 (98 vs 117)

^a Adapted from 2013 WHI trial (CE n=1,639; placebo=1,674). WHI publications can be viewed at www.nhlbi.nih.gov/whi.

^b Results are based on centrally adjudicated data.

^c In the WHI studies, hazard ratios were estimated using Cox proportional hazards models comparing treatment to placebo; however, they are described here as relative risks. Nominal confidence intervals unadjusted for multiple looks and multiple comparisons.

^d Not included in “global index.”

^e Includes metastatic and non-metastatic breast cancer with the exception of in situ cancer.

^f All deaths, except from breast or colorectal cancer, definite or probable CHD, PE or cerebrovascular disease.

^g A subset of the events was combined in a “global index,” defined as the earliest occurrence of CHD events, invasive breast cancer, stroke, PE, colorectal cancer, hip fracture, or death due to other causes.

Women’s Health Initiative Estrogen-Alone Memory Study

The WHIMS estrogen-alone ancillary study of WHI enrolled 2,947 predominantly healthy hysterectomized postmenopausal women 65 to 79 years of age (45% were 65 to 69 years of age, 36% were 70 to 74 years of age, and 19% were 75 years of age and older) to evaluate the effects of daily CE (0.625 mg)-alone on the incidence of probable dementia (primary outcome) compared to placebo. Probable dementia as defined in this study included Alzheimer's disease (AD), vascular dementia (VaD) and mixed type (having features of both AD and VaD). The most common classification of probable dementia in the treatment group and the placebo group was AD.

After an average follow-up of 5.2 years, the relative risk of probable dementia for CEalone versus placebo was 1.49 (95% CI, 0.83-2.66). The absolute risk of probable dementia for CE-alone versus placebo was 37 versus 25 cases per 10,000 women years. Since the ancillary study was conducted in women 65 to 79 years of age, it is unknown whether these findings apply to younger postmenopausal.

INDICATIONS AND USAGE

Vivelle is indicated in the following:

- Treatment of moderate to severe vasomotor symptoms associated with the menopause.
- Treatment of moderate to severe symptoms of vulvar and vaginal atrophy associated with the menopause. When prescribing solely for the treatment of symptoms of vulvar and vaginal atrophy, topical vaginal products should be considered.
- Treatment of hypoenestrogenism due to hypogonadism, castration, or primary ovarian failure.
- Prevention of postmenopausal osteoporosis. When prescribing solely for the prevention of postmenopausal osteoporosis, therapy should only be considered for women at significant risks of osteoporosis and non-estrogen medications should be carefully considered.

The mainstays for decreasing the risk of postmenopausal osteoporosis are weight-bearing exercise, adequate calcium and vitamin D intake, and when indicated, pharmacologic therapy. Postmenopausal women require an average of 1500 mg/day of elemental calcium. Therefore, when not contraindicated, calcium supplementation may be helpful for women with suboptimal dietary intake. Vitamin D supplementation of 400-800 IU/day may also be required to ensure adequate daily intake in postmenopausal women.

CONTRAINDICATIONS

Vivelle is contraindicated in women with any of the following conditions:

- Abnormal genital bleeding of unknown etiology.
- Known, suspected or history of cancer of the breast.
- Known or suspected estrogen-dependent neoplasia.
- Active deep vein thrombosis, pulmonary embolism or a history of these conditions.
- Active or recent (e.g., within the past year) arterial thromboembolic disease (e.g., stroke, myocardial infarction).

- Liver dysfunction or disease.
- Vivelle should not be used in patients with known hypersensitivity to its ingredients.

WARNINGS

1 Cardiovascular Disorders

Vivelle is contraindicated in women with active DVT, PE, stroke, or a history of these conditions [see Contraindications]. Immediately discontinue Vivelle if a PE, DVT, or stroke, occurs or is suspected. If feasible, discontinue Vivelle at least 4 to 6 weeks before surgery of the type associated with an increased risk of thromboembolism, or during periods of prolonged immobilization.

The safety and efficacy of Vivelle for the prevention of cardiovascular disorders have not been established.

The Women's Health Initiative (WHI) estrogen-alone trial reported increased risks of pulmonary embolism (PE), deep vein thrombosis (DVT), and stroke, in postmenopausal women (50 to 79 years of age, average age 63.4 years) during 7.2 years of treatment with daily oral conjugated estrogens (CE) [0.625 mg] relative to placebo. Analyses were also conducted in women aged 50-59 years, a group of women more likely to present with new onset of moderate to severe VMS compared to women of other age groups in the trial (see CLINICAL STUDIES).

Only daily oral 0.625 mg CE was studied in the WHI estrogen-alone trial. Therefore, the relevance of the WHI findings regarding adverse cardiovascular events to lower CE doses, other routes of administration, or other estrogen products is not known. Without such data, it is not possible to definitively exclude these risks or determine the extent of these risks for other products.

Venous Thromboembolism

In women aged 50-59 years, the WHI estrogen-alone trial reported a relative risk for PE of 1.53 (95% confidence interval [CI], 0.63, 3.75) for CE compared to placebo, with an absolute risk difference of 4 per 10,000 women-years (WYs; 10 versus 6). The relative risk for DVT was 1.66 (95% CI 0.75, 3.67) for CE compared to placebo, with a risk difference of 5 per 10,000 WYs (13 versus 8).

In the overall study population of women aged 50-79 years, the WHI estrogen-alone trial reported a relative risk of PE of 1.35 (95% CI 0.89, 2.05) for CE compared to placebo, with a risk difference of 4 per 10,000 WYs (14 versus 10). The relative risk for DVT was 1.48 (95% 1.06, 2.07) for CE compared to placebo, with a risk difference of 7 per 10,000 WYs (23 versus 15).

Stroke

In women aged 50-59 years, the WHI estrogen-alone trial reported a relative risk for stroke of 0.99 (95% 0.53, 1.85) for CE compared to placebo, with a risk difference of -1 per 10,000 WYs (16 versus 17).

In the overall study population of women aged 50-79 years, the WHI estrogen-alone trial reported a relative risk for stroke of 1.35 (95%, 1.07, 1.70) for CE compared to placebo, with a risk difference of 11 per 10,000 WYs (45 versus 34).

2 Malignant Neoplasms

Endometrial Cancer

In Vivelle-treated menopausal women with a uterus with persistent or recurring abnormal genital bleeding of unknown etiology, perform adequate diagnostic measures, including directed or random endometrial sampling when indicated, to assess for endometrial cancer.

An increased risk of endometrial cancer has been reported with the use of unopposed estrogen therapy in women with a uterus. The reported endometrial cancer risk among unopposed estrogen users is about 2 to 12 times greater than in non-users and appears dependent on duration of treatment and on estrogen dose. Most studies show no significant increased risk associated with the use of estrogens for less than 1 year. The greatest risk appears to be associated with prolonged use, with increased risks of 15- to 24-fold for 5 to 10 years or more. This risk has been shown to persist for at least 8 to 15 years after estrogen therapy is discontinued. There is no evidence that the use of natural estrogens results in a different endometrial risk profile than synthetic estrogens of equivalent estrogen dose.

Adding a progestogen to estrogen-alone therapy has been shown to reduce the risk of endometrial hyperplasia, which may be a precursor to endometrial cancer. There are, however, possible risks associated with the use of progestogens plus estrogens that differ from those of estrogen-alone regimens [*see Warnings (3)*].

Breast Cancer

Surveillance measures for breast cancer, such as breast examinations and mammography, are recommended. The use of estrogen-alone therapy has been reported to result in an increase in abnormal mammograms requiring further evaluation.

In the WHI estrogen-alone trial, after an average follow-up of 7.1 years, daily CE-alone was not associated with an increased risk of invasive breast cancer. Among women 50- 59 years old, the relative risk was 0.82 (95% CI, 0.50, 1.34) for CE compared to placebo, with a risk difference of -5 per 10,000 WYs (24 versus 29). In the overall study population of women aged 50-79 years (average age 63.4 years), the relative risk was 0.79 (95% CI, 0.61, 1.02), with a risk difference of -7 per 10,000 WYs (28 versus 35). [*see Clinical Studies (14.3)*]. However, a large meta-analysis including 24 prospective studies of postmenopausal women comparing current use of estrogen-only products with use duration of 5 to 14 years (average of 9 years) versus never use reported a relative risk for breast cancer of 1.33 (95% CI, 1.28 to 1.38).

Ovarian Cancer

A large meta-analysis including 17 prospective studies of postmenopausal women compared current use of estrogen-only products versus never use and reported a relative risk for ovarian cancer of 1.37 (95% CI, 1.26 to 1.50). The duration of hormone therapy use that was associated with an increased risk of ovarian cancer is unknown.⁶

3 Risks Associated with Concomitant Use of Estrogen Plus Progestogen

If Vivelle is administered with a progestogen, there are possible risks associated with the use of progestogens plus estrogens that differ from those of estrogen-alone regimens. Refer to prescribing information for progestogens indicated for the prevention of endometrial hyperplasia in non-hysterectomized menopausal women receiving estrogens.

Geriatric Use

There have not been sufficient numbers of geriatric women involved in clinical studies utilizing Vivelle to determine whether those over 65 years of age differ from younger subjects in their response to Vivelle.

The Women's Health Initiative Studies

In the WHI estrogen-alone trial (daily CE [0.625 mg]-alone versus placebo), there was a higher relative risk of stroke in women greater than 65 years of age (see CLINICAL STUDIES).

The Women's Health Initiative Memory Study

In the WHIMS ancillary studies of postmenopausal women 65 to 79 years of age, there was an increased risk of probable dementia in women receiving estrogen-alone (see CLINICAL STUDIES).

Since the trial was conducted in women 65 to 79 years of age, it is unknown whether these findings apply to younger menopausal women. The safety and efficacy of Vivelle for the prevention of dementia has not been established.

PRECAUTIONS

General

- ***Addition of a Progestin When a Woman Has Not Had a Hysterectomy.*** Studies of the addition of a progestin for 10 or more days of a cycle of estrogen administration, or daily with estrogen in a continuous regimen, have reported a lowered incidence of endometrial hyperplasia than would be induced by estrogen treatment alone. Endometrial hyperplasia may be a precursor to endometrial cancer. There are, however, possible risks that may be associated with the use of progestins with estrogens compared to estrogen alone regimens. These include a possible increased risk of breast cancer.
- ***Elevated Blood Pressure.*** In a small number of case reports, substantial increases in blood pressure have been attributed to idiosyncratic reactions to estrogens. In a large, randomized, placebo-controlled clinical trial, a generalized effect of estrogens on blood pressure was not seen. Blood pressure should be monitored at regular intervals with estrogen use.

- ***Hypertriglyceridemia.*** In patients with pre-existing hypertriglyceridemia, estrogen therapy may be associated with elevations of plasma triglycerides leading to pancreatitis and other complications.
- ***Impaired Liver Function and Past History of Cholestatic Jaundice.*** Although transdermally administered estrogen therapy avoids first-pass hepatic metabolism, estrogens may be poorly metabolized in patients with impaired liver function. For patients with a history of cholestatic jaundice associated with past estrogen use or with pregnancy, caution should be exercised and in the case of recurrence, medication should be discontinued.
- ***Hypothyroidism.*** Estrogen administration leads to increased thyroid-binding globulin (TBG) levels. Patients with normal thyroid function can compensate for the increased TBG by making more thyroid hormone, thus maintaining free T₄ and T₃ serum concentrations in the normal range. Patients dependent on thyroid hormone replacement therapy who are also receiving estrogens may require increased doses of their thyroid replacement therapy. These patients should have their thyroid function monitored in order to maintain their free thyroid hormone levels in an acceptable range.
- ***Fluid Retention.*** Because estrogens may cause some degree of fluid retention, patients with conditions that might be influenced by this factor, such as cardiac or renal dysfunction, warrant careful observation when estrogens are prescribed.
- ***Hypocalcemia.*** Estrogens should be used with caution in individuals with severe hypocalcemia.
- ***Exacerbation of Endometriosis.*** Endometriosis may be exacerbated with administration of estrogens. A few cases of malignant transformation of residual endometrial implants have been reported in women treated post-hysterectomy with estrogen alone therapy. For patients known to have residual endometriosis post-hysterectomy, the addition of progestin should be considered.
- ***Exacerbation of Other Conditions.*** Estrogens may cause an exacerbation of asthma, diabetes mellitus, epilepsy, migraine or porphyria, systemic lupus erythematosus, and hepatic hemangiomas and should be used with caution in women with these conditions.

Patient Information

Physicians are advised to discuss the Patient Information leaflet with patients for whom they prescribe Vivelle.

Laboratory Tests

Estrogen administration should be initiated at the lowest dose approved for the indication and then guided by clinical response, rather than by serum hormone levels (e.g., estradiol, FSH).

Drug/Laboratory Test Interactions

- Accelerated prothrombin time, partial thromboplastin time, and platelet aggregation time; increased platelet count; increased factors II, VII antigen, VIII antigen, VIII coagulant activity, IX, X, XII, VII-X complex, II-VII-X complex; and beta-thromboglobulin; decreased levels of anti-factor Xa and antithrombin III; decreased antithrombin III

activity; increased levels of fibrinogen and fibrinogen activity; increased plasminogen antigen and activity.

- Increased thyroid-binding globulin (TBG) leading to increased circulating total thyroid hormone levels as measured by protein-bound iodine (PBI), T₄ levels (by column or by radioimmunoassay) or T₃ levels by radioimmunoassay. T₃ resin uptake is decreased, reflecting the elevated TBG. Free T₄ and free T₃ concentrations are unaltered. Patients on thyroid replacement therapy may require higher doses of thyroid hormone.
- Other binding proteins may be elevated in serum, i.e., corticosteroid binding globulin (CBG), sex hormone binding globulin (SHBG), leading to increased circulating corticosteroids and sex steroids, respectively. Free hormone concentrations may be decreased. Other plasma proteins may be increased (angiotensinogen/renin substrate, alpha-1-antitrypsin, ceruloplasmin).
- Increased plasma HDL and HDL₂ cholesterol subfraction concentrations, reduced LDL cholesterol concentration, increased triglycerides levels.
- Impaired glucose tolerance.
- Reduced response to metyrapone test.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term continuous administration of estrogen, with and without progestin, in women with and without a uterus, has shown an increased risk of endometrial cancer, breast cancer, and ovarian cancer (see [BOXED WARNINGS](#), [WARNINGS](#) and [PRECAUTIONS](#)).

Long-term, continuous administration of natural and synthetic estrogens in certain animal species increases the frequency of carcinomas of the breast, uterus, cervix, vagina, testis, and liver.

Pregnancy

Vivelle should not be used during pregnancy (see [CONTRAINDICATIONS](#)).

Nursing Mothers

Estrogen administration to nursing mothers has been shown to decrease the quantity and quality of the milk. Detectable amounts of estrogens have been identified in the milk of mothers receiving this drug. Caution should be exercised when Vivelle is administered to a nursing woman.

Pediatric Use

Estrogen therapy has been used for the induction of puberty in adolescents with some forms of pubertal delay. Safety and effectiveness in pediatric patients have not otherwise been established.

Large and repeated doses of estrogen over an extended time period have been shown to accelerate epiphyseal closure, which could result in short adult stature if treatment is initiated before the completion of physiologic puberty in normally developing children. If estrogen is administered to patients whose bone growth is not complete, periodic monitoring

of bone maturation and effects on epiphyseal centers is recommended during estrogen administration.

Estrogen treatment of prepubertal girls also induces premature breast development and vaginal cornification and may induce vaginal bleeding (see [INDICATIONS](#) and [DOSAGE AND ADMINISTRATION](#)).

Geriatric Use

The safety and effectiveness in geriatric patients have not been established.

In the Women's Health Initiative Memory Study, including 4,532 women 65 years of age and older, followed for an average of four years, 82% (n=3,729) were 65 to 74 while 18% (n=803) were 75 and over. Most women (80%) had no prior hormone therapy use. Women treated with conjugated estrogens plus medroxyprogesterone acetate were reported to have a two-fold increase in the risk of developing probable dementia. Alzheimer's disease was the most common classification of probable dementia in both the conjugated estrogens plus medroxyprogesterone acetate group and the placebo group. Ninety percent of the cases of probable dementia occurred in the 54% of women that were older than 70 (see [WARNINGS, Dementia](#)). It is unknown whether these findings apply to estrogen alone therapy.

ADVERSE REACTIONS

See [BOXED WARNINGS](#), [WARNINGS](#) and [PRECAUTIONS](#).

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse events that appear to be related to drug use and for approximating rates.

The following adverse events have been reported with Vivelle therapy:

Table 3
Summary of Most Frequently Reported Adverse Experiences/Medical Events
Regardless of Relationship Reported at a Frequency =5%

	Vivelle 0.025 mg/ day [†] (N=47) N (%)	Vivelle 0.0375 mg/ day [†] (N=130) N (%)	Vivelle 0.05 mg/ day [†] (N=103) N (%)	Vivelle 0.075 mg/ day [†] (N=46) N (%)	Vivelle 0.1 mg/ day [†] (N=132) N (%)	Placebo (N=157) N (%)
Gastrointestinal Disorders						
Constipation	2 (4.3)	5 (3.8)	4 (3.9)	3 (6.5)	2 (1.5)	4 (2.5)
Dyspepsia	4 (8.5)	12 (9.2)	3 (2.9)	2 (4.3)	0	10 (6.4)
Nausea	2 (4.3)	8 (6.2)	4 (3.9)	0	7 (5.3)	5 (3.2)
General Disorders and Administration Site Conditions***						

Influenza-like Illness	3 (6.4)	6 (4.6)	8 (7.8)	0	3 (2.3)	10 (6.4)
Pain NOS*	0	8 (6.2)	0	2 (4.3)	7 (5.3)	7 (4.5)
Infections and Infestations						
Influenza	4 (8.5)	4 (3.1)	6 (5.8)	0	10 (7.6)	14 (8.9)
Nasopharyngitis	3 (6.4)	16 (12.3)	10 (9.7)	9 (19.6)	11 (8.3)	24 (15.3)
Sinusitis NOS*	4 (8.5)	17 (13.1)	13 (12.6)	3 (6.5)	7 (5.3)	16 (10.2)
Upper Respiratory Tract Infection NOS*	3 (6.4)	8 (6.2)	11 (10.7)	4 (8.7)	6 (4.5)	9 (5.7)
Investigations						
Weight Increased	4 (8.5)	5 (3.8)	2 (1.9)	2 (4.3)	0	3 (1.9)
Musculoskeletal and Connective Tissue Disorders						
Arthralgia	0	11 (8.5)	4 (3.9)	2 (4.3)	5 (3.8)	9 (5.7)
Back Pain	4 (8.5)	10 (7.7)	9 (8.7)	4 (8.7)	14 (10.6)	10 (6.4)
Neck Pain	3 (6.4)	4 (3.1)	4 (3.9)	0	6 (4.5)	2 (1.3)
Pain in Limb	0	10 (7.7)	7 (6.8)	2 (4.3)	6 (4.5)	9 (5.7)
Nervous System Disorders						
Headache NOS*	7 (14.9)	35 (26.9)	32 (31.1)	23 (50.0)	34 (25.8)	37 (23.6)
Sinus Headache	0	12 (9.2)	5 (4.9)	5 (10.9)	2 (1.5)	8 (5.1)
Psychiatric Disorders						
Anxiety NEC**	3 (6.4)	5 (3.8)	0	0	2 (1.5)	4 (2.5)
Depression	5 (10.6)	4 (3.1)	7 (6.8)	0	4 (3.0)	6 (3.8)
Insomnia	3 (6.4)	6 (4.6)	4 (3.9)	2 (4.3)	2 (1.5)	9 (5.7)
Reproductive System and Breast Disorders						
Breast Tenderness	8 (17.0)	10 (7.7)	8 (7.8)	3 (6.5)	17 (12.9)	0
Dysmenorrhea	0	0	0	3 (6.5)	0	0
Intermenstrual Bleeding	3 (6.4)	9 (6.9)	6 (5.8)	0	14 (10.6)	7 (4.5)
Respiratory, Thoracic and Mediastinal Disorders						
Sinus Congestion	0	4 (3.1)	3 (2.9)	3 (6.5)	6 (4.5)	7 (4.5)
Vascular Disorders						
Hot Flashes NOS*	3 (6.4)	0	3 (2.9)	0	0	6 (3.8)
Hypertension NOS*	2 (4.3)	0	3 (2.9)	0	0	2 (1.3)

† Represents milligrams of estradiol delivered daily by each system

* NOS represents not otherwise specified

** NEC represents not elsewhere classified

*** Application site erythema and application site irritation were observed in a small number of patients (3.2% or less of patients across treatment groups)

The following additional adverse reactions have been reported with estrogens and/or progestin therapy.

- **Genitourinary System.** Changes in vaginal bleeding pattern and abnormal withdrawal bleeding or flow; breakthrough bleeding; spotting; dysmenorrhea, increase in size of uterine leiomyomata; vaginitis, including vaginal candidiasis; change in amount of cervical secretion; changes in cervical ectropion; ovarian cancer; endometrial hyperplasia; endometrial cancer.
- **Breasts.** Tenderness, enlargement, pain, nipple discharge, galactorrhea; fibrocystic breast changes; breast cancer.
- **Cardiovascular.** Deep and superficial venous thrombosis; pulmonary embolism; thrombophlebitis; myocardial infarction; stroke; increase in blood pressure.
- **Gastrointestinal.** Nausea, vomiting; abdominal cramps, bloating; cholestatic jaundice; increased incidence of gallbladder disease; pancreatitis, enlargement of hepatic hemangiomas.
- **Skin.** Chloasma or melasma, which may persist when drug is discontinued; erythema multiforme; erythema nodosum; hemorrhagic eruption; loss of scalp hair; hirsutism; pruritus, rash.
- **Eyes.** Retinal vascular thrombosis; intolerance to contact lenses.
- **Central Nervous System.** Headache; migraine; dizziness; mental depression; chorea; nervousness; mood disturbances; irritability; exacerbation of epilepsy, dementia.
- **Miscellaneous.** Increase or decrease in weight; reduced carbohydrate tolerance; aggravation of porphyria; edema; arthralgias; leg cramps; changes in libido; anaphylactoid/anaphylactic reactions including urticaria and angioedema; hypocalcemia; exacerbation of asthma; increased triglycerides.

Post-Marketing Adverse Events

Although a causal relationship with Vivelle has not been established, adverse events reported from marketing experience include: isolated reports of anaphylaxis, rare elevated liver function tests, and reports of leg pain.

OVERDOSAGE

Serious ill effects have not been reported following acute ingestion of large doses of estrogen-containing drug products by young children. Overdosage of estrogen may cause nausea and vomiting, and withdrawal bleeding may occur in females.

DOSAGE AND ADMINISTRATION

Initiation of Therapy

The timing of Vivelle initiation can affect the overall benefit-risk profile. Consider initiating Vivelle in women < 60 years old or < 10 years since menopause onset (see CLINICAL STUDIES, WARNINGS, GERIATRIC USE).

When estrogen is prescribed for a menopausal woman with a uterus, the addition of a progestogen has been shown to reduce the risk of endometrial cancer. There are possible risks associated with the use of progestogens plus estrogens that differ from those of estrogen-alone regimens. See prescribing information for progestogens indicated for the prevention of endometrial hyperplasia in non-hysterectomized menopausal women receiving estrogens (see WARNINGS).

For treatment of moderate to severe vasomotor symptoms and vulvar and vaginal atrophy associated with the menopause, start therapy with Vivelle estradiol transdermal system 0.0375 mg/day applied to the skin twice weekly. For the prevention of postmenopausal osteoporosis, start therapy with Vivelle 0.025 mg/day applied to the skin twice weekly. The dosage may be adjusted as necessary. Reproductive system associated adverse events were encountered more frequently in the highest dose group (0.1 mg/day) than in other active treatment groups or in placebo-treated patients.

In women not currently taking oral estrogens or in women switching from another estradiol transdermal therapy, treatment with the Vivelle estradiol transdermal system may be initiated at once. In women who are currently taking oral estrogens, treatment with the Vivelle estradiol transdermal system should be initiated one week after withdrawal of oral hormone therapy, or sooner if menopausal symptoms reappear in less than one week.

Therapeutic Regimen

Vivelle may be given continuously in patients who do not have an intact uterus. In those patients with an intact uterus, Vivelle may be given continuously or on a cyclic schedule (e.g., three weeks on drug followed by one week off drug) with a progestin.

HOW SUPPLIED

Vivelle estradiol transdermal system 0.025 mg/day - each 7.25 cm² system contains 2.17 mg of estradiol USP for nominal* delivery of 0.025 mg of estradiol per day.

Patient Calendar Pack of 8 systems.....NDC xxxx-xx-xx

Vivelle estradiol transdermal system 0.0375 mg/day - each 11.0 cm² system contains 3.28 mg of estradiol USP for nominal* delivery of 0.0375 mg of estradiol per day.

Patient Calendar Pack of 8 systems.....NDC xxxx-xxxx-xx

Vivelle estradiol transdermal system 0.05 mg/day - each 14.5 cm² system contains 4.33 mg of estradiol USP for nominal* delivery of 0.05 mg of estradiol per day.

Patient Calendar Pack of 8 systems.....NDC xxxx-xxxx-xx

Vivelle estradiol transdermal system 0.075 mg/day - each 22.0 cm² system contains 6.57 mg of estradiol USP for nominal* delivery of 0.075 mg of estradiol per day.

Patient Calendar Pack of 8 systems.....NDC xxxx-xxxx-xx

Vivelle estradiol transdermal system 0.1 mg/day - each 29.0 cm² system contains 8.66 mg of estradiol USP for nominal* delivery of 0.1 mg of estradiol per day.

Patient Calendar Pack of 8 systems.....NDC xxxx-xxxx-xx

***See DESCRIPTION**

Do not store above 30°C (86°F). Do not store unpouched. Apply immediately upon removal from the protective pouch.

Rev. 10/2026

PATIENT INFORMATION

Vivelle®

estradiol transdermal system

Rx only

Read this PATIENT INFORMATION before you start using Vivelle® (estradiol transdermal system) and read all the information that you get each time you refill Vivelle. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

The Vivelle patch that your healthcare provider has prescribed for you releases small amounts of an estrogen hormone through the skin.

This leaflet describes the risks and benefits of treatment with Vivelle. Vivelle is not for everyone. Talk to your healthcare provider if you have any questions or concerns about this medication.

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT VIVELLE (AN ESTROGEN HORMONE)?

- Estrogens increase the chances of getting cancer of the uterus.

Report any unusual vaginal bleeding right away while you are taking estrogens. Vaginal bleeding after menopause may be a warning sign of cancer of the uterus (womb). Your healthcare provider should check any unusual vaginal bleeding to find out the cause.

-

You and your healthcare provider should talk regularly about whether you still need treatment with Vivelle

WHAT IS VIVELLE ?

Vivelle is a patch that contains the estrogen hormone, estradiol. When applied to the skin as directed below, Vivelle releases estrogen through the skin into the bloodstream.

WHAT IS VIVELLE USED FOR?

Vivelle is used after menopause to:

- Reduce moderate to severe hot flashes

Estrogens are hormones made by a woman's ovaries. The ovaries normally stop making estrogens when a woman is between 45 and 55 years old. This drop in body estrogen levels

causes the “change of life” or menopause (the end of monthly menstrual periods). Sometimes, both ovaries are removed during an operation before natural menopause takes place. The sudden drop in estrogen levels causes “surgical menopause.”

When the estrogen levels begin dropping, some women develop very uncomfortable symptoms, such as feelings of warmth in the face, neck, and chest or sudden strong feelings of heat and sweating (“hot flashes” or “hot flushes”). In some women the symptoms are mild, and they will not need estrogens. In other women, symptoms can be more severe. You and your healthcare provider should talk regularly about whether you still need treatment with Vivelle.

- Treat moderate to severe dryness, itching and burning in or around the vagina

You and your healthcare provider should talk regularly about whether you still need treatment with Vivelle to control these problems. If you use Vivelle only to treat your dryness, itching, and burning in and around your vagina, talk with your healthcare provider about whether a topical vaginal product would be better for you.

- Treat certain conditions in which a young woman’s ovaries do not produce enough estrogens naturally
- Help reduce your chances of getting osteoporosis (thin, weak bones)

Osteoporosis from menopause is a thinning of the bones that makes them weaker and easier to break. If you use Vivelle only to prevent osteoporosis from menopause, talk with your healthcare provider about whether a different treatment or medicine without estrogens might be better for you. You and your healthcare provider should talk regularly about whether you should continue with Vivelle.

Weight-bearing exercise, like walking or running, and taking calcium and vitamin D supplements may also lower your chances of getting postmenopausal osteoporosis. It is important to talk about exercise and supplements with your healthcare provider before starting them.

WHO SHOULD NOT TAKE VIVELLE?

Do not start taking Vivelle if you:

- Have unusual vaginal bleeding
- Currently have or have had certain cancers

Estrogens may increase the chances of getting certain types of cancers, including cancer of the breast or uterus. If you have or had cancer, talk with your healthcare provider about whether you should take Vivelle.

- Had a stroke or heart attack in the recent past (for example, in the past year)
- Currently have or have had blood clots
- Currently have or have had liver problems
- Are allergic to Vivelle or any of its ingredients

See the end of this leaflet for a list of ingredients in Vivelle.

- Think you may be, or know that you are, pregnant

TELL YOUR HEALTHCARE PROVIDER:

- If you are breastfeeding
The hormone in Vivelle can pass into your milk.
- About all of your medical problems

Your healthcare provider may need to check you more carefully if you have certain conditions such as asthma (wheezing), epilepsy (seizures), migraine, endometriosis, lupus, or problems with your heart, liver, thyroid, kidneys, or have high calcium levels in your blood.


- About all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Some medicines may affect how Vivelle works. Vivelle may also affect how other medicines work.
- If you are going to have surgery or will be on bed rest
You may need to stop taking estrogens.

HOW SHOULD I TAKE VIVELLE?

- Start at the lowest dose and talk to your healthcare provider about how well that dose is working for you.
- Estrogens should be used at the lowest dose possible for your treatment, only as long as needed. The lowest effective dose of Vivelle has not been determined for any indication. You and your healthcare provider should talk regularly (for example, every three to six months) about the dose you are taking and whether you still need treatment with Vivelle.

How and Where to Apply Vivelle

Each system is individually sealed in a protective pouch. Tear open this pouch at the indentation (do not use scissors) and remove the system.




A stiff protective liner covers the adhesive side of the system - the side that will be placed against your skin. This liner must be removed before applying the system. Hold the unit with the protective liner facing you.



Peel off one side of the protective liner and discard it. Try to avoid touching the sticky side of the system with your fingers.



Using the other half of the liner as a handle, apply the sticky side of the system to a dry area of the skin on the trunk of the body (including the abdomen or buttocks). Press the sticky side on the skin and smooth down.




Fold back the remaining side of the system. Grasp the straight edge of the protective liner and pull it off the system.



Press the system firmly in place.



OR



Some women may find that it is more comfortable to wear Vivelle on the buttocks. *Do not apply Vivelle to your breasts.* The sites of application must be rotated, with an interval of at least one week allowed between applications to a particular site. The area selected should not be oily, damaged, or irritated. Avoid the waistline since tight clothing may rub the system off. Apply the system immediately after opening the pouch and removing the protective liner. Press the system firmly in place with the palm of your hand for about 10 seconds, making sure there is good contact, especially around the edges.

The Vivelle system should be worn continuously until it is time to replace it with a new system. You may wish to experiment with different locations when applying a new system, to find ones that are most comfortable for you and where clothing will not rub on the system.

When to Apply Vivelle

The Vivelle system should be replaced twice weekly. Your Vivelle package contains a calendar checklist to help you remember a schedule. Mark the 2-day schedule you plan to follow. Always change the system on the two days of the week you have marked.

When changing the system, remove the used Vivelle system and discard it. Any adhesive that might remain on your skin can be easily rubbed off. Then place the new Vivelle system on a different skin site. (The same skin site should not be used again for at least one week after removal of the system.)

Please note: Contact with water when you are bathing, swimming, or showering will not affect the system. In the event that a system should fall off, put this same system back on and continue to follow your original treatment schedule. If necessary, you may apply a new system but continue to follow your original schedule.

WHAT ARE THE POSSIBLE SIDE EFFECTS OF ESTROGENS?

Less Common but Serious Side Effects Include:

- Breast Cancer
- Cancer of the Uterus
- Stroke
- Heart Attack
- Blood Clots

- Dementia
- Gallbladder Disease
- Ovarian Cancer

These are Some of the Warning Signs of Serious Side Effects:

- Breast Lumps
- Unusual Vaginal Bleeding
- Dizziness and Faintness
- Changes in Speech
- Severe Headaches
- Chest Pain
- Shortness of Breath
- Pains in Your Legs
- Changes in Vision
- Vomiting

Call your healthcare provider right away if you get any of these warning signs, or have any other unusual symptom that concerns you.

Common Side Effects Include:

- Headache
- Breast Pain
- Irregular Vaginal Bleeding or Spotting
- Stomach/Abdominal Cramps, Bloating
- Nausea and Vomiting
- Hair Loss

Other Side Effects Include:

- High Blood Pressure
- Liver Problems
- High Blood Sugar
- Fluid Retention
- Enlargement of Benign Tumors of the Uterus (“Fibroids”)

- Vaginal Yeast Infection

Other side effects of Vivelle may be possible. If you have questions, talk to your healthcare provider or pharmacist.

WHAT CAN I DO TO LOWER MY CHANCES OF A SERIOUS SIDE EFFECT WITH VIVELLE?

- Talk with your healthcare provider regularly about whether you should continue taking Vivelle.
- If you have a uterus, talk to your healthcare provider about whether the addition of a progestin is right for you.
- See your healthcare provider right away if you get vaginal bleeding while taking Vivelle.
- Have a breast exam and mammogram (breast X-ray) every year unless your healthcare provider tells you something else. If members of your family have had breast cancer or if you have ever had breast lumps or an abnormal mammogram, you may need to have breast exams more often.
- If you have high blood pressure, high cholesterol (fat in the blood), diabetes, are overweight, or if you use tobacco, you may have higher chances for getting heart disease. Ask your healthcare provider for ways to lower your chances for getting heart disease.

GENERAL INFORMATION ABOUT SAFE AND EFFECTIVE USE OF VIVELLE

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not take Vivelle for conditions for which it was not prescribed. Do not give Vivelle to other people, even if they have the same symptoms you have. It may harm them.

Keep Vivelle out of the reach of children.

This leaflet provides a summary of the most important information about Vivelle. If you would like more information, talk with your healthcare provider or pharmacist. You can ask for information about Vivelle that is written for health professionals. You can get more information by calling the toll-free number 1-800-525-8747

WHAT ARE THE INGREDIENTS IN VIVELLE?

The Vivelle system comprises three layers. Proceeding from the visible surface toward the surface attached to the skin, these layers are (1) a translucent flexible film consisting of an ethylene vinyl alcohol copolymer film, a polyurethane film, urethane polymer and epoxy resin, (2) an adhesive formulation containing estradiol USP, acrylic adhesive, polyisobutylene, ethylene vinyl acetate copolymer, 1,3 butylene glycol, styrene-butadiene rubber, oleic acid NF, lecithin, propylene glycol, bentonite NF, mineral oil USP, and dipropylene glycol, and (3) a polyester release liner that is attached to the adhesive surface and must be removed before the system can be used.

The active component of the system is estradiol. The remaining components of the system are pharmacologically inactive.

Rev. October 2026

Distributed by

Sandoz Inc., Princeton, NJ, 08540