

Patient Information
Retin-A Micro® (ret-in-A MY-kroe)
(tretinoin) Gel microsphere, 0.1%, 0.08%, 0.06%, and 0.04%
for topical use

Important information: Retin-A Micro is for use on skin only. Do not get Retin-A Micro in your eyes, mouth, vagina or the corners of your nose.

What is Retin-A Micro?

Retin-A Micro is a prescription medicine used on the skin (topical) to treat acne. Acne is a condition in which the skin has blackheads, whiteheads, and other pimples.

It is not known if Retin-A Micro is safe and effective in the treatment of other conditions.

It is not known if Retin-A Micro is safe and effective in children under 12 years of age.

What should I tell my doctor before using Retin-A Micro?

Before using Retin-A Micro, tell your doctor about all of your medical conditions, including if you:

- have a skin condition called eczema.
- have a sunburn. You should not use Retin-A Micro until your skin has healed.
- have any other medical condition.
- are pregnant or plan to become pregnant. It is not known if Retin-A Micro will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if Retin-A Micro passes into your breast milk.

Tell your doctor about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, herbal supplements, and any skin products that you use.

Especially tell your doctor if you use any other medicines to treat your acne, including medicated cleansers or soaps.

Using other topical acne products may increase the irritation of your skin when used with Retin-A Micro.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

How should I use Retin-A Micro?

- Use Retin-A Micro exactly as your doctor tells you to use it.
- Your doctor may change your dose of Retin-A Micro if you have skin irritation.
- Before you apply Retin-A Micro, gently wash the affected skin area with a mild, non-medicated soap. Rinse and pat your skin dry.
- Apply Retin-A Micro 1 time a day in the evening, or as prescribed by your doctor.
- Do not use more Retin-A Micro than you need to cover the affected area and do not apply Retin-A Micro more than 1 time a day. Using too much Retin-A Micro or using it too often will not give you faster or better results, and you may get skin redness, peeling, or discomfort.
- You may have a brief feeling of warmth or slight stinging after applying Retin-A Micro.
- You may use moisturizers and cosmetics.
- Early in your treatment, you may get new pimples. At this stage, it is important to continue using Retin-A Micro.
- Your acne may not get better right away. Use Retin-A Micro even after your acne improves. Your acne may get better after two weeks of treatment, but more than seven weeks of Retin-A Micro treatment are needed before you get the full benefit.

Applying Retin-A Micro:

- Retin-A Micro comes in a tube and a pump. If you have been prescribed the:

Tube: Squeeze the gel from the tube onto a fingertip. Apply a thin layer to cover the affected area, as prescribed by your doctor. Spread Retin-A Micro evenly over the affected area.

Pump: Fully depress the pump twice to dispense Retin-A Micro onto a fingertip. Apply a thin layer to cover the affected area, as prescribed by your doctor. Spread Retin-A Micro evenly over the affected area.

- Wash your hands after applying Retin-A Micro.

What should I avoid while using Retin-A Micro?

- Avoid washing your skin too often and scrubbing the affected skin area.
- You should avoid sunlamps, tanning beds, and ultraviolet light during treatment with Retin-A Micro.
- Minimize exposure to sunlight.
- If you have to be in the sunlight or are sensitive to sunlight, use a sunscreen with SPF (sun protection factor) of 15 or more and wear a wide-brimmed hat or other protective clothing to cover the treated areas.
- If you do get sunburned, stop using Retin-A Micro until your skin has healed and is back to normal.
- Cold weather and wind may irritate skin treated with Retin-A Micro. Talk to your doctor about ways to manage skin irritation.
- Avoid contact with the peels of limes.

What are the possible side effects of Retin-A Micro?

Retin-A Micro may cause serious side effects, including:

Skin irritation. Retin-A Micro may cause skin dryness, redness, swelling, and blistering. If you develop these symptoms, your doctor may tell you to stop using Retin-A Micro for a while, change your dose, decrease the number of times you apply Retin-A Micro, or completely stop treatment with Retin-A Micro. It is not known if Retin-A Micro is effective when used less than 1 time a day.

The most common side effects of Retin-A Micro include skin burning and itching.

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of Retin-A Micro. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Retin-A Micro?

- Store Retin-A Micro at room temperature between 68° to 77°F (20° to 25°C).
- Store Retin-A Micro pump upright.

Keep Retin-A Micro and all medicines out of the reach of children.

General information about the safe and effective use of Retin-A Micro.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use Retin-A Micro for a condition for which it was not prescribed. Do not give Retin-A Micro to other people, even if they have the same symptoms you have. It may harm them.

If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about Retin-A Micro that is written for health professionals.

For more information, call 1-800-321-4576.

What are the ingredients of Retin-A Micro?

Active ingredient: tretinoin

Inactive ingredients: benzyl alcohol, butylated hydroxytoluene, carbomer 974P, cyclomethicone and dimethicone copolyol, disodium EDTA, glycerin, methyl methacrylate/glycol dimethacrylate crosspolymer, PPG-20 methyl glucose ether distearate, propylene glycol, propylene glycol dicaprylate/dicaprate, purified water, sorbic acid, and trolamine

Distributed by:

Bausch Health US, LLC
Bridgewater, NJ 08807 USA

Manufactured by:

Bausch Health Companies Inc.
Laval, Quebec H7L 4A8, Canada

Retin-A Micro is a trademark of Bausch Health Companies Inc. or its affiliates.

Microsponge is a registered trademark of AMCOL International Corporation.

All other product/brand names and/or logos are trademarks of the respective owners.

© 2025 Bausch Health Companies Inc. or its affiliates

p/n

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: 4/2025

BAUSCH Health

DESCRIPTION: Retin-A Micro (Tretinoin Gel) - 0.04% - 45g - Carton - US - Laval

(b) (4)

(b) (4)

Each gram contains: Tretinoin 0.04% in benzyl alcohol, butylated hydroxytoluene, carbomer 974P, cyclomethicone and dimethicone copolyol, disodium EDTA, glycerin, methyl methacrylate/glycol dimethacrylate crosspolymer (MICROSPONGE® System), PPG-20 methyl glucose ether distearate, propylene glycol, propylene glycol dicaprylate/dicaprate, purified water, sorbic acid, and trolamine.

Microsponge is a registered trademark of AMCOL International Corporation.
Retin-A Micro and Ortho Dermatologics are trademarks of Bausch Health Companies Inc. or its affiliates.



3 01875 14445 6

Important: The opening of this product is covered by a metal seal. If this seal has been punctured or is not visible, do not use and return to place of purchase.
To open: To puncture the seal, reverse the cap and place the puncture-top onto the tube. Push down firmly until seal is open. To close, screw the cap back onto the tube.

For Topical Use Only **WARNING: Keep out of reach of children.**
Rx only **Store at 20° to 25°C (68° to 77°F); excursions permitted from**
Usual Dosage: See prescribing information. **15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].**



RETIN-A MICRO®
(tretinoin) Gel microsphere
0.04%

NDC 0187-5144-45

For Topical Use Only
Rx only

Net Wt. 45 g

Distributed by:
Bausch Health US, LLC
Bridgewater, NJ 08807 USA

Manufactured by:
Bausch Health Companies Inc.
Laval, Quebec H7L 4A8, Canada
Product of Germany

© 2024 Bausch Health Companies Inc. or its affiliates


Ortho | Dermatologics 9457804
50104769C

**LOT and EXP
LASER BURNED
ON BLACK FLAP
SANS-SERIF FONT; 6pts**
**LOT0000000
EXPYYYY/MM**

BAUSCH Health

DESCRIPTION: Retin-A Micro (Tretinoin Gel) - 0.04% - 45g - Tube - US - Laval

(b) (4)



(b) (4)

Usual Dosage:
See prescribing information.

WARNING: Keep out of reach of children.
See side of label for lot number and expiration date.

Store at 20° to 25°C (68° to 77°F); excursions permitted from 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Store pump upright.

Each gram contains:
Tretinoin 0.04% in benzyl alcohol, butylated hydroxytoluene, carbomer 974P, cyclomethicone and dimethicone copolyol, disodium EDTA, glycerin, methyl methacrylate/glycol dimethacrylate crosspolymer (MICROSPONGE® System), PPG-20 methyl glucose ether distearate, propylene glycol, propylene glycol dicaprylate/dicaprate, purified water, sorbic acid, and trolamine.

Distributed by:
Bausch Health US, LLC
Bridgewater, NJ 08807 USA

Manufactured by:
Bausch Health Companies Inc.
Laval, Quebec H7L 4A8, Canada

Microsponge is a registered trademark of AMCOL International Corporation.
© 2024 Bausch Health Companies Inc. or its affiliates
9450806

NDC 0187-5144-50

RETIN-A MICRO®
(tretinoin) Gel microsphere
0.04%
PUMP

For Topical Use Only
Rx only

Net Wt. 50 g

Ortho | Dermatologics

00000000
YYYY/MM

50104774E

3 018751 14450 10

OVERLAPPING AREA


LOT and EXP PRINTED ON WHITE AREA,
SANS-SERIF FONT; 7pts
LOT0000000
EXPYYYY/MM

AREA FOR LOT/EXP and DATAMATRIX MUST BE WHITE

(b) (4)

LOT and EXP
LASER BURNED
ON BLACK FLAP
SANS-SERIF FONT; 9pts

LOT0000000
EXPYYYY/MM



NDC 0187-5144-50

RETIN-A MICRO[®]
(tretinoin) Gel microsphere
0.04%
PUMP

For Topical Use Only
Rx only

Rx only
Usual Dosage: See prescribing information.
WARNING: Keep out of reach of children.
Store at 20° to 25°C (68° to 77°F); excursions permitted from 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Store pump upright.
Each gram contains: Tretinoin 0.04% in benzyl alcohol, butylated hydroxytoluene, carbomer 974P, cyclomethicone and dimethicone copolyol, disodium EDTA, glycerin, methyl methacrylate/glycol dimethacrylate crosspolymer (MICROSPONGE[®] System), PPG-20 methyl glucose ether distearate, propylene glycol, propylene glycol dicaprylate/dicaprate, purified water, sorbic acid, and tromamine.
 Microsponge is a registered trademark of AMCOL International Corporation.
Distributed by:
 Bausch Health US, LLC
 Bridgewater, NJ 08807 USA
Manufactured by:
 Bausch Health Companies Inc.
 Laval, Quebec H7L 4A8, Canada
 Product of Germany
 Retin-A Micro and Ortho Dermatologics are trademarks of Bausch Health Companies Inc. or its affiliates.
 © 2024 Bausch Health Companies Inc. or its affiliates



9450906

Net Wt. 50 g

Ortho | Dermatologics



BAUSCH Health

DESCRIPTION: Retin-A Micro (Tretinoin Gel) - 0.04% - 50g - Pump - Carton - US - Laval

(b) (4)



David
Lewis

Digitally signed by David Lewis

Date: 4/22/2025 10:33:58AM

GUID: 508da72000029f287fa31e664741b577