Important: Read all product information before using. Keep this box for important information. This product is intended for use in adults.

Drug Facts

Active ingredient (in each 10 mL) — **Purposes**

Ibuprofen 200 mg (NSAID)* ..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - minor pain of arthritis
  - backache
  - menstrual cramps
  - muscular aches
  - toothache
  - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if
- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product
- take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
- **this product does not contain directions or complete warnings for use in children under 12 years of age.**
- **do not take more than directed**
- **the smallest effective dose should be used**
- shake well before using
- mL = milliliter
- remove bottle cap and squeeze your dose into the dosing cup
- use only enclosed dosing cup. Do not use any other dosing device
- replace original bottle cap to maintain child resistance

adults and children 12 years and older	■ take 10 mL every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 10 mL, an additional 10 mL may be used ■ do not exceed 60 mL (1200 mg) in 24 hours, unless directed by a doctor
children under 12 years	■ Do not use

Other information
- **each 10 mL contains:** sodium 4 mg
- store between 20-25°C (68-77°F)
- **do not use if carton is opened. Do not use if bottle wrap imprinted with "Sealed For Your Safety" is broken or missing**

Inactive ingredients

acesulfame potassium, anhydrous citric acid, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions?

Call **1-877-896-3665** (toll-free) or **215-273-8755** (collect) or visit www.motrin.com

OPEN HERE ➡

Motrin® IB

Ibuprofen Oral Suspension
200 mg per 10 mL
Pain Reliever/Fever Reducer **(NSAID)**

NDC 50580-769-60

NEW

Motrin® IB

Ibuprofen Oral Suspension
200 mg per 10 mL
Pain Reliever/Fever Reducer **(NSAID)**

Adult Liquid

Alcohol Free
Dye Free

Wild Berry Flavor

8 fl oz (240 mL)

Motrin® IB

STEP 1. See *Drug Facts* for complete directions and dosing

STEP 2. Shake well before using

STEP 3. Remove bottle cap

STEP 4. Squeeze the dose into the dosing cup

STEP 5. Replace original bottle cap to maintain child resistance

Distributed by:
Kenvue Brands LLC
Summit, NJ 07901
© Kenvue Brands LLC 2026
Pat. www.kenvuepats.com

Important: Read all product information before using. Keep this box for important information. This product is intended for use in adults.

Drug Facts

Active Ingredient (In each 10 mL) — **Purposes**

Ibuprofen 200 mg (NSAID)* ..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. When using this product ■ take with food or milk if stomach upset occurs. Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ this product does not contain directions or complete warnings for use in children under 12 years of age. ■ **do not take more than directed** ■ **the smallest effective dose should be used** ■ shake well before using ■ mL = milliliter ■ remove bottle cap and squeeze your dose into the dosing cup ■ use only enclosed dosing cup. Do not use any other dosing device. ■ replace original bottle cap to maintain child resistance. Adults and children 12 years and older ■ take 10 mL every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 10 mL, an additional 10 mL may be used ■ do not exceed 60 mL (1200 mg) in 24 hours, unless directed by a doctor. Children under 12 years ■ Do not use.

Other information ■ each 10 mL contains: sodium 4 mg ■ store between 20-25°C (68-77°F) ■ **do not use if bottle wrap imprinted with "Sealed For Your Safety" is broken or missing.**

Inactive Ingredients acesulfame potassium, anhydrous citric acid, flavors, glycerin, polyvinyl alcohol-90, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions? Call 1-877-895-3685 (toll-free) or 215-273-8755 (collect) or visit www.motrin.com

Distributed by: Kenvue Brands LLC Summit, NJ 07901

© Kenvue Brands LLC 2026

OOOOOOOO

NDC 50580-769-08

Motrin® IB

Ibuprofen Oral Suspension 200 mg per 10 mL

Pain Reliever/Fever Reducer (NSAID)

Adult Liquid

Wild Berry Flavor

8 fl oz (240 mL)