



NPA019378

REGULATORY LEGEND	
	Actual Size
DRUG FACTS TITLE	9 pt
DRUG FACTS CONTINUED	8 pt
HEADINGS	8 pt
SUBHEADINGS/BODY TEXT	6 pt
LEADING	6.5 pt
BULLETS	5 pt
# OF CHARACTERS PER INCH	33
SPACE BEFORE BULLET	2 ems
BARLINES, HAIRLINES	1.5, 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	13.5 pt
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

Reference ID: 5822361



NPA019382

REGULATORY LEGEND	
	Actual Size
DRUG FACTS TITLE	9 pt
DRUG FACTS CONTINUED	8 pt
HEADINGS	8 pt
SUBHEADINGS/BODY TEXT	6 pt
LEADING	6.5 pt
BULLETS	5 pt
# OF CHARACTERS PER INCH	33
SPACE BEFORE BULLET	2 ems
BARLINES, HAIRLINES	1.5, 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	27 pt
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A



NPA019380

REGULATORY LEGEND	
	Actual Size
DRUG FACTS TITLE	9 pt
DRUG FACTS CONTINUED	8 pt
HEADINGS	8 pt
SUBHEADINGS/BODY TEXT	6, 6 pt
LEADING	6.5 pt
BULLETS	5 pt
# OF CHARACTERS PER INCH	27
SPACE BEFORE BULLET	2 ems
BARLINES, HAIRLINES	1.5, 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	31.5 pt
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	N/A

Do not use if printed sleeve is broken or missing.

Adults and ages 12+

HALEON

Advil

LIQUID

Ibuprofen Oral Suspension

200 mg per 10 mL Pain Reliever/
Fever Reducer (NSAID)

White Grape
Flavor

ALCOHOL-FREE
DYE-FREE

8 FL.OZ (240 mL)

Active ingredient (in each 10 mL).....Ibuprofen 200 mg

Uses

• temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis.

• temporarily reduces fever

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you, are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery

Ask a doctor before use if stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers or fever reducers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin; taking any other drug

When using this product take with food or milk if stomach upset occurs

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Each 10 mL contains: sodium 7 mg

Store at 20-25° C (68-77° F).

Directions

This product does not contain directions or complete warnings for use by children under 12 years of age. Do not take more than directed. The smallest effective dose should be used. Shake well before using. Adults and children 12 years and over: take 10 mL every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 10 mL, 20 mL may be taken. Do not exceed 60 mL in 24 hours, unless directed by a doctor. Measure only with the dosing cup provided. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension. Children under 12 years: ask a doctor

Questions or comments?
1-855-297-3031

Distributed by: Haleon, Warren, NJ 07060.

For most recent product information, visit www.Advil.com

Trademarks are owned by or licensed to the Haleon group of companies.

©2026 Haleon group of companies or its licensor.

NPA019379

NPA019379

REGULATORY LEGEND	
	Actual Size
DRUG FACTS TITLE	N/A
DRUG FACTS CONTINUED	N/A
HEADINGS	6 pt
SUBHEADINGS/BODY TEXT	5 pt
LEADING	5.4 pt
BULLETS	5 pt
# OF CHARACTERS PER INCH	48
SPACE BEFORE BULLET	N/A
BARLINES, HAIRLINES	1.5 pt, N/A
SPACE BETWEEN HAIRLINES AND BOX END	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	15.4 pt

Reference ID: 5822361