Reference ID: 5828079

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredients (in each extended release tablet) **Purpose**

Cetirizine HCl 5 mg.....Antihistamine
Pseudoephedrine HCl 120 mg.....Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
  - nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- **do not use more than directed**
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- store between 20° to 25°C (68° to 77°F)
- **do not use if blister unit is torn or broken**

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions? call **1-800-343-7805** (toll-free) or **215-273-8755** (collect)

3 00450 20427 1

LOT: XXXXXXX
EXP: YYYY/MMM

Made in Belgium
Distributed by: Kenvue Brands LLC
Summit, NJ 07901
© Kenvue Brands LLC 2026
www.zyrtec.com **30061090**

The trade dress of this ZYRTEC® package is subject to trademark protection.
www.kenvuepats.com

ZYRTEC-D® ALLERGY + SINUS

Original Prescription Strength

ZYRTEC-D®

NDC 50580-719-12

Cetirizine HCl **5 mg** /antihistamine
Pseudoephedrine HCl **120 mg** /nasal decongestant
Extended Release Tablets

Indoor & Outdoor Allergies

ALLERGY + SINUS

NASAL CONGESTION + SINUS PRESSURE

12 hour relief of

Sneezing

Runny Nose

Itchy, Watery Eyes

Itchy Throat or Nose

Actual Size

12 EXTENDED RELEASE TABLETS

30061090

5006 1090

Reference ID: 5828079

ZYRTEC-D® ALLERGY + SINUS

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredients (in each extended release tablet) Purpose

Cetirizine HCl 5 mg.....Antihistamine
Pseudoephedrine HCl 120 mg.....Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
  - nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- drowsiness may occur
- **do not use more than directed**
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

If pregnant or breast-feeding:
- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- store between 20° to 25°C (68° to 77°F)
- **do not use if blister unit is torn or broken**

The trade dress of this ZYRTEC® package 30061092 is subject to trademark protection.

www.kenvuepats.com

LOT: XXXXXXX

EXP: YYYY/MMM

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions? call **1-800-343-7805** (toll-free) or **215-273-8755** (collect)

Made in Belgium
Distributed by: Kenvue Brands LLC
Summit, NJ 07901
© Kenvue Brands LLC 2026

www.zyrtec.com

3 00450 20424 0