



Bayer, the Bayer Cross, Claritin, and the Claritin Vial are registered trademarks of Bayer.
 © 2025 Bayer. Diet by: Bayer HealthCare LLC
 Whippany, NJ 07981 P.O. Box 111111, Whippany, NJ 07981
 00000000 Bayer

Juniors Non-Drowsy*
Claritin
 Chewables

MAX Strength 10mg 40 CHEWABLE TABLETS

MAX Strength 10mg

Juniors Ages 6+ years

Claritin
 Chewables

Indoor & Outdoor Allergies

24hr Non-Drowsy* Relief of:

DYE FREE
 GRAPE FLAVORED

40 CHEWABLE TABLETS



*When taken as directed. See Drug Facts Panel.

MAX Strength 10mg

Juniors Ages 6+ years

Claritin
 Chewables

DYE FREE
 GRAPE FLAVORED

40 CHEWABLE TABLETS

00000000



NO VARNISH

Drug Facts

Active ingredient (in each chewable tablet) Purpose
 Loratadine 10 mgAntihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ Itchy, watery eyes
 ■ sneezing ■ Itching of the nose or throat

Warnings
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ chew or crush tablets completely before swallowing.

adults and children 6 years and over	chew 1 tablet daily; not more than 1 chewable tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

■ phenylethanolic: contains phenylalanine 3 mg per chewable tablet
 ■ safety sealed: do not use if the individual blister unit imprinted with Claritin® Chewables is open or torn
 ■ store between 20° to 25°C (68° to 77°F)

Inactive ingredients

aspartame, citric acid anhydrous, colloidal silicon dioxide, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Questions or comments?
 1-800-CLARITIN (1-800-252-7484)

LOT/EXP/MADE IN
 LOT/EXP AREA
 COUNTRY OF ORIGIN AREA
 NO VARNISH
 NO INK

NO VARNISH



00000000

Glue Area



Juniors Non-Drowsy*
Claritin
 Chewables

MAX Strength 10mg 60 CHEWABLE TABLETS

MAX Strength 10mg

Juniors Ages 6+ years

Claritin
 Chewables

Indoor & Outdoor Allergies

24hr Non-Drowsy* Relief of:

DYE FREE
 GRAPE FLAVORED

60 CHEWABLE TABLETS



*When taken as directed. See Drug Facts Panel.

MAX Strength 10mg

Juniors Ages 6+ years

Claritin
 Chewables

DYE FREE
 GRAPE FLAVORED

60 CHEWABLE TABLETS

Drug Facts

Active ingredient (in each chewable tablet) Purpose
 Loratadine 10 mgAntihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ Itchy, watery eyes
 ■ sneezing ■ Itching of the nose or throat

Warnings
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ chew or crush tablets completely before swallowing.	
adults and children 6 years and over	chew 1 tablet daily; not more than 1 chewable tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

■ phenylethanolic: contains phenylalanine 3 mg per chewable tablet
 ■ safety sealed: do not use if the individual blister unit imprinted with Claritin® Chewables is open or torn
 ■ store between 20° to 25°C (68° to 77°F)

Inactive ingredients

aspartame, citric acid anhydrous, colloidal silicon dioxide, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Questions or comments?
 1-800-CLARITIN (1-800-252-7484)

LOT/EXP AREA
 COUNTRY OF ORIGIN AREA
 NO VARNISH
 NO INK

LOT/EXP MADE IN

NO VARNISH



00000000

00000000



NO VARNISH

Glue Area

Juniors ⁶⁺ years

Non-Drowsy*
Claritin®
Chewables

loratadine chewable tablets
10 mg/antihistamine



ACTUAL SIZE

MAX Strength 10mg

24hr Non-Drowsy* Relief of:


Sneezing • Runny Nose • Itchy, Watery Eyes • Itchy Throat or Nose

Indoor & Outdoor Allergies



* When taken as directed. See Drug Facts Panel.

80
CHEWABLE
TABLETS



Bayer
000 00000
Pat. Reg. U.S. Pat. & Tm. Off. © 2020 Bayer HealthCare LLC, Whippany, NJ 07981
Bayer, the Bayer Cross, Claritin, the Claritin Vials, and registered trademarks of Bayer
© 2020 Bayer
Dist. by: Bayer HealthCare LLC
Whippany, NJ 07981

Drug Facts (continued)	
Active ingredient (in each chewable tablet)	Loratadine 10 mg; Antihistamine
Purpose	Chew or crush tablets completely before swallowing.
Directions	adults and children 6 years and over 1 chewable tablet in 24 hours ask a doctor children under 6 years of age ask a doctor consumers with liver or kidney disease
Warnings	Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Other information	phenylethanolamine 3 mg per chewable tablet ask a doctor Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Questions or comments?	1-800-CLARITIN (1-800-252-7484)

• Sneezing
• Runny Nose
• Itchy, Watery Eyes
• Itchy Throat or Nose
24hr Non-Drowsy* Relief of:



ACTUAL SIZE

Chewables

Claritin®

Non-Drowsy*

Juniors ⁶⁺ years

Juniors ⁶⁺ years

Non-Drowsy*
Claritin®
Chewables

loratadine chewable tablets
10 mg/antihistamine



ACTUAL SIZE

MAX Strength 10mg

24hr Non-Drowsy* Relief of:


Sneezing • Runny Nose • Itchy, Watery Eyes • Itchy Throat or Nose

Indoor & Outdoor Allergies



* When taken as directed. See Drug Facts Panel.

90
CHEWABLE
TABLETS



Bayer
000 00000
Bayer, the Bayer Cross, Claritin,
the Claritin Vials, are registered
trademarks of Bayer.
© 2020 Bayer.
Distributed by: Bayer Healthcare LLC
Whisper, NJ 07981
Bayer Pharmaceuticals Inc.
P.O. Box 100
Parsippany, NJ 07981

Drug Facts (continued)	
Active ingredient (in each chewable tablet)	Loratadine 10 mg; Antihistamine
Purpose	Chew or crush tablets completely before swallowing.
Directions	adults and children 6 years and over 1 chewable tablet in 24 hours children under 6 years of age ask a doctor consumers with liver or kidney disease ask a doctor
Warnings	Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Other information	phenylethanolamine 3 mg per chewable tablet may cause drowsiness. Do not use if the individual blister unit is printed with "Do not use if you have ever had an allergic reaction to this product or any of its ingredients." Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Questions or comments?	1-800-CLARITIN (1-800-252-7484)

24hr Non-Drowsy* Relief of:
• Sneezing
• Runny Nose
• Itchy, Watery Eyes
• Itchy Throat or Nose

ACTUAL SIZE
Claritin®
Non-Drowsy*
Chewables

Juniors ⁶⁺ years

Juniors ⁶⁺ **AGES**

Non-Drowsy*
Claritin®
Chewables

loratadine chewable tablets
10 mg/antihistamine



ACTUAL SIZE

MAX Strength 10mg


24hr Non-Drowsy* Relief of:
Sneezing • Runny Nose • Itchy, Watery Eyes • Itchy Throat or Nose

Indoor & Outdoor Allergies



*When taken as directed. See Drug Facts Panel.

100
CHEWABLE
TABLETS



Bayer

000-00000
Pat. : patents, lwevll.bayer.com
Whippany, NJ 07981
Dist. by: Bayer Healthcare LLC
© 2025 Bayer.
the Claritin tests are registered
trademarks of Bayer.
Bayer, the Bayer Cross, Claritin

Drug Facts (continued)	
Active ingredient (in each chewable tablet) Loratadine 10 mg, Antihistamine	Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Purpose Loratadine 10 mg, Antihistamine	Uses Importantly relieves these symptoms due to hay fever or other upper respiratory allergies: • itchy, watery eyes • sneezing • itching of the nose or throat
Directions • chew or crush tablets completely before swallowing.	Other information • pregnancy category: contains phenylbutane 3 mg per chewable tablet • safety seal: do not use if the individual blister unit imprinted with • storage between 20° to 25° C (68° to 77° F) • Claritin Chewables is open to tampering.
Questions or comments? 1-800-CLARITIN (1-800-252-7444)	Inactive ingredients aspartame, citric acid, erythrosine, colloidal silicon dioxide, flavor, glycolate, stearic acid

- Sneezing
 - Runny Nose
 - Itchy, Watery Eyes
 - Itchy Throat or Nose
- 24hr Non-Drowsy* Relief of:**

ACTUAL SIZE



Chewables

Claritin®

Non-Drowsy*

Juniors
Ages **6+** years

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
05/20/2026 08:49:51 AM