




LOT: XXXXX
EXP: YYY/MMM



Active ingredient (in each 5 mL): Cetirizine HCl 5 mg. **Purpose:** Antihistamine. **Uses:** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat. **Warnings:** **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **When using this product:** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. **Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding, not recommended ■ if pregnant, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions:** ■ use only with enclosed dosing cup ■ find right dose on chart below ■ mL = milliliter ■ adults and children 6 years and over: 5 mL or 10 mL once daily depending upon severity of symptoms; do not take more than 10 mL in 24 hours. ■ adults 65 years and over: 5 mL once daily; do not take more than 5 mL in 24 hours. ■ children 2 to under 6 years of age: 2.5 mL once daily. If needed, dose can be increased to a maximum of 5 mL once daily or 2.5 mL every 12 hours. Do not give more than 5 mL in 24 hours. ■ children under 2 years of age: ask a doctor ■ consumers with liver or kidney disease: ask a doctor. **Other information:** ■ store between 20° to 25°C (68° to 77°F) ■ do not use if bottle wrap imprinted with "Sealed For Your Safety" is broken or missing Active made in Switzerland. Dist. by: Kenvue Brands LLC, Summit, NJ 07901 © Kenvue Brands LLC 2026 30060419





Uses

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product

- drowsiness may occur
- avoid alcoholic drinks

- Be careful when driving a motor vehicle or operating machinery
- In rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens.



NDC 50580-730 05

if pregnant or breast-feeding: ■ if breast-feeding: not recommended
■ if pregnant: ask a health professional before use.

Directions
■ use only with enclosed dosing cup ■ ml = milliliter ■ adults and children

Other information

- store between 20° to 25°C (68° to 77°F)
- do not use if bottle wrap imprinted with "Sealed For Your Safety" is broken or missing

Questions? call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)
Dosing cup should be washed and left to air dry after each use.



Drug Facts	
Active ingredient (in each 5 mL)	Purpose Antihistamine
Uses Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: - runny nose - sneezing - itchy, watery eyes - itching of throat or throat	
Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to other antihistamines containing hydroxyzine.	
Ask a doctor before you use if you have liver or kidney disease or if your doctor should determine if you need a different dose.	
Ask a doctor or pharmacist before use if you are taking other prescription or over-the-counter drugs, or sedatives.	
When using this product - When drowsiness may occur - Do not drink alcohol - Do not drink grapefruit juice - Alcohol, sedatives, and tranquilizers may increase drowsiness - Do not operate a motor vehicle or other machinery until you know how you react to this medicine - Do not operate machinery or drive a motor vehicle or other operating machine after taking this medicine - Do not drink grapefruit juice during or after therapy - Consult your physician if you experience prolonged or persistent hives or other allergic reaction	
Stop use and ask a doctor if: you have an allergic reaction to this product; or seek medical attention if these happen: - hives - swelling of the face, lips, or tongue - difficulty breathing	
If pregnant or breast-feeding: • breast-feeding: not recommended • pregnant: ask a health professional for advice before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)	

Drug Facts (continued)	
Directions	
• only with enclosed dosing cup • right dose on chart below • 15 mL = 1 tablespoon	
• children • ages 1 to 5 • ask a doctor	• 5 mL, or 10 mL, once daily depending on severity of symptoms; do not take more than 15 mL in 24 hours.
• ask a doctor • ages 6 years and over	• 5 mL once daily; do not take more than 5 mL in 24 hours.
• children • ages 6 to 11 • ask a doctor	• 2.5 mL once daily. If needed, dose can be increased to a maximum of 5 mL, once daily for 2 to 3 days. • Do not give more than 5 mL in 24 hours.
• children under 2 years of age • ask a doctor	
• consumers with liver or kidney disease • ask a doctor	
Other information	
• store between 20° to 25° (68° to 77° F) • do not use if cap is opened. Do not use if cap is damaged or if seal is broken. • Your Safety is protected or missing • use bottom cap for lot number and expiration date	
Inactive ingredients	
anhydrous citric acid, fencone, propylene glycol, polyethylene glycol, sodium benzoate, sorbitol, sodium lauryl sulfate, sucrose.	
Questions? 1-800-343-7805 (toll-free) 1-213-273-8755 (local)	

Children's ZYRTEC
 Allergy

Children's ZYRTEC
 Allergy

Bubble Gum Flavor

Children's ZYRTEC
 Cetirizine HCl oral solution
 1 mg/ml (antihistamine)

Children's ZYRTEC
 Cetirizine HCl oral solution
 1 mg/ml (antihistamine)

Indoor & Outdoor Allergies

2 yrs & older

24 Hour

Relief of

Sneezing

Runny Nose

Itchy, Watery Eyes

Itchy Throat or Nose

Dye-Free Sugar-Free Alcohol-Free

Indoor & Outdoor Allergies

2 yrs & older

24 Hour

Relief of

Sneezing

Runny Nose

Itchy, Watery Eyes

Itchy Throat or Nose

Dye-Free Sugar-Free Alcohol-Free

Bubble Gum Flavor

4 fl oz (118 ml)

Dosing cup included

Bubble Gum Flavor

4 fl oz (118 ml)

Dosing cup included

LOT: XXXXXXXX
EXP: YYYY/MMM



Active ingredient (in each 5 mL) **Purpose**
Cetirizine HCl 5 mg.....Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ sneezing
■ itchy, watery eyes ■ itching of the nose or throat



LOT: XXXXXX
EXP: YYY/MMM

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.
When using this product
■ drowsiness may occur
■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens.

NDC 50580-721-05

Children's

ZYRTEC®

Cetirizine HCl oral solution
1 mg/mL antihistamine

ALLERGY

24
Hour

Relief of

Sneezing

Runny Nose

Itchy, Watery Eyes

Itchy Throat or Nose

Dye-Free
Sugar-Free
Alcohol-Free

Indoor & Outdoor Allergies

2 yrs & older

Bubble Gum Flavor

4 fl oz (118 mL)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding:
■ if breast-feeding: not recommended
■ if pregnant: ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions
■ use only with enclosed dosing cup ■ mL = milliliter ■ adults and children 6 years and over: 5 mL or 10 mL once daily depending upon severity of symptoms; do not take more than 10 mL in 24 hours. ■ adults 65 years and over: 5 mL once daily; do not take more than 5 mL in 24 hours. ■ children 2 to under 6 years of age: 2.5 mL once daily. If needed, dose can be increased to a maximum of 5 mL once daily or 2.5 mL every 12 hours. Do not give more than 5 mL in 24 hours. ■ children under 2 years of age: ask a doctor ■ consumers with liver or kidney disease: ask a doctor
Other information
■ store between 20° to 25°C (68° to 77°F)
■ do not use if bottle wrap imprinted with "Sealed For Your Safety" is broken or missing
Inactive ingredients
anhydrous citric acid, flavors, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose
Questions? call 1-800-343-7805 (toll-free) or 215-273-8755 (collect).
Dosing cup should be washed and left to air dry after each use.
Active ingredient made in Switzerland
Distributed by: **Kenvue Brands LLC** Summit, NJ 07901
© Kenvue Brands LLC 2025 www.zyrtec.com 30060994



Active ingredient (in each 5 mL)
Cetirizine HCl 5 mg.....
Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.
Warnings Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms: ■ trouble swallowing ■ dizziness or loss of consciousness ■ swelling of tongue ■ swelling in or around mouth ■ trouble speaking ■ drooling ■ wheezing or problems breathing These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times. **Do not use ■ to prevent** hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. ■ if you have ever had an allergic reaction to this product or any of its ingredients or to

Purpose
Antihistamine

LOT: XXXXXX
EXP: YYYY/MMMM

NDC 50580-128 04

Children's

ZYRTEC®

Cetirizine HCl 1 mg/mL HIVES

oral solution antihistamine

6 yrs & older

HIVES RELIEF

24 Hour relief

Reduces Itching Due to Hives

Reduces Hives

Grape Flavor

4 fl oz (118 mL)

an antihistamine containing hydroxyzine. **Ask a doctor before use if you have** ■ liver or kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch **Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping cetirizine. Consult your healthcare provider if this happens. **Stop use and ask a doctor if** ■ an allergic reaction to this product occurs. Seek medical help right away. ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted more than 6 weeks **If pregnant or breast-feeding:** ■ if breast-feeding, not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions:** ■ use only with enclosed dosing cup ■ find right dose on chart below ■ mL = milliliter **Adults and children 6 years and over:** 5 mL or 10 mL once daily depending upon severity of symptoms; do not take more than 10 mL in 24 hours. **Adults 65 years and over:** 5 mL once daily; do not take more than 5 mL in 24 hours. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information** ■ store between 20° to 75° (68° to 77°F) ■ **do not use if bottle wrap imprinted with "Sealed For Your Safety" is broken or missing.** **Inactive ingredients** anhydrous citric acid, flavors, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose. **Questions?** call 1-800-343-7808 (toll-free) or 215-273-8755 (collect). **Dosing cup should be washed and left to air dry after each use.** Active ingredient made in Switzerland. The trade dress of this ZYRTEC® package is subject to trademark protection. Distributed by: **KENVUE BRANDS LLC** Summit, NJ 07901 © Kenvue Brands LLC 2025 **zyrtec.com** Pat. www.kenvuepats.com 30059467

Reference ID: 5833116

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

DOROTHY CHANG
07/13/2026 02:19:30 PM