

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use HYCOFENIX® safely and effectively. See [full prescribing information](#) for HYCOFENIX.

HYCOFENIX (hydrocodone bitartrate, pseudoephedrine hydrochloride, and guaifenesin) oral solution, CII
Initial U.S. Approval: 2015

WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; MEDICATION ERRORS; CYTOCHROME P450 3A4 INTERACTION; CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS; INTERACTION WITH ALCOHOL; NEONATAL OPIOID WITHDRAWAL SYNDROME

See full prescribing information for complete boxed warning.

- HYCOFENIX exposes users to risks of addiction, abuse, and misuse, which can lead to overdose and death. Assess patient's risk before prescribing and monitor closely for these behaviors and conditions. (5.1)
- Serious, life-threatening, or fatal respiratory depression may occur. Monitor closely, especially upon initiation or when used in patients at higher risk. (5.2)
- Accidental ingestion of HYCOFENIX, especially by children, can result in a fatal overdose of hydrocodone. (5.2)
- Ensure accuracy when prescribing, dispensing, and administering HYCOFENIX. Dosing errors can result in accidental overdose and death. (2.1, 5.5)
- Concomitant use with CYP3A4 inhibitors (or discontinuation of CYP3A4 inducers) can result in a fatal overdose of hydrocodone. Avoid the use of HYCOFENIX in patients taking CYP3A4 inhibitors or inducers. (5.7, 7.2, 7.3)
- Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Avoid the use of HYCOFENIX in patients taking benzodiazepines, other CNS depressants, or alcohol. (5.8, 7.4)
- Instruct patients not to consume alcohol or any products containing alcohol while taking HYCOFENIX because co-ingestion can result in fatal plasma hydrocodone levels. (5.8, 7.1)
- HYCOFENIX is not recommended for use in pregnant women. Advise pregnant women using opioids for an extended period of time of the risk of Neonatal Opioid Withdrawal Syndrome, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery. (5.14, 8.1)

RECENT MAJOR CHANGES

Boxed Warning	12/2025
Indications and Usage (1)	12/2025
Dosage and Administration (2.3)	12/2025
Warnings and Precautions (5.1, 5.2, 5.8, 5.9)	12/2025

INDICATIONS AND USAGE

HYCOFENIX is a combination of hydrocodone, an opioid agonist; pseudoephedrine, an alpha-adrenergic agonist; and guaifenesin, an expectorant, indicated for the symptomatic relief of cough, nasal congestion, and to loosen mucus associated with the common cold in patients 18 years of age and older. (1)

Limitations of Use

- Not indicated for pediatric patients under 18 years of age. (1)
- Because of the risks of addiction, abuse, misuse, overdose, and death, which can occur at any dosage or duration and persist over the course of therapy, reserve HYCOFENIX for use in adult patients for whom alternative treatment options are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of cough. (1, 5.1)

DOSAGE AND ADMINISTRATION

- Adults 18 years of age and older: 10 mL every 4 to 6 hours as needed, not to exceed 4 doses (40 mL) in 24 hours. (2.2)

- Measure HYCOFENIX with an accurate milliliter measuring device. (2.1, 5.5)
- Do not increase the dose or dosing frequency. (2.1)
- Prescribe for the shortest duration consistent with treatment goals. (2.3)
- Reevaluate patients with unresponsive cough in 5 days or sooner for possible underlying pathology. (2.3)
- Reevaluate patient prior to refilling. (2.3)
- Do not rapidly reduce or abruptly discontinue in a physically-dependent patient. (2.3)

DOSAGE FORMS AND STRENGTHS

Oral solution: Each 5 mL contains hydrocodone bitartrate 2.5 mg; pseudoephedrine hydrochloride 30 mg; and guaifenesin 200 mg. (3)

CONTRAINDICATIONS

- Children younger than 6 years of age. (4)
- Significant respiratory depression. (4)
- Acute or severe bronchial asthma in an unmonitored setting or in absence of resuscitative equipment. (4)
- Known or suspected gastrointestinal obstruction, including paralytic ileus. (4)
- Patients with narrow angle glaucoma, urinary retention, severe hypertension, or severe coronary artery disease (4)
- Hypersensitivity to hydrocodone, pseudoephedrine, guaifenesin, or any of the inactive ingredients in HYCOFENIX. (4)

WARNINGS AND PRECAUTIONS

- Life-threatening respiratory depression in patients with chronic pulmonary disease or in elderly, cachectic, or debilitated patients:** Monitor closely, particularly during initiation of therapy. (5.4)
- Activities requiring mental alertness:** Avoid engaging in hazardous tasks requiring mental alertness such as driving or operating machinery. (5.6)
- Risks of use in patients with head injury, impaired consciousness, increased intracranial pressure, or brain tumors:** Avoid use. May increase intracranial pressure and obscure the clinical course of head injuries. (5.10)
- Cardiovascular and central nervous system effects:** Use with caution in patients with cardiovascular disorders. (5.11)
- Seizures in patients with seizure disorders:** Monitor during therapy. (5.12)
- Severe hypotension:** Monitor during initiation of therapy. Avoid use in patients with circulatory shock. (5.13)
- Adrenal insufficiency:** If diagnosed, treat with physiologic replacement of corticosteroids, and wean patient off of the opioid. (5.15)

ADVERSE REACTIONS

Common adverse reactions include: Sedation (somnolence, mental clouding, lethargy), impaired mental and physical performance, lightheadedness, dizziness, headache, dry mouth, nausea, vomiting, constipation, tachycardia, arrhythmias including premature ventricular contractions, CNS stimulation including anxiety, restlessness, nervousness, tremor, and irritability. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Chartwell RX Sciences, LLC. at 1-845-268-5000 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Serotonergic Drugs:** Concomitant use may result in serotonin syndrome. Discontinue if serotonin syndrome is suspected. (7.5)
- Monoamine Oxidase Inhibitors (MAOIs):** Can potentiate the effects of hydrocodone. Avoid concomitant use in patients receiving MAOIs or within 14 days of stopping an MAOI. (7.6)
- Muscle relaxants:** Avoid concomitant use. (7.7)
- Diuretics:** Hydrocodone may reduce the efficacy of diuretics. Monitor for reduced effect. (7.8)
- Anticholinergic drugs:** Concurrent use may cause paralytic ileus. (5.9, 7.9)
- Antihypertensive drugs:** Concomitant use may interfere with antihypertensive effects. (7.10)
- Digitalis:** Concomitant use may increase ectopic pacemaker activity. (7.11)

USE IN SPECIFIC POPULATIONS

- Pregnancy:** Avoid use in pregnant women. May cause fetal harm. (8.1)
- Lactation:** Breastfeeding not recommended. (8.2)
- Renal Impairment:** Use with caution in patients with severe renal impairment. (8.6)
- Hepatic Impairment:** Use with caution in patients with severe hepatic impairment. (8.7)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; MEDICATION ERRORS; CYTOCHROME P450 3A4 INTERACTION; CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS; INTERACTION WITH ALCOHOL; NEONATAL OPIOID WITHDRAWAL SYNDROME

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Dosage and Administration Instructions
- 2.2 Recommended Dosage
- 2.3 Monitoring, Maintenance, and Discontinuation of Therapy

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Addiction, Abuse, and Misuse
- 5.2 Life-Threatening Respiratory Depression
- 5.3 Risks of Use in Pediatric Populations
- 5.4 Risks of Use in Other At-Risk Populations
- 5.5 Risks of Accidental Overdose and Death due to Medication Errors
- 5.6 Activities Requiring Mental Alertness: Risks of Driving and Operating Machinery
- 5.7 Risks from Concomitant Use or Discontinuation of Cytochrome P450 3A4 Inhibitors and Inducers
- 5.8 Risks from Concomitant Use with Benzodiazepines or other CNS Depressants
- 5.9 Risks of Gastrointestinal Complications
- 5.10 Risks of Use in Patients with Head Injury, Impaired Consciousness, Increased Intracranial Pressure, or Brain Tumors
- 5.11 Cardiovascular and Central Nervous System Effects
- 5.12 Increased Risk of Seizures in Patients with Seizure Disorders
- 5.13 Severe Hypotension
- 5.14 Neonatal Opioid Withdrawal Syndrome
- 5.15 Adrenal Insufficiency
- 5.16 Drug/Laboratory Test Interactions

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

- 7.1 Alcohol
- 7.2 Inhibitors of CYP3A4 and CYP2D6
- 7.3 CYP3A4 Inducers
- 7.4 Benzodiazepines, and Other CNS Depressants
- 7.5 Serotonergic Drugs
- 7.6 Monoamine Oxidase Inhibitors (MAOIs)
- 7.7 Muscle Relaxants
- 7.8 Diuretics
- 7.9 Anticholinergic Drugs
- 7.10 Antihypertensive Drugs
- 7.11 Digitalis

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

9 DRUG ABUSE AND DEPENDENCE

- 9.1 Controlled Substance
- 9.2 Abuse
- 9.3 Dependence

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; MEDICATION ERRORS; CYTOCHROME P450 3A4 INTERACTION; CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS; INTERACTION WITH ALCOHOL; NEONATAL OPIOID WITHDRAWAL SYNDROME

Addiction, Abuse, and Misuse

HYCOFENIX exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Reserve HYCOFENIX for use in adult patients for whom the benefits of cough suppression are expected to outweigh the risks, and in whom an adequate assessment of the etiology of the cough has been made. Assess each patient's risk prior to prescribing HYCOFENIX, prescribe HYCOFENIX for the shortest duration that is consistent with individual patient treatment goals, monitor all patients regularly for the development of addition or abuse, and refill only after reevaluation of the need for continued treatment. [see *Warnings and Precautions (5.1)*]

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of HYCOFENIX. Monitor for respiratory depression, especially during initiation of HYCOFENIX therapy or when used in patients at higher risk [see *Warnings and Precautions (5.2)*].

Accidental Ingestion

Accidental ingestion of even one dose of HYCOFENIX, especially by children, can result in a fatal overdose of hydrocodone [see *Warnings and Precautions (5.2)*].

Risk of Medication Errors

Ensure accuracy when prescribing, dispensing, and administering HYCOFENIX. Dosing errors can result in accidental overdose and death. Always use an accurate milliliter measuring device when measuring and administering HYCOFENIX [see *Dosage and Administration (2.1)*, *Warnings and Precautions (5.5)*].

Cytochrome P450 3A4 Interaction

The concomitant use of HYCOFENIX with all cytochrome P450 3A4 inhibitors may result in an increase in hydrocodone plasma concentrations, which could increase or prolong adverse drug effects and may cause potentially fatal respiratory depression. In addition, discontinuation of a concomitantly used cytochrome P450 3A4 inducer may result in an increase in hydrocodone plasma concentration. Avoid the use of HYCOFENIX in patients taking a CYP3A4 inhibitor or inducer [see *Warnings and Precautions (5.7)*, *Drug Interactions (7.2, 7.3)*].

Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Avoid the use of HYCOFENIX in patients taking benzodiazepines, other CNS depressants, or alcohol. [see *Warning and Precautions (5.8)*, *Drug Interactions (7.4)*]

Interaction with Alcohol

Instruct patients not to consume alcoholic beverages or use prescription or non-prescription products that contain alcohol while taking HYCOFENIX. The co-ingestion of alcohol with HYCOFENIX may result in increased plasma levels and a potentially fatal overdose of hydrocodone [see *Warnings and Precautions (5.8)* and *Drug Interactions (7.1)*].

Neonatal Opioid Withdrawal Syndrome

HYCOFENIX is not recommended for use in pregnant women [see *Use in Specific Populations (8.1)*]. Advise pregnant women using opioids for an extended period of time of the risk of Neonatal Opioid Withdrawal Syndrome, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery [see *Warnings and Precautions (5.14)*]

1 INDICATIONS AND USAGE

HYCOFENIX is indicated for the symptomatic relief of cough, nasal congestion, and to loosen mucus associated with the common cold in patients 18 years of age and older.

Limitations of Use:

- Not indicated for pediatric patients under 18 years of age [see *Use in Specific Populations (8.4)*].
- Contraindicated in pediatric patients less than 6 years of age [see *Contraindications (4)*].
- Because of the risks of addiction, abuse, misuse, overdose, and death, which can occur at any dosage or duration and persist over the course of therapy [see *Warnings and Precautions (5.1)*], reserve HYCOFENIX for use in adult patients for whom alternative treatment options are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of cough.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Instructions

Administer HYCOFENIX by the oral route only.

Always use an accurate milliliter measuring device when administering HYCOFENIX to ensure that the dose is measured and administered accurately. A household teaspoon is not an accurate measuring device and could lead to overdosage [see *Warnings and Precautions (5.5)*]. For prescriptions where a measuring device is not provided, a pharmacist can provide an appropriate measuring device and can provide instructions for measuring the correct dose. Do not overfill. Rinse the measuring device with water after each use.

Advise patients not to increase the dose or dosing frequency of HYCOFENIX because serious adverse events such as respiratory depression may occur with overdosage [see *Warnings and Precautions (5.2)*, *Overdosage (10)*]. The dosage of HYCOFENIX should not be increased if cough fails to respond; an unresponsive cough should be reevaluated for possible underlying pathology [see *Dosage and Administration (2.3)*, *Warnings and Precautions (5.4)*].

2.2 Recommended Dosage

Adults 18 years of age and older: 10 mL every 4 to 6 hours as needed, not to exceed 4 doses (40 mL) in 24 hours.

2.3 Monitoring, Maintenance, and Discontinuation of Therapy

Prescribe HYCOFENIX for the shortest duration that is consistent with individual patient treatment goals [see *Warnings and Precautions (5.1)*].

Monitor patients closely for respiratory depression, especially within the first 24-72 hours of initiating therapy [see *Warnings and Precautions (5.2)*].

Reevaluate patients with unresponsive cough in 5 days or sooner for possible underlying pathology, such as foreign body or lower respiratory tract disease [see *Warnings and Precautions (5.4)*]. If a patient requires a refill, reevaluate the cause of the cough and assess the need for continued treatment with HYCOFENIX, the relative incidence of adverse reactions, and the development of addiction, abuse, or misuse [see *Warnings and Precautions (5.1)*].

Do not rapidly reduce or abruptly discontinue HYCOFENIX in a physically-dependent patient [see *Drug Abuse and Dependence (9.3)*]. When a patient who has been taking HYCOFENIX regularly and may be physically dependent no longer requires therapy with HYCOFENIX, taper the dose gradually, by 25% to 50% every 2 to 4 days, while monitoring carefully for signs and symptoms of withdrawal. If the patient develops these signs or symptoms, raise the dose to the previous level and taper more slowly, either by increasing the interval between decreases, decreasing the amount of change in dose, or both.

3 DOSAGE FORMS AND STRENGTHS

Oral solution: Each 5 mL contains hydrocodone bitartrate, USP, 2.5 mg; pseudoephedrine hydrochloride, USP, 30 mg; and guaifenesin, USP, 200 mg [see *Description (11)*].

4 CONTRAINDICATIONS

HYCOFENIX is contraindicated for:

- All children younger than 6 years of age [see *Warnings and Precautions (5.2, 5.3), Use in Specific Populations (8.4)*].

HYCOFENIX is also contraindicated in patients with:

- Significant respiratory depression [see *Warnings and Precautions (5.2)*].
- Acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment [see *Warnings and Precautions (5.4)*].
- Known or suspected gastrointestinal obstruction, including paralytic ileus [see *Warnings and Precautions (5.9)*].
- Narrow angle glaucoma, urinary retention, severe hypertension, or severe coronary artery disease [see *Warnings and Precautions (5.11)*].
- Hypersensitivity to hydrocodone, pseudoephedrine, guaifenesin, or any of the inactive ingredients in HYCOFENIX [see *Adverse Reactions (6)*].

5 WARNINGS AND PRECAUTIONS

5.1 Addiction, Abuse, and Misuse

HYCOFENIX contains hydrocodone, a Schedule II controlled substance. As an opioid, HYCOFENIX exposes users to the risks of addiction, abuse, and misuse [see *Drug Abuse and Dependence (9)*], which can lead to overdose and death [see *Overdosage (10)*]. **Reserve HYCOFENIX for use in adult patients for whom the benefits of cough suppression are expected to outweigh the risks, and in whom an adequate assessment of the etiology of the cough has been made. Assess each patient's risk prior to prescribing HYCOFENIX, prescribe HYCOFENIX for the shortest duration that is consistent with individual patient treatment**

goals, monitor all patients regularly for the development of addiction or abuse, and refill only after reevaluation of the need for continued treatment.

Although the risk of addiction in any individual is unknown, it can occur in patients appropriately prescribed HYCOFENIX. Addiction can occur at recommended dosages and if the drug is misused or abused. The risk of opioid-related overdose or overdose-related death is increased with higher opioid doses, and this risk persists over the course of therapy. In postmarketing studies, addiction, abuse, misuse, and fatal and non-fatal opioid overdose were observed in patients with long-term opioid use [*see Adverse Reactions (6)*]. Risks are increased in patients with a personal or family history of substance abuse (including drug or alcohol abuse or addiction) or mental illness (e.g., major depression).

Opioids are sought by drug abusers and people with addiction disorders and are subject to criminal diversion. Consider these risks when prescribing or dispensing HYCOFENIX. Strategies to reduce these risks include prescribing the drug in the smallest appropriate quantity and advising the patient on the proper disposal of unused drug [*see Patient Counseling Information (17)*]. Contact local state professional licensing board or state controlled substances authority for information on how to prevent and detect abuse or diversion of this product.

5.2 Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression has been reported with the use of opioids, including hydrocodone, one of the active ingredients in HYCOFENIX. Hydrocodone produces dose-related respiratory depression by directly acting on the brain stem respiratory center that controls respiratory rhythm and may produce irregular and periodic breathing. Respiratory depression, if not immediately recognized and treated, may lead to respiratory arrest and death. Management of respiratory depression includes discontinuation of HYCOFENIX, close observation, supportive measures, and use of opioid overdose reversal agents (e.g., naloxone or nalmefene), depending on the patient's clinical status [*see Overdosage (10)*]. Carbon dioxide (CO₂) retention from opioid- induced respiratory depression can exacerbate the sedating effects of opioids.

While serious, life-threatening, or fatal respiratory depression can occur at any time during the use of HYCOFENIX, the risk is greatest during the initiation of therapy, when HYCOFENIX is used concomitantly with other drugs that may cause respiratory depression [*see Warnings and Precautions (5.8)*], in patients with chronic pulmonary disease or decreased respiratory reserve, and in patients with altered pharmacokinetics or altered clearance (e.g. elderly, cachectic, or debilitated patients) [*see Warnings and Precautions (5.4)*].

To reduce the risk of respiratory depression, proper dosing of HYCOFENIX is essential [*see Dosage and Administration (2.1)*, *Warnings and Precautions (5.5)*]. Monitor patients closely, especially within the first 24-72 hours of initiating therapy or when used in patients at higher risk.

Overdose of hydrocodone in adults has been associated with fatal respiratory depression, and the use of hydrocodone in children younger than 6 years of age has been associated with fatal respiratory depression when used as recommended. Accidental ingestion of even one dose of HYCOFENIX, especially by children, can result in respiratory depression and death.

5.3 Risks with Use in Pediatric Populations

Children are particularly sensitive to the respiratory depressant effects of hydrocodone [*see Warnings and Precautions (5.2)*]. Because of the risk of life-threatening respiratory depression and death, HYCOFENIX is contraindicated in children less than 6 years of age [*see Contraindications (4)*].

Use of HYCOFENIX in children also exposes them to the risks of addiction, abuse, and misuse [*see Drug Abuse and Dependence (9)*], which can lead to overdose and death [*see Warnings and Precautions (5.1)*, *Overdosage (10)*]. Because the benefits of symptomatic treatment of cough associated with allergies or the common cold do not outweigh the risks of use of hydrocodone in pediatric patients, HYCOFENIX is not indicated for use in patients younger than 18 years of age [*see Indications (1)*, *Use in Specific Populations (8.4)*].

5.4 Risks with Use in Other At-Risk Populations

Unresponsive Cough

The dosage of HYCOFENIX should not be increased if cough fails to respond; an unresponsive cough should be reevaluated in 5 days or sooner for possible underlying pathology, such as foreign body or lower respiratory tract disease [see *Dosage and Administration* (2.3)].

Asthma and Other Pulmonary Disease

The use of HYCOFENIX in patients with acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment is contraindicated [see *Contraindications* (4)].

Opioid analgesics and antitussives, including hydrocodone, one of the active ingredients in HYCOFENIX, should not be used in patients with acute febrile illness associated with productive cough or in patients with chronic respiratory disease where interference with ability to clear the tracheobronchial tree of secretions would have a deleterious effect on the patient's respiratory function.

HYCOFENIX-treated patients with significant chronic obstructive pulmonary disease or cor pulmonale, and those with a substantially decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory depression are at increased risk of decreased respiratory drive including apnea, even at recommended dosages of HYCOFENIX [see *Warnings and Precautions* (5.2)].

Elderly, Cachectic, or Debilitated Patients: Life-threatening respiratory depression is more likely to occur in elderly, cachectic, or debilitated patients because they may have altered pharmacokinetics or altered clearance compared to younger, healthier patients [see *Warnings and Precautions* (5.2)].

Because of the risk of respiratory depression, avoid the use of opioid antitussives, including HYCOFENIX in patients with compromised respiratory function, patients at risk of respiratory failure, and in elderly, cachectic, or debilitated patients. If HYCOFENIX is prescribed, monitor such patients closely, particularly when initiating HYCOFENIX and when HYCOFENIX is given concomitantly with other drugs that depress respiration [see *Warnings and Precautions* (5.8)].

5.5 Risk of Accidental Overdose and Death due to Medication Errors

Dosing errors can result in accidental overdose and death. To reduce the risk of overdose and respiratory depression, ensure that the dose of HYCOFENIX is communicated clearly and dispensed accurately [see *Dosage and Administration* (2.1)].

Advise patients to always use an accurate milliliter measuring device when measuring and administering HYCOFENIX. Inform patients that household teaspoon is not an accurate measuring device and such use could lead to overdosage and serious adverse reactions [see *Overdosage* (10)]. For prescriptions where a measuring device is not provided, a pharmacist can provide an appropriate calibrated measuring device and can provide instructions for measuring the correct dose.

5.6 Activities Requiring Mental Alertness: Risks of Driving and Operating Machinery

Hydrocodone, one of the active ingredients in HYCOFENIX, may produce marked drowsiness and impair the mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery. Advise patients to avoid engaging in hazardous tasks requiring mental alertness and motor coordination after ingestion of HYCOFENIX. Avoid concurrent use of HYCOFENIX with alcohol or other central nervous system depressants because additional impairment of central nervous system performance may occur [see *Warnings and Precautions* (5.8)].

5.7 Risks from Concomitant Use or Discontinuation of Cytochrome P450 3A4 Inhibitors and Inducers

Concomitant use of HYCOFENIX with a CYP3A4 inhibitor, such as macrolide antibiotics (e.g., erythromycin), azole-antifungal agents (e.g., ketoconazole), and protease inhibitors (e.g., ritonavir), may increase plasma concentrations of hydrocodone and prolong opioid adverse reactions, which may cause potentially fatal

respiratory depression [*see Warnings and Precautions (5.2)*], particularly when an inhibitor is added after a stable dose of HYCOFENIX is achieved. Similarly, discontinuation of a CYP3A4 inducer, such as rifampin, carbamazepine, and phenytoin, in HYCOFENIX-treated patients may increase hydrocodone plasma concentrations and prolong opioid adverse reactions.

Concomitant use of HYCOFENIX with CYP3A4 inducers or discontinuation of an CYP3A4 inhibitor could decrease hydrocodone plasma concentrations, decrease opioid efficacy or, possibly, lead to a withdrawal syndrome in a patient who had developed physical dependence to hydrocodone.

Avoid the use of HYCOFENIX in patients who are taking a CYP3A4 inhibitor or inducer. If concomitant use of HYCOFENIX with a CYP3A4 inhibitor or inducer is necessary, monitor patients for signs and symptoms that may reflect opioid toxicity and opioid withdrawal [*see Drug Interactions (7.2, 7.3)*].

5.8 Risks from Concomitant Use with Benzodiazepines or other CNS Depressants

Concomitant use of opioids, including HYCOFENIX, with benzodiazepines, gabapentinoids (gabapentin or pregabalin), or other CNS depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Because of these risks, avoid use of opioid cough medications in patients taking benzodiazepines, other CNS depressants, or alcohol [*see Drug Interactions (7.1, 7.4)*].

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioids alone. Because of similar pharmacologic properties, it is reasonable to expect similar risk with concomitant use of opioid cough medications and benzodiazepines, gabapentinoids (gabapentin or pregabalin), other CNS depressants, or alcohol.

Advise both patients and caregivers about the risks of respiratory depression and sedation if HYCOFENIX is used with benzodiazepines, gabapentinoids (gabapentin or pregabalin), alcohol, or other CNS depressants [*see Patient Counseling Information (17)*].

Patients must not consume alcoholic beverages, or prescription or non-prescription products containing alcohol, while on HYCOFENIX therapy. The co-ingestion of alcohol with HYCOFENIX may result in increased plasma levels and a potentially fatal overdose of hydrocodone [*see Drug Interactions (7.1, 7.4)*].

5.9 Risks of Gastrointestinal Complications

HYCOFENIX is contraindicated in patients with known or suspected gastrointestinal obstruction, including paralytic ileus [*see Contraindications (4)*]. The use of hydrocodone in HYCOFENIX may obscure the diagnosis or clinical course of patients with acute abdominal conditions.

The concurrent use of anticholinergics with HYCOFENIX may produce paralytic ileus [*see Drug Interactions (7.9)*].

The hydrocodone in HYCOFENIX may result in constipation or obstructive bowel disease, especially in patients with underlying intestinal motility disorders. Use with caution in patients with underlying intestinal motility disorders.

The hydrocodone in HYCOFENIX may cause spasm of the sphincter of Oddi, resulting in an increase in biliary tract pressure. Opioids may cause increases in serum amylase [*see Warnings and Precautions (5.16)*]. Monitor patients with biliary tract disease, including acute pancreatitis for worsening symptoms.

Cases of opioid-induced esophageal dysfunction (OIED) have been reported in patients taking opioids. The risk of OIED may increase as the dose and/or duration of opioids increases. Regularly evaluate patients for signs and symptoms of OIED (e.g., dysphagia, regurgitation, non-cardiac chest pain), and if necessary, adjust opioid therapy as clinically appropriate [*see Clinical Pharmacology (12.2)*].

5.10 Risks of Use in Patients with Head Injury, Impaired Consciousness, Increased Intracranial Pressure, or Brain Tumors

Avoid the use of HYCOFENIX in patients with head injury, intracranial lesions, or a pre-existing increase in intracranial pressure. In patients who may be susceptible to the intracranial effects of CO₂ retention (e.g., those with evidence of increased intracranial pressure or brain tumors), HYCOFENIX may reduce respiratory drive, and the resultant CO₂ retention can further increase intracranial pressure. Furthermore, opioids produce adverse reactions that may obscure the clinical course of patients with head injuries.

5.11 Cardiovascular and Central Nervous System Effects

The pseudoephedrine contained in HYCOFENIX can produce cardiovascular and central nervous system effects in some patients such as, insomnia, dizziness, weakness, tremor, transient elevations in blood pressure, or arrhythmias. In addition, central nervous system stimulation with convulsions or cardiovascular collapse with accompanying hypotension has been reported. Therefore, HYCOFENIX is contraindicated in patients with severe hypertension or coronary artery disease [*see Contraindications (4)*], and should, be used with caution in patients with other cardiovascular disorders.

5.12 Increased Risk of Seizures in Patients with Seizure Disorders

The hydrocodone in HYCOFENIX may increase the frequency of seizures in patients with seizure disorders, and may increase the risk of seizures occurring in other clinical settings associated with seizures. Monitor patients with a history of seizure disorders for worsened seizure control during HYCOFENIX therapy.

5.13 Severe Hypotension

HYCOFENIX may cause severe hypotension including orthostatic hypotension and syncope in ambulatory patients. There is increased risk in patients whose ability to maintain blood pressure has already been compromised by a reduced blood volume or concurrent administration of certain CNS depressant drugs (e.g., phenothiazines or general anesthetics) [*see Drug Interactions (7.4)*]. Monitor these patients for signs of hypotension after initiating HYCOFENIX.

In patients with circulatory shock, HYCOFENIX may cause vasodilation that can further reduce cardiac output and blood pressure. Avoid the use of HYCOFENIX in patients with circulatory shock.

5.14 Neonatal Opioid Withdrawal Syndrome

HYCOFENIX is not recommended for use in pregnant women. Prolonged use of HYCOFENIX during pregnancy can result in withdrawal in the neonate. Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. Observe newborns for signs of neonatal opioid withdrawal syndrome and manage accordingly. Advise pregnant women using opioids for a prolonged period of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available. [*see Use in Specific Populations (8.1), Patient Counseling Information (17)*]

5.15 Adrenal Insufficiency

Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use. Presentation of adrenal insufficiency may include non-specific symptoms and signs including nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. If adrenal insufficiency is suspected, confirm the diagnosis with diagnostic testing as soon as possible. If adrenal insufficiency is diagnosed, treat with physiologic replacement doses of corticosteroids. Wean the patient off of the opioid to allow adrenal function to recover and continue corticosteroid treatment until adrenal function recovers. Other opioids may be tried as some cases reported use of a different opioid without recurrence of adrenal insufficiency. The information available does not identify any particular opioids as being more likely to be associated with adrenal insufficiency.

5.16 Drug/Laboratory Test Interactions

Because opioid agonists may increase biliary tract pressure, with resultant increase in plasma amylase or lipase levels, determination of these enzyme levels may be unreliable for 24 hours after administration of a dose of HYCOFENIX.

6 ADVERSE REACTIONS

The following serious adverse reactions are described, or described in greater detail, in other sections:

- Addiction, abuse, and misuse [see *Warnings and Precautions (5.1)*, *Drug Abuse and Dependence (9.3)*]
- Life-threatening respiratory depression [see *Warnings and Precautions (5.2, 5.3, 5.4, 5.8)*, *Overdosage (10)*]
- Accidental overdose and death due to medication errors [see *Warnings and Precautions (5.5)*]
- Decreased mental alertness with impaired mental and/or physical abilities [see *Warnings and Precautions (5.6)*]
- Interactions with benzodiazepines and other CNS depressants [see *Warnings and Precautions (5.8)*, *Drug Interactions (7.1, 7.4)*]
- Paralytic ileus, gastrointestinal adverse reactions [see *Warnings and Precautions (5.9)*]
- Increased intracranial pressure [see *Warnings and Precautions (5.10)*]
- Obscured clinical course in patients with head injuries [see *Warnings and Precautions (5.10)*]
- Cardiovascular and central nervous system effects [see *Warnings and Precautions (5.11)*]
- Seizures [see *Warnings and Precautions (5.12)*]
- Severe hypotension [see *Warnings and Precautions (5.13)*]
- Neonatal Opioid Withdrawal Syndrome [see *Warnings and Precautions (5.14)*]
- Adrenal insufficiency [see *Warnings and Precautions (5.15)*]

The following adverse reactions have been identified during clinical studies, in the literature, or during post-approval use of hydrocodone, pseudoephedrine, and/or guaifenesin. Because these reactions may be reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The most common adverse reactions to HYCOFENIX include: Sedation (somnolence, mental clouding, lethargy), impaired mental and physical performance, lightheadedness, dizziness, headache, dry mouth, nausea, vomiting, constipation, tachycardia, arrhythmias including premature ventricular contractions, CNS stimulation including anxiety, restlessness, nervousness, tremor, and irritability.

Other reactions include:

Anaphylaxis: Anaphylaxis has been reported with hydrocodone, one of the ingredients in HYCOFENIX.

Body as a whole: Coma, death, fatigue, falling injuries, lethargy, weakness, hyperthermia.

Cardiovascular: Peripheral edema, increased blood pressure, decreased blood pressure, tachycardia, chest pain, palpitation, syncope, orthostatic hypotension, prolonged QT interval, hot flush, atrial fibrillation, myocardial infarction.

Central Nervous System: Ataxia, facial dyskinesia, insomnia, increased intracranial pressure, migraine, seizure, tremor.

Dermatologic: Flushing, hyperhidrosis, pruritus, rash. Cases of severe skin reactions such as acute generalized exanthematous pustulosis (AGEP) have been reported with pseudoephedrine-containing products.

Endocrine/Metabolic: Cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of opioids with serotonergic drugs. Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use. Cases of androgen deficiency have occurred with chronic use of opioids [see *Clinical Pharmacology (12.2)*].

Gastrointestinal: Abdominal pain, bowel obstruction, decreased appetite, diarrhea, difficulty swallowing, GERD, indigestion, pancreatitis, paralytic ileus, biliary tract spasm (spasm of the sphincter of Oddi), dysgeusia, ischemic colitis.

Genitourinary: Urinary tract infection, ureteral spasm, spasm of vesicle sphincters, urinary retention.

Laboratory: Increases in serum amylase.

Musculoskeletal: Arthralgia, backache, muscle spasm, weakness.

Ophthalmic: Miosis (constricted pupils), visual disturbances, mydriasis (dilated pupils).

Psychiatric: Agitation, anxiety, confusion, fear, dysphoria, depression, hyperactivity, confusion, hallucinations, hyperexcitability.

Reproductive: Hypogonadism, infertility.

Respiratory: Bronchitis, cough, dyspnea, nasal congestion, nasopharyngitis, respiratory depression, sinusitis, upper respiratory tract infection.

Other: Drug abuse, drug dependence, opioid withdrawal syndrome.

Hypoglycemia: Cases of hypoglycemia have been reported in patients taking opioids. Most reports were in patients with at least one predisposing risk factor (e.g., diabetes).

Opioid-induced esophageal dysfunction (OIED): Cases of OIED have been reported in patients taking opioids, and may occur more frequently in patients taking higher doses of opioids, and/or in patients taking opioids longer term.

7 DRUG INTERACTIONS

No specific drug interaction studies have been conducted with HYCOFENIX.

7.1 Alcohol

Concomitant use of alcohol with HYCOFENIX can result in an increase of hydrocodone plasma levels and potentially fatal overdose of hydrocodone. Instruct patients not to consume alcoholic beverages or use prescription or nonprescription products containing alcohol while on HYCOFENIX therapy [see *Warnings and Precautions (5.8)*, *Clinical Pharmacology (12.3)*].

7.2 Inhibitors of CYP3A4 and CYP2D6

The concomitant use of HYCOFENIX and CYP3A4 inhibitors, such as macrolide antibiotics (e.g., erythromycin), azole-antifungal agents (e.g. ketoconazole), or protease inhibitors (e.g., ritonavir), can increase the plasma concentration of hydrocodone, resulting in increased or prolonged opioid effects. These effects could be more pronounced with concomitant use of HYCOFENIX and CYP2D6 and CYP3A4 inhibitors, particularly when an inhibitor is added after a stable dose of HYCOFENIX is achieved [see *Warnings and Precautions (5.7)*]. After stopping a CYP3A4 inhibitor, as the effects of the inhibitor decline, the hydrocodone plasma concentration will decrease [see *Clinical Pharmacology (12.3)*], resulting in decreased opioid efficacy or a withdrawal syndrome in patients who had developed physical dependence to hydrocodone.

Avoid the use of HYCOFENIX while taking a CYP3A4 or CYP2D6 inhibitor. If concomitant use is necessary, monitor patients for respiratory depression and sedation at frequent intervals.

7.3 CYP3A4 Inducers

The concomitant use of HYCOFENIX and CYP3A4 inducers such as rifampin, carbamazepine, or phenytoin, can decrease the plasma concentration of hydrocodone [see *Clinical Pharmacology (12.3)*], resulting in decreased efficacy or onset of a withdrawal syndrome in patients who have developed physical dependence to hydrocodone [see *Warnings and Precautions (5.7)*]. After stopping a CYP3A4 inducer, as the effects of the inducer decline, the hydrocodone plasma concentration will increase [see *Clinical Pharmacology (12.3)*], which could increase or prolong both the therapeutic effects and adverse reactions, and may cause serious respiratory depression.

Avoid the use of HYCOFENIX in patients who are taking CYP3A4 inducers. If concomitant use of a CYP3A4 inducer is necessary, follow the patient for reduced efficacy.

7.4 Benzodiazepines, and Other CNS Depressants

Due to additive pharmacologic effect, the concomitant use of benzodiazepines or other CNS depressants, including alcohol, other sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics,

antipsychotics, gabapentinoids (gabapentin or pregabalin), and other opioids, can increase the risk of hypotension, respiratory depression, profound sedation, coma, and death. Avoid the use of HYCOFENIX in patients who are taking benzodiazepines, gabapentinoids (gabapentin or pregabalin), or other CNS depressants [see *Warnings and Precautions* (5.8)], and instruct patients to avoid consumption of alcohol while on HYCOFENIX [see *Drug Interactions* (7.1), *Patient Counseling Information* (17)].

7.5 Serotonergic Drugs

The concomitant use of opioids with other drugs that affect the serotonergic neurotransmitter system has resulted in serotonin syndrome. If concomitant use is warranted, carefully observe the patient, particularly during treatment initiation. Discontinue HYCOFENIX if serotonin syndrome is suspected.

7.6 Monoamine Oxidase Inhibitors (MAOIs)

Avoid the use of HYCOFENIX in patients who are taking monoamine oxidase inhibitors (MAOIs) or have taken MAOIs within 14 days. The use of MAOIs or tricyclic antidepressants with hydrocodone, one of the active ingredients in HYCOFENIX, may increase the effect of either the antidepressant or hydrocodone. MAOI interactions with opioids may manifest as serotonin syndrome or opioid toxicity (e.g., respiratory depression, coma). An increase in blood pressure or hypertensive crisis may also occur when pseudoephedrine containing preparations are used with MAOIs.

7.7 Muscle Relaxants

Hydrocodone may enhance the neuromuscular blocking action of skeletal muscle relaxants (e.g., cyclobenzaprine, metaxalone) and produce an increased degree of respiratory depression. Avoid the use of HYCOFENIX in patients taking muscle relaxants. If concomitant use is necessary, monitor patients for signs of respiratory depression that may be greater than otherwise expected.

7.8 Diuretics

Opioids can reduce the efficacy of diuretics by inducing the release of antidiuretic hormone. Monitor patients for signs of diminished diuresis and/or effects on blood pressure and increase the dosage of the diuretic as needed.

7.9 Anticholinergic Drugs

The concomitant use of anticholinergic drugs with HYCOFENIX may increase risk of urinary retention and/or severe constipation, which may lead to paralytic ileus [see *Warnings and Precautions* (5.9)]. Monitor patients for signs of urinary retention or reduced gastric motility when HYCOFENIX is used concomitantly with anticholinergic drugs.

7.10 Antihypertensive drugs

Due to the antagonistic pharmacologic effects of pseudoephedrine, one of the active ingredients in HYCOFENIX, the concomitant use of HYCOFENIX with antihypertensive drugs which interfere with sympathetic activity (e.g., methyldopa, mecamylamine, and reserpine) may reduce their antihypertensive effects. Use HYCOFENIX with caution in patients who are taking antihypertensive drugs.

7.11 Digitalis

Increased ectopic pacemaker activity can occur when pseudoephedrine is used concomitantly with digitalis. Use HYCOFENIX with caution in patients who are taking digitalis.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

HYCOFENIX is not recommended for use in pregnant women, including during or immediately prior to labor. Prolonged use of opioids during pregnancy may cause neonatal opioid withdrawal syndrome [see *Warnings and Precautions* (5.14), *Clinical Considerations*].

There are no available data with HYCOFENIX use in pregnant women to inform a drug-associated risk for adverse developmental outcomes. Published studies with hydrocodone have reported inconsistent findings and have important methodological limitations (*see Data*).

Reproductive toxicity studies have not been conducted with HYCOFENIX; however, studies are available with individual active ingredients or related active ingredients (*see Data*).

In animal reproduction studies, hydrocodone administered by the subcutaneous route to pregnant hamsters during the period of organogenesis produced a teratogenic effect at a dose approximately 70 times the maximum recommended human dose (MRHD) (*see Data*).

Guaifenesin administered by the oral route to pregnant rats during the period of organogenesis was embryolethal at a dose approximately 2 times the MRHD and produced teratogenic effects at a dose approximately 3 times the MRHD (*see Data*).

Based on the animal data, advise pregnant women of the potential risk to a fetus.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Clinical Considerations

Fetal/Neonatal Adverse Reactions

Prolonged use of opioid analgesics during pregnancy for medical or nonmedical purposes can result in physical dependence in the neonate and neonatal opioid withdrawal syndrome shortly after birth.

Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high pitched cry, tremor, vomiting, diarrhea and failure to gain weight. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by the newborn. Observe newborns for symptoms of neonatal opioid withdrawal syndrome and manage accordingly [*see Warnings and Precautions (5.14)*].

Maternal use of pseudoephedrine can cause fetal tachycardia.

Labor or Delivery

Opioids cross the placenta and may produce respiratory depression and psycho-physiologic effects in neonates. An opioid overdose reversal agent, such as naloxone or nalmefene, must be available for reversal of opioid-induced respiratory depression in the neonate. Opioids, including HYCOFENIX, can prolong labor through actions which temporarily reduce the strength, duration, and frequency of uterine contractions. However, this effect is not consistent and may be offset by an increased rate of cervical dilation, which tends to shorten labor. Monitor neonates exposed to opioids during labor for signs of excess sedation and respiratory depression.

Data

Human Data

Hydrocodone

A limited number of pregnancies have been reported in published observational studies and postmarketing reports describing hydrocodone use during pregnancy. However, these data cannot definitely establish or exclude any drug-associated risk during pregnancy. Methodological limitations of these observational studies include small sample size and lack of details regarding dose, duration and timing of exposure.

Pseudoephedrine

The majority of studies examining the use of pseudoephedrine in pregnancy did not find an association with an increased risk of congenital anomalies. Some studies reported an association with an increased risk of gastroschisis. However, several similar studies did not find a statistically significant association.

Methodological limitations of these studies included small sample size, recall bias and lack of information regarding dose and timing of exposure.

Animal Data

Reproductive toxicity studies have not been conducted with HYCOFENIX; however, studies are available with individual active ingredients or related active ingredients.

Hydrocodone

In an embryofetal development study in pregnant hamsters dosed on gestation day 8 during the period of organogenesis, hydrocodone induced cranioschisis, a malformation, at approximately 70 times the MRHD (on a mg/m² basis with a maternal subcutaneous dose of 102 mg/kg). Reproductive toxicology studies were also conducted with codeine, an opiate related to hydrocodone. In an embryofetal development study in pregnant rats dosed throughout the period of organogenesis, codeine increased resorptions and decreased fetal weights at a dose approximately 95 times the MRHD of hydrocodone (on a mg/m² basis with a maternal oral dose of codeine at 120 mg/kg/day); however, these effects occurred in the presence of maternal toxicity. In embryofetal development studies with pregnant rabbits and mice dosed throughout the period of organogenesis, codeine produced no adverse developmental effects at doses approximately 50 and 240 times, respectively, the MRHD of hydrocodone (on a mg/m² basis with maternal oral doses of codeine at 30 mg/kg/day in rabbits and 600 mg/kg/day in mice).

Pseudoephedrine

Animal studies with pseudoephedrine are not available.

Guaifenesin

In an embryofetal development study in pregnant rats dosed throughout the period of organogenesis, guaifenesin resulted in fetal death at doses approximately 2 times the MRHD (on a mg/m² basis with maternal oral doses of 350 mg/kg/day and higher). Guaifenesin also induced hemorrhagic spots and decreases in fetal weight and lengths of full body, skull, fore- and hind-limbs, and tail at doses 2 times the MRHD (on a mg/m² basis with maternal oral doses of 250 mg/kg/day and higher). Limb and tail defects, increased intercostal space, and improper development of limbs were observed at doses 3 times the MRHD (on a mg/m² basis with maternal oral doses of 500 mg/kg/day and higher).

8.2 Lactation

Risk Summary

Because of the potential for serious adverse reactions, including excess sedation, respiratory depression, and death in a breastfed infant, advise patients that breastfeeding is not recommended during treatment with HYCOFENIX.

There are no data on the presence of HYCOFENIX in human milk, the effects of HYCOFENIX on the breastfed infant, or the effects of HYCOFENIX on milk production; however, data are available with hydrocodone, pseudoephedrine, and guaifenesin.

Hydrocodone

Hydrocodone is present in breast milk. Published cases report variable concentrations of hydrocodone and hydromorphone (an active metabolite) in breast milk with administration of immediate-release hydrocodone to nursing mothers in the early post-partum period with relative infant doses of hydrocodone ranging between 1.4 and 3.7%. There are case reports of excessive sedation and respiratory depression in breastfed infants exposed to hydrocodone. No information is available on the effects of hydrocodone on milk production.

Pseudoephedrine

Pseudoephedrine is present in human milk. Pseudoephedrine has been reported to decrease milk production (*see Data*). Pseudoephedrine has been reported to cause “irritability” in a breastfed infant (*see Clinical Considerations and Data*).

Guaifenesin

No information is available on the levels of guaifenesin in breast milk or on milk production.

Clinical Considerations

Infants exposed to HYCOFENIX through breast milk should be monitored for excess sedation, respiratory depression, and irritability. Withdrawal symptoms can occur in breastfed infants when maternal administration of an opioid is stopped, or when breastfeeding is stopped.

Data

Pseudoephedrine

In a study of eight lactating women, who were 8 to 76 weeks postpartum and received a single dose of 60 mg of pseudoephedrine, the mean 24-hour milk production was reduced by 24%. In the same study, the estimated mean relative infant dose from breast milk (assuming mean milk consumption of 150 ml/kg/day and a maternal dosing regimen of 60 mg pseudoephedrine four times per day) was calculated to be 4.3% of the weight-adjusted maternal dose.

8.3 Females and Males of Reproductive Potential

Infertility

Chronic use of opioids, such as hydrocodone, a component of HYCOFENIX, may cause reduced fertility in females and males of reproductive potential. It is not known whether these effects on fertility are reversible [*see Adverse Reactions (6), Clinical Pharmacology (12.2)*].

8.4 Pediatric Use

HYCOFENIX is not indicated for use in patients younger than 18 years of age because the benefits of symptomatic treatment of cough associated with allergies or the common cold do not outweigh the risks for use of hydrocodone in these patients [*see Indications (1), Warnings and Precautions (5.3)*].

Life-threatening respiratory depression and death have occurred in children who received hydrocodone [*see Warnings and Precautions (5.2)*]. Because of the risk of life-threatening respiratory depression and death, HYCOFENIX is contraindicated in children less than 6 years of age [*see Contraindications (4)*].

8.5 Geriatric Use

Clinical studies have not been conducted with HYCOFENIX in geriatric populations.

Use caution when considering the use of HYCOFENIX in patients 65 years of age or older. Elderly patients may have increased sensitivity to hydrocodone; greater frequency of decreased hepatic, renal, or cardiac function; or concomitant disease or other drug therapy [*see Warnings and Precautions (5.4)*].

Respiratory depression is the chief risk for elderly patients treated with opioids, including HYCOFENIX. Respiratory depression has occurred after large initial doses of opioids were administered to patients who were not opioid-tolerant or when opioids were co-administered with other agents that depress respiration [*see Warnings and Precautions (5.4, 5.8)*].

Hydrocodone is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, monitor these patients closely for respiratory depression, sedation, and hypotension.

8.6 Renal Impairment

The pharmacokinetics of HYCOFENIX has not been characterized in patients with renal impairment. Patients with renal impairment may have higher plasma concentrations than those with normal function [see *Clinical Pharmacology* (12.3)]. Pseudoephedrine is primarily excreted unchanged in the urine. Therefore, pseudoephedrine may accumulate in patients with renal impairment. HYCOFENIX should be used with caution in patients with severe impairment of renal function, and patients should be monitored closely for signs of hydrocodone toxicity (respiratory depression, sedation, and hypotension) and pseudoephedrine toxicity.

8.7 Hepatic Impairment

The pharmacokinetics of HYCOFENIX has not been characterized in patients with hepatic impairment. Patients with severe hepatic impairment may have higher plasma concentrations than those with normal hepatic function [see *Clinical Pharmacology* (12.3)]. Therefore, HYCOFENIX should be used with caution in patients with severe impairment of hepatic function, and patients should be monitored closely for respiratory depression, sedation, and hypotension.

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

HYCOFENIX contains hydrocodone, a Schedule II controlled substance.

9.2 Abuse

Hydrocodone

HYCOFENIX contains hydrocodone, a substance with a high potential for abuse similar to other opioids including morphine and codeine. HYCOFENIX can be abused and is subject to misuse, addiction, and criminal diversion [see *Warnings and Precautions* (5.1)].

All patients treated with opioids require careful monitoring for signs of abuse and addiction, since use of opioid analgesic and antitussive products carries the risk of addiction even under appropriate medical use.

Prescription drug abuse is the intentional non-therapeutic use of a prescription drug, even once, for its rewarding psychological or physiological effects.

Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that develop after repeated substance use and includes: a strong desire to take the drug, difficulties in controlling its use, persisting in its use despite harmful consequences, a higher priority given to drug use than to other activities and obligations, increased tolerance, and sometimes a physical withdrawal.

“Drug-seeking” behavior is very common in persons with substance use disorders. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing, or referral, repeated “loss” of prescriptions, tampering with prescriptions, and reluctance to provide prior medical records or contact information for other treating health care provider(s). “Doctor shopping” (visiting multiple prescribers to obtain additional prescriptions) is common among drug abusers and people suffering from untreated addiction. Preoccupation with achieving adequate pain relief can be appropriate behavior in a patient with poor pain control.

Abuse and addiction are separate and distinct from physical dependence and tolerance. Health care providers should be aware that addiction may not be accompanied by concurrent tolerance and symptoms of physical dependence in all addicts. In addition, abuse of opioids can occur in the absence of true addiction.

HYCOFENIX, like other opioids, can be diverted for non-medical use into illicit channels of distribution. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests, as required by state and federal law, is strongly advised.

Proper assessment of the patient, proper prescribing practices, periodic re-evaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

Guaifenesin

Abuse of guaifenesin has been linked to the formation of kidney stones composed of the major metabolite β -(2-methoxyphenoxy) lactic acid.

Risks Specific to Abuse of HYCOFENIX

HYCOFENIX is for oral use only. Abuse of HYCOFENIX poses a risk of overdose and death. The risk is increased with concurrent use of HYCOFENIX with alcohol and other central nervous system depressants [see *Warnings and Precautions (5.8)*, *Drug Interactions (7.1, 7.4)*].

Parenteral drug abuse is commonly associated with transmission of infectious diseases such as hepatitis and HIV.

9.3 Dependence

Psychological dependence, physical dependence, and tolerance may develop upon repeated administration of opioids; therefore, HYCOFENIX should be prescribed and administered for the shortest duration that is consistent with individual patient treatment goals and patients should be reevaluated prior to refills [see *Dosage and Administration (2.3)*, *Warnings and Precautions (5.1)*].

Physical dependence, the condition in which continued administration of the drug is required to prevent the appearance of a withdrawal syndrome, assumes clinically significant proportions only after several weeks of continued oral opioid use, although some mild degree of physical dependence may develop after a few days of opioid therapy.

Do not rapidly reduce or abruptly discontinue HYCOFENIX in a physically-dependent patient. If HYCOFENIX is rapidly reduced or abruptly discontinued in a physically-dependent patient, a withdrawal syndrome may occur. Withdrawal also may be precipitated through the administration of drugs with opioid antagonist activity (e.g., naloxone, nalmefene), mixed agonist/antagonist analgesics (e.g., pentazocine, butorphanol, nalbuphine), or partial agonists (e.g., buprenorphine). Some or all of the following can characterize this syndrome: restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, myalgia, and mydriasis. Other signs and symptoms also may develop, including irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, or increased blood pressure, respiratory rate, or heart rate.

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal signs [see *Use in Specific Populations (8.1)*].

10 OVERDOSAGE

Clinical Presentation

Hydrocodone

Acute overdose with hydrocodone is characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-Stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and, in some cases, pulmonary edema, bradycardia, partial or complete airway obstruction, atypical snoring, hypotension, hypoglycemia, circulatory collapse, cardiac arrest, and death. Toxic leukoencephalopathy has been reported after opioid overdose and can present hours, days, or weeks after apparent recovery from the initial intoxication.

Hydrocodone may cause miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origin may produce similar findings). Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations [see *Clinical Pharmacology (12.2)*].

Pseudoephedrine

Overdosage with sympathomimetics such as pseudoephedrine can cause excessive CNS stimulation resulting in nervousness, anxiety, tremor, restlessness and insomnia. Other effects may include headache, tachycardia, palpitations, precordial pain, hypertension, pallor, mydriasis, nausea, vomiting, sweating, thirst, urinary retention (difficulty in micturition), muscle weakness and tenseness, giddiness, anxiety, restlessness, hyperglycemia, and insomnia. Many patients can present a toxic psychosis with delusion and hallucinations. Severe overdosage may cause tachypnea or hyperpnea, hallucinations, convulsions, delirium, or coma, but in some individuals there may be CNS depression with somnolence, stupor, respiratory depression, or respiratory failure. Arrhythmias (including ventricular fibrillation) may lead to hypotension and circulatory collapse. Severe hypokalemia can occur, probably due to compartmental shift rather than depletion of potassium.

Guaifenesin

Overdosage with guaifenesin can cause depression of the central nervous system. While present in polypharmacy overdoses, one case of overdose with only significant levels of guaifenesin has been reported. Symptoms included slurred speech, shallow respirations, reduced heart rate with rhythm sinus bradycardia, followed by asystole.

Treatment of Overdose

Treatment of overdosage is driven by the overall clinical presentation, and consists of discontinuation of HYCOFENIX together with institution of appropriate therapy. Give primary attention to the reestablishment of adequate respiratory exchange through provision of a patent and protected airway and the institution of assisted or controlled ventilation. Employ other supportive measures (including oxygen and vasopressors) in the management of circulatory shock and pulmonary edema as indicated. Cardiac arrest or arrhythmias will require advanced life-support techniques. Gastric emptying may be useful in removing unabsorbed drug.

For clinically significant respiratory or circulatory depression secondary to hydrocodone overdose, administer an opioid overdose reversal agent such as naloxone or nalmefene. An opioid overdose reversal agent should not be administered in the absence of clinically significant respiratory depression. Because the duration of opioid reversal is expected to be less than the duration of action of hydrocodone in HYCOFENIX, carefully monitor the patient until spontaneous respiration is reliably reestablished. If the response to an opioid overdose reversal agent is suboptimal or only brief in nature, administer additional reversal agent as directed by the product's prescribing information.

Hemodialysis is not routinely used to enhance the elimination of hydrocodone, pseudoephedrine or guaifenesin from the body.

Adrenergic receptor blocking agents (beta-blockers), such as propranolol, may be used for the treatment of cardiac toxicity due to pseudoephedrine.

11 DESCRIPTION


HYCOFENIX (hydrocodone bitartrate, pseudoephedrine hydrochloride, and guaifenesin) oral solution contains hydrocodone, an opioid agonist; pseudoephedrine, an alpha-adrenergic agonist; and guaifenesin, an expectorant. Each 5 mL of HYCOFENIX contains 2.5 mg of hydrocodone bitartrate, 30 mg of pseudoephedrine hydrochloride, and 200 mg of guaifenesin for oral administration.

HYCOFENIX also contains the following inactive ingredients: black raspberry flavor, citric acid, D&C Red #33, FD&C Blue #1, glycerin, methylparaben, polyethylene glycol propylparaben, purified water, saccharin sodium, sodium citrate, and sorbitol.

Hydrocodone Bitartrate


The chemical name for hydrocodone bitartrate is morphinan-6-one, 4,5-epoxy-3-methoxy-17-methyl-, (5 α)-, [R-(R*,R*)]-2,3-dihydroxybutanedioate (1:1), hydrate (2:5). It is also known as 4,5 α -Epoxy-3-methoxy-17-

methylmorphinan-6-one tartrate (1:1) hydrate (2:5). It occurs as a fine white crystal or crystalline powder, which is derived from the opium alkaloid, thebaine, and it has the following chemical structure:




Pseudoephedrine Hydrochloride

The chemical name for pseudoephedrine hydrochloride is benzenemethanol, α -[1-(methylamino)ethyl]-, [S-(R*,R*)] hydrochloride and has the following chemical structure:



Guaifenesin

The chemical name for guaifenesin is 3-(2-methoxyphenoxy)-1,2-propanediol. It occurs as a white powder, and it has the following chemical structure:



12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Hydrocodone

Hydrocodone is an opioid agonist with relative selectivity for the mu-opioid receptor, although it can interact with other opioid receptors at higher doses. The precise mechanism of action of hydrocodone and other opiates is not known; however, hydrocodone is believed to act centrally on the cough center. In excessive doses, hydrocodone will depress respiration.

Pseudoephedrine

Pseudoephedrine is a sympathomimetic amine that exerts a decongestant action on the nasal mucosa via alpha adrenergic activity. Pseudoephedrine produces peripheral effects similar to those of ephedrine and central effects similar to, but less intense than, amphetamines. It has the potential for excitatory side effects.

Guaifenesin

Guaifenesin is an expectorant, the action of which promotes or facilitates the removal of secretions from the respiratory tract. The precise mechanism of action of guaifenesin is not known; however, it is thought to act as an expectorant by increasing the volume and reducing the viscosity of secretions in the trachea and bronchi. In turn, this may increase the efficiency of the cough reflex and facilitate removal of the secretions.

12.2 Pharmacodynamics

Hydrocodone

Effects on the Central Nervous System

Hydrocodone produces respiratory depression by direct action on brain stem respiratory centers. The respiratory depression involves a reduction in the responsiveness of the brain stem respiratory centers to both increases in carbon dioxide tension and to electrical stimulation.

Hydrocodone causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origins may produce similar findings). Marked mydriasis rather than miosis may be seen due to hypoxia in overdose situations.

Effects on the Gastrointestinal Tract and Other Smooth Muscle

Hydrocodone causes a reduction in motility associated with an increase in smooth muscle tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone may be increased to the point of spasm resulting in constipation. Other opioid-induced effects may include a reduction in biliary and pancreatic secretions, spasm of sphincter of Oddi, transient elevations in serum amylase, and opioid-induced esophageal dysfunction (OIED).

Effects on the Cardiovascular System

Hydrocodone produces peripheral vasodilation which may result in orthostatic hypotension or syncope. Manifestations of histamine release and/or peripheral vasodilation may include pruritus, flushing, red eyes and sweating and/or orthostatic hypotension.

Effects on the Endocrine System

Opioids inhibit the secretion of adrenocorticotrophic hormone (ACTH), cortisol, and luteinizing hormone (LH) in humans [see *Adverse Reactions* (6)]. They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon.

Chronic use of opioids may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The causal role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date [see *Adverse Reactions* (6)].

Effects on the Immune System

Opioids have been shown to have a variety of effects on components of the immune system in in vitro and animal models. The clinical significance of these findings is unknown. Overall, the effects of opioids appear to be modestly immunosuppressive.

Concentration–Adverse Reaction Relationships

There is a relationship between increasing hydrocodone plasma concentration and increasing frequency of dose-related opioid adverse reactions such as nausea, vomiting, CNS effects, and respiratory depression. In opioid-tolerant patients, the situation may be altered by the development of tolerance to opioid-related adverse reactions.

Pseudoephedrine

Interaction of pseudoephedrine with alpha-1 adrenergic receptors on vascular smooth muscle cells causes activation of the cells and results in vasoconstriction.

12.3 Pharmacokinetics

Absorption

Following a single 10 mL oral dose administration of 5 mg hydrocodone bitartrate, 60 mg pseudoephedrine hydrochloride, and 400 mg guaifenesin administered to 37 healthy adults, the geometric mean Cmax and AUC0-inf for hydrocodone were 9.0 ng/mL and 61.2 ng•hr/mL, respectively. The median time to maximum concentration for hydrocodone was about 1.67 hours. Food has no significant effect on the extent of absorption of hydrocodone.

Pseudoephedrine is rapidly and almost completely absorbed from the gastrointestinal tract. Following a single 10 mL oral dose administration of 5 mg hydrocodone bitartrate, 60 mg pseudoephedrine hydrochloride, and 400 mg guaifenesin administered to 37 healthy adults, the geometric mean Cmax and AUC0-inf for pseudoephedrine were 0.19 mcg/mL and 1.9 mcg•hr/mL, respectively. The median time to maximum concentration for pseudoephedrine was about 2.5 hours post administration. Food has no significant effect on the extent of absorption of pseudoephedrine.

Following a single 10 mL oral dose administration of 5 mg hydrocodone bitartrate, 60 mg pseudoephedrine hydrochloride, and 400 mg guaifenesin administered to 36 healthy adults, the geometric mean Cmax and AUC0-inf for guaifenesin were 2.0 mcg/mL and 2.6 mcg•hr/mL, respectively. The median time to maximum concentration for guaifenesin was about 25 minutes. The effect of food on guaifenesin systemic exposure is not considered to be clinically meaningful.

Distribution

Although the extent of protein binding of hydrocodone in human plasma has not been definitively determined, structural similarities to related opioid analgesics suggest that hydrocodone is not extensively protein bound. As most agents in the 5-ring morphinan group of semi-synthetic opioids bind plasma protein to a similar degree (range 19% [hydromorphone] to 45% [oxycodone]), hydrocodone is expected to fall within this range.

Pseudoephedrine hydrochloride is extensively distributed into extravascular sites. The apparent volume of distribution (V/F) of pseudoephedrine ranged between 2.6 and 3.5 L/kg.

Elimination

Metabolism

Hydrocodone exhibits a complex pattern of metabolism, including N-demethylation, O-demethylation, and 6keto reduction to the corresponding 6- α -and 6- β -hydroxy metabolites. CYP3A4 mediated N-demethylation to norhydrocodone is the primary metabolic pathway of hydrocodone with a lower contribution from CYP2D6 mediated O-demethylation to hydromorphone. Hydromorphone is formed from the O-demethylation of hydrocodone and may contribute to the total analgesic effect of hydrocodone. Therefore, the formation of these and related metabolites can, in theory, be affected by other drugs [see *Drug Interactions* (7)]. Published in vitro studies have shown that N-demethylation of hydrocodone to form norhydrocodone can be attributed to CYP3A4 while O-demethylation of hydrocodone to hydromorphone is predominantly catalyzed by CYP2D6 and to a lesser extent by an unknown low affinity CYP enzyme.

Excretion

Hydrocodone and its metabolites are eliminated primarily in the kidneys. The mean plasma half-life of hydrocodone is approximately 4 hours.

About 43-96% of an administered dose of pseudoephedrine is excreted unchanged in the urine. The remainder is apparently metabolized in the liver to inactive compounds by N-demethylation, parahydroxylation and oxidative deamination. Pseudoephedrine has been shown to have a mean elimination half-life of 4-6 hours which is dependent on urine pH. The elimination half-life is decreased at urine pH lower than 6 and may be increased at urine pH higher than 8.

The mean plasma half-life of guaifenesin is approximately 1 hour.

Specific Populations

Renal Impairment

No data are available on the pharmacokinetics of pseudoephedrine in renally-impaired subjects. Pseudoephedrine is primarily excreted unchanged in the urine. A decrease in renal function is, therefore, likely to decrease the clearance of pseudoephedrine, prolonging the half-life and resulting in accumulation. Therefore, pseudoephedrine may accumulate in patients with renal impairment.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity, mutagenicity, and fertility studies have not been conducted with HYCOFENIX; however, published information is available for the individual active ingredients or related active ingredients.

Hydrocodone

Carcinogenicity studies were conducted with codeine, an opiate related to hydrocodone. Two-year studies in F344/N rats and B6C3F1 mice were conducted to assess the carcinogenic potential of codeine. No evidence of tumorigenicity was observed in male and female rats at codeine dietary doses up to 70 and 80 mg/kg/day (approximately equivalent to 55 and 65 times the MRHD of hydrocodone on a mg/m² basis, respectively). No evidence of tumorigenicity was observed in male and female mice at codeine dietary doses up to 400 mg/kg/day (approximately equivalent to 160 times the MRHD of hydrocodone on a mg/m² basis).

Mutagenicity studies with hydrocodone have not been conducted.

Fertility studies with hydrocodone have not been conducted.

Pseudoephedrine

Carcinogenicity studies were conducted with ephedrine sulfate, a structurally related drug. Two-year studies in F344/N rats and B6C3F1 mice were conducted to assess the carcinogenic potential of ephedrine sulfate. No evidence of tumorigenicity was observed in male and female rats at ephedrine sulfate dietary doses up to 9 and 11 mg/kg/day (approximately equivalent to 0.4 and 0.5 times the MRHD of pseudoephedrine on a mg/m² basis, respectively). No evidence of tumorigenicity was observed in male and female mice at ephedrine sulfate dietary doses up to 29 and 25 mg/kg/day (approximately equivalent to 0.7 and 0.6 times the MRHD of pseudoephedrine on a mg/m² basis, respectively).

Mutagenicity studies with pseudoephedrine have not been conducted.

Fertility studies with pseudoephedrine have not been conducted.

Guaifenesin

Carcinogenicity, mutagenicity, and fertility studies with guaifenesin have not been conducted.

16 HOW SUPPLIED/STORAGE AND HANDLING

HYCOFENIX (hydrocodone bitartrate, pseudoephedrine hydrochloride, and guaifenesin) oral solution is supplied as a violet-colored, black raspberry flavored liquid containing 2.5 mg hydrocodone bitartrate, 30 mg pseudoephedrine hydrochloride, and 200 mg guaifenesin in each 5 mL. It is available in:

White HDPE bottles of 16 fl. oz. (473 mL): NDC XXXXX-XXX-XX

White HDPE bottles of 4 fl. oz. (118 mL): NDC XXXXX-XXX-XX

Store solution at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]

Dispense in a tight, light-resistant container, as defined in the USP, with a child-resistant closure.

Ensure that patients have an oral dosing dispenser that measures the appropriate volume in milliliters. Counsel patients on how to utilize an oral dosing dispenser and correctly measure the oral suspension as prescribed.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Addiction, Abuse, and Misuse

Inform patients that the use of HYCOFENIX, even when taken as recommended, can result in addiction, abuse, and misuse, which can lead to overdose and death [*see Warnings and Precautions (5.1)*]. Instruct patients not to share HYCOFENIX with others and to take steps to protect HYCOFENIX from theft or misuse.

Important Dosing and Administration Instructions

Instruct patients how to measure and take the correct dose of HYCOFENIX. Advise patients to measure HYCOFENIX with an accurate milliliter measuring device. Patients should be informed that a household teaspoon is not an accurate measuring device and could lead to overdosage. Advise patients to ask their pharmacist to recommend an appropriate measuring device and for instructions for measuring the correct dose [*see Dosage and Administration (2.1), Warnings and Precautions (5.5)*]. Advise patients not to increase the dose or dosing frequency of HYCOFENIX because serious adverse events such as respiratory depression may occur with overdosage [*see Warnings and Precautions (5.2), Overdosage (10)*].

Life-Threatening Respiratory Depression

Inform patients of the risk of life-threatening respiratory depression, including information that the risk is greatest when starting HYCOFENIX and that it can occur even at recommended dosages [*see Warnings and Precautions (5.2)*]. Advise patients how to recognize respiratory depression and to seek medical attention if breathing difficulties develop.

Accidental Ingestion

Inform patients that accidental ingestion, especially by children, may result in respiratory depression or death [*see Warnings and Precautions (5.2)*]. Instruct patients to take steps to store HYCOFENIX securely and to properly dispose of unused HYCOFENIX in accordance with the local state guidelines and/or regulations.

Activities Requiring Mental Alertness

Advise patients to avoid engaging in hazardous tasks that require mental alertness and motor coordination such as operating machinery or driving a motor vehicle as HYCOFENIX may produce marked drowsiness [*see Warnings and Precautions (5.6)*].

Interactions with Benzodiazepines and Other Central Nervous System Depressants, Including Alcohol

Inform patients and caregivers that potentially fatal additive effects may occur if HYCOFENIX is used with benzodiazepines or other CNS depressants, including alcohol (e.g., non-benzodiazepine sedative/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, gabapentinoids [gabapentin or pregabalin], and other opioids). Advise patients to avoid concomitant use of HYCOFENIX with benzodiazepines or other CNS depressants and instruct patients not to consume alcoholic beverages, as well as prescription and over-the-counter products that contain alcohol, during treatment with HYCOFENIX [*see Warnings and Precautions (5.8), Drug Interactions (7.1, 7.4)*].

Constipation

Advise patients of the potential for severe constipation [*see Warnings and Precautions (5.9), Adverse Reactions (6)*].

Cardiovascular and CNS Effects

Inform patients that the pseudoephedrine contained in HYCOFENIX can produce cardiovascular and central nervous system effects in some patients such as, insomnia, dizziness, weakness, tremor, transient elevations in blood pressure, or arrhythmias.

Anaphylaxis

Inform patients that anaphylaxis has been reported with ingredients contained in HYCOFENIX. Advise

patients how to recognize such a reaction and when to seek medical attention [see *Contraindications (4)*, *Adverse Reactions (6)*].

MAOI Interaction

Inform patients not to take HYCOFENIX while using or within 14 days of stopping any drugs that inhibit monoamine oxidase. Patients should not start MAOIs while taking HYCOFENIX [see *Drug Interactions (7.6)*].

Hypotension

Inform patients that HYCOFENIX may cause orthostatic hypotension and syncope. Instruct patients how to recognize symptoms of low blood pressure and how to reduce the risk of serious consequences should hypotension occur (e.g., sit or lie down, carefully rise from a sitting or lying position) [see *Warnings and Precautions (5.13)*].

Pregnancy

Advise patients that use of HYCOFENIX is not recommended during pregnancy [see *Use in Specific Populations (8.1)*].

Neonatal Opioid Withdrawal Syndrome

Inform female patients of reproductive potential that use of HYCOFENIX during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated [see *Warnings and Precautions (5.14)*, *Use in Specific Populations (8.1)*].

Embryo-Fetal Toxicity

Inform female patients of reproductive potential that HYCOFENIX can cause fetal harm and to inform their healthcare provider of a known or suspected pregnancy [see *Use in Specific Populations (8.1)*].

Lactation

Advise women that breastfeeding is not recommended during treatment with HYCOFENIX [see *Use in Specific Populations (8.2)*].

Infertility

Inform patients that chronic use of opioids, such as hydrocodone, a component of HYCOFENIX, may cause reduced fertility. It is not known whether these effects on fertility are reversible [see *Use in Specific Populations (8.3)*].

Adrenal Insufficiency

Inform patients that HYCOFENIX could cause adrenal insufficiency, a potentially life-threatening condition. Adrenal insufficiency may present with non-specific symptoms and signs such as nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. Advise patients to seek medical attention if they experience a constellation of these symptoms [see *Warnings and Precautions (5.15)*].

Serotonin Syndrome

Inform patients that HYCOFENIX could cause a rare but potentially life-threatening condition resulting from concomitant administration of serotonergic drugs. Warn patients of the symptoms of serotonin syndrome and to seek medical attention right away if symptoms develop. Instruct patients to inform their physicians if they are taking, or plan to take serotonergic medications. [see *Adverse Reactions (6)*, *Drug Interactions (7.6)*].

Disposal of Unused HYCOFENIX

Advise patients to properly dispose of unused HYCOFENIX. Advise patients to throw the drug in the household trash following these steps. 1) Remove them from their original containers and mix them with an undesirable substance, such as used coffee grounds or kitty litter (this makes the drug less appealing to children and pets, and unrecognizable to people who may intentionally go through the trash seeking drugs). 2) Place the

mixture in a sealable bag, empty can, or other container to prevent the drug from leaking or breaking out of a garbage bag, or to dispose of in accordance with local state guidelines and/or regulations.

Manufactured for:
Chartwell RX Sciences, LLC.
Congers, NY 10920

LXXXXX

MEDICATION GUIDE
HYCOFENIX® (Hye-koh-fee-niks)
(hydrocodone bitartrate, pseudoephedrine hydrochloride, and guaifenesin) oral
solution, C-II

What is the most important information I should know about HYCOFENIX? HYCOFENIX is not for children under 18 years of age.

HYCOFENIX can cause serious side effects, including:

- **Addiction, abuse and misuse.** Taking HYCOFENIX or other medicines that contain an opioid can cause addiction, abuse and misuse, which can lead to overdose and death. This can happen even if you take HYCOFENIX exactly as prescribed by your healthcare provider. Your risk of addiction, abuse, and misuse is increased if you or a family member has a history of drug or alcohol abuse or addiction, or mental health problems.
 - **Do not** share your HYCOFENIX with other people.
 - Keep HYCOFENIX in a safe place away from children.
- **Life-threatening breathing problems (respiratory depression).** HYCOFENIX can cause breathing problems (respiratory depression) that can happen at any time during treatment and can lead to death. Your risk of breathing problems is greatest when you first start taking HYCOFENIX, are taking other medicines that can cause breathing problems, have certain lung problems, are elderly, or have certain other health problems. **Children are at higher risk for respiratory depression.**

Breathing problems can happen even if you take HYCOFENIX exactly as prescribed by your healthcare provider.

Call your healthcare provider or get emergency medical help right away if anyone taking HYCOFENIX has any of the symptoms below:

- increased sleepiness
- shallow breathing
- confusion
- limpness
- difficulty breathing

Keep HYCOFENIX in a safe place away from children. Accidental use of even 1 dose of HYCOFENIX, especially by a child, is a medical emergency and can cause breathing problems (respiratory depression) which can lead to death. If a child accidentally takes HYCOFENIX, get emergency medical help right away.

- **Overdose and death due to medicine dosing errors.** Overdose and death can happen if you measure the wrong dose of HYCOFENIX. Always use an accurate milliliter (mL) measuring device to measure the correct amount of HYCOFENIX. **Do not** use a household teaspoon to measure your medicine. You may accidentally take too much. You can ask your pharmacist for the measuring device and how to measure the correct dose.
- **Breathing problems (respiratory depression) that can lead to death and opioid withdrawal** can happen if you start taking or stop taking other medicines while taking HYCOFENIX, including:
 - certain antibiotics
 - certain medicines to treat a fungal infection
 - certain medicines to treat Human Immunodeficiency Virus (HIV)-1 infection, Acquired Immune Deficiency Syndrome (AIDS), or Hepatitis C
 - rifampin
 - carbamazepine
 - phenytoin

Tell your healthcare provider if you take any of these medicines. Ask your healthcare provider or pharmacist if you are not sure if your medicine is one listed above.

- **Severe drowsiness, breathing problems (respiratory depression), coma, and death** can happen in people who take HYCOFENIX with benzodiazepines, gabapentinoids (gabapentin or pregabalin), or other central nervous system depressants, including alcohol.
 - **Do not** take benzodiazepines or any medicine that can cause drowsiness or sleepiness during treatment with HYCOFENIX.
 - **Do not** drink alcohol or take prescription or over-the-counter medicines that contain alcohol during treatment with HYCOFENIX.
- **Opioid withdrawal in a newborn.** Use of HYCOFENIX during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated. You should not take HYCOFENIX if you are pregnant. Tell your healthcare provider right away if you are pregnant or think you may be pregnant.

What is HYCOFENIX?

- HYCOFENIX is a prescription medicine used in adults to treat cough, a stuffy nose (nasal congestion), and loosen mucus that you can have with a common cold. HYCOFENIX contains 3 medicines, hydrocodone, pseudoephedrine and guaifenesin. Hydrocodone is an opioid (narcotic) cough suppressant. Pseudoephedrine is a decongestant. Guaifenesin is an expectorant.
- **HYCOFENIX is a federal controlled substance (C-II) because it contains hydrocodone that can be abused or lead to dependence.** Keep HYCOFENIX in a safe place to prevent misuse and abuse. Selling or giving away HYCOFENIX may harm others, and is against the law. Tell your healthcare provider if you have abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take HYCOFENIX?

HYCOFENIX is not for children under 18 years of age. See “What is the most important information I should know about HYCOFENIX?”

Do not take HYCOFENIX if you:

- have severe breathing problems (respiratory depression) or breathing problems caused by asthma. See “[What is the most important information I should know about HYCOFENIX?](#)”
- have a blockage (obstruction) in your bowel such as a paralytic ileus.
- have a type of glaucoma called “narrow angle glaucoma”.
- have problems emptying your bladder (urinary retention).
- have severe high blood pressure or certain heart problems (severe coronary artery disease).
- are allergic to hydrocodone, pseudoephedrine, guaifenesin, or any of the ingredients in HYCOFENIX. See the end of this Medication Guide for a complete list of ingredients.

Ask your healthcare provider if you have any questions about this information.

Before you take HYCOFENIX, tell your healthcare provider about all of your medical conditions, including if you:

- have a drug addiction
- have lung or breathing problems
- have a fever and are coughing up mucus
- have had a recent head injury
- have had a brain tumor or other brain problems
- have or have had seizures
- have pain in your stomach-area (abdomen)
- have constipation or other bowel problems
- have bile duct or pancreas problems
- have glaucoma (increased pressure in your eyes)
- have prostate problems
- have problems with your urinary tract or difficulty urinating
- have kidney or liver problems
- have heart or blood vessel (cardiovascular) problems
- have adrenal gland problems
- plan to have surgery
- have low blood pressure
- are pregnant or plan to become pregnant. HYCOFENIX can harm your unborn baby. See “[What is the most important information I should know about HYCOFENIX?](#)”
- are breastfeeding or plan to breastfeed. Hydrocodone and pseudoephedrine pass into your breast milk and can cause serious side effects in your baby including increased sleepiness, irritability, breathing problems (respiratory depression), and death. You and your healthcare provider should decide if you will take HYCOFENIX or breastfeed. You should not do both. See “[What should I avoid while taking HYCOFENIX?](#)”
- plan to have children. HYCOFENIX may affect the ability to have a child in females and males (fertility problems). It is not known if these fertility problems will be reversible, even after you stop taking HYCOFENIX. Talk to your healthcare provider if this is a concern for you.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

Taking HYCOFENIX with certain other medicines can cause side effects or affect how well HYCOFENIX or the other medicines work. Do not start or stop taking other medicines without talking to your healthcare provider.

Especially tell your healthcare provider if you:

- take pain medicines such as opioids (narcotics).
- take cold or allergy medicines that contain antihistamines or cough suppressants.
- drink alcohol.
- take muscle relaxants.
- take certain medicines used to treat mood, anxiety, psychotic or thought disorders, or depression, including monoamine oxidase inhibitors (MAOIs), tricyclics, selective serotonin reuptake inhibitors (SSRIs), selective serotonin-norepinephrine reuptake inhibitors (SNRIs), or antipsychotics.
- take medicines to lower your blood pressure.
- take water pills (diuretics).
- take medicines called “anticholinergics” used to treat certain health problems including asthma, chronic obstructive pulmonary disease (COPD), or stomach problems.
- take a medicine called “digitalis” used to treat certain heart problems.

Ask your healthcare provider if you are not sure if you take one of these medicines.

How should I take HYCOFENIX?

- See “[What is the most important information I should know about HYCOFENIX?](#)”
- Take HYCOFENIX exactly as your healthcare provider tells you to take it. Do not change your dose without talking to your healthcare provider.
- Take HYCOFENIX using an accurate milliliter (mL) measuring device. If you do not have one, ask your pharmacist to give you a measuring device to help you measure the correct amount of HYCOFENIX. **Do not use a household teaspoon to measure your medicine. You may accidentally take too much.**
- **Do not** overfill the measuring device.

- Rinse your measuring device with water after each use.
- If you take too much HYCOFENIX, call your healthcare provider or go to the nearest hospital emergency room right away.
- Tell your healthcare provider if your cough does not get better within 5 days of treatment with HYCOFENIX.

What should I avoid while taking HYCOFENIX?

- Avoid driving a car or operating machinery during treatment with HYCOFENIX. HYCOFENIX can cause you to be drowsy, slow your thinking and motor skills, and affect your vision.
- **Do not** drink alcohol during treatment with HYCOFENIX. Drinking alcohol can increase your chances of having serious side effects.

Avoid the use of HYCOFENIX if you:

- are pregnant. Use of HYCOFENIX during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated. Tell your healthcare provider right away if you are pregnant or think you may be pregnant.
- are breastfeeding. Use of HYCOFENIX while breastfeeding can cause severe breathing problems (respiratory depression) in your breastfed infant that could be life-threatening.
- take a medicine called a monoamine oxidase inhibitor (MAOI). Avoid taking an MAOI within 14 days after you stop taking HYCOFENIX. Avoid starting HYCOFENIX if you stopped taking an MAOI in the last 14 days.

What are the possible side effects of HYCOFENIX? HYCOFENIX

can cause serious side effects, including:

- See "[What is the most important information I should know about HYCOFENIX?](#)"
- **Bowel problems including severe constipation or stomach pain.** See "[Who should not take HYCOFENIX?](#)"
- **Increased pressure in your head (intracranial).** Avoid the use of HYCOFENIX if you have a head injury or have been told that you have changes in the tissue of your brain (brain lesions) or increased pressure in your head.
- **Heart and blood vessel (cardiovascular) and central nervous system (CNS) effects.** Cardiovascular and CNS effects can happen in some people during treatment with HYCOFENIX, including trouble sleeping (insomnia), dizziness, weakness, tremors, temporary increases in blood pressure, abnormal heart beats (arrhythmias), seizures, and feeling faint. Call your healthcare provider right away if you have any of these symptoms.
- **Low blood pressure.** A sudden drop in blood pressure can happen in some people during treatment with HYCOFENIX and this may cause you to feel dizzy, faint, lightheaded, or weak, especially when you stand up (orthostatic hypotension). Your risk of having this problem may be increased if you take HYCOFENIX with certain other medicines that lower blood pressure. If you have any of these symptoms while taking HYCOFENIX, sit or lie down. Do not change your body position too fast. Get up slowly from sitting or lying down.
- **Adrenal gland problems.** HYCOFENIX can cause serious and life-threatening adrenal gland problems. Your healthcare provider may do blood tests to check for adrenal gland problems. Call your healthcare provider right away if you have any of these symptoms:
 - nausea
 - vomiting
 - not wanting to eat (anorexia)
 - fatigue
 - weakness
 - dizziness
 - low blood pressure

The most common side effects of HYCOFENIX include:

• sleepiness	• nausea
• confusion	• vomiting
• coordination problems	• constipation
• decrease in mental and physical performance	• fast or irregular heart beat
• lack of energy	• irritability
• lightheadedness	• anxiety
• dizziness	• restlessness
• headache	• nervousness
• dry mouth	• tremor

These are not all the possible side effects of HYCOFENIX.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store HYCOFENIX?

- Store HYCOFENIX at room temperature between 68°F to 77°F (20°C to 25°C).
- Store HYCOFENIX in a tightly closed container, in a dry, cool place away from heat or direct sunlight.
- **Keep HYCOFENIX and all medicines out of the reach of children.**

How should I store/ dispose of HYCOFENIX?

Remove unused HYCOFENIX from the container and mix it with an undesirable, non-toxic substance such as cat litter or used coffee grounds to make it less appealing to children and pets. Place the mixture in a container such as a

sealed plastic bag and throw it away in the household trash. You can also follow your state or local guidelines on how to safely throw away HYCOFENIX.

General information about the safe and effective use of HYCOFENIX.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use HYCOFENIX for a condition for which it was not prescribed. Do not give HYCOFENIX to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about HYCOFENIX that is written for health professionals.

What are the ingredients in HYCOFENIX?

Active Ingredients: hydrocodone bitartrate, pseudoephedrine hydrochloride, and guaifenesin

Inactive Ingredients: black raspberry flavor, citric acid, D&C Red #33, FD&C Blue #1, glycerin, methylparaben, polyethylene glycol propylparaben, purified water, saccharin sodium, sodium citrate, and sorbitol

Manufactured for: Chartwell RX Sciences, LLC. For more information, go to
www.ChartwellPharma.com or call 1-845-268-5000.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: 12/2025