

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use **LEVOTHYROXINE SODIUM FOR INJECTION** safely and effectively. See full prescribing information for **LEVOTHYROXINE SODIUM FOR INJECTION**.

LEVOTHYROXINE SODIUM for injection, for intravenous use
Initial U.S. Approval: 1969

WARNING: NOT FOR TREATMENT OF OBESITY OR FOR WEIGHT LOSS

Thyroid hormones, including **Levothyroxine Sodium for Injection**, should not be used for the treatment of obesity or for weight loss. (5.3)
Larger doses may produce serious or even life threatening manifestations of toxicity. (6)

INDICATIONS AND USAGE

Levothyroxine Sodium is an L-thyroxine product. Levothyroxine (T₄) Sodium for Injection is indicated for the treatment of myxedema coma. (1)

Important Limitations of Use:

The relative bioavailability of this drug has not been established. Use caution when converting patients from oral to intravenous levothyroxine. (1)

DOSAGE AND ADMINISTRATION

- An initial intravenous loading dose of Levothyroxine Sodium for Injection between 300 to 500 mcg followed by once daily intravenous maintenance doses between 50 and 100 mcg should be administered, as clinically indicated, until the patient can tolerate oral therapy. (2.1)
- Reconstitute the lyophilized Levothyroxine Sodium for Injection by aseptically adding 5 mL of 0.9% Sodium Chloride Injection, USP. Shake vial to ensure complete mixing. Reconstituted drug product is preservative free. Use immediately after reconstitution. Discard any unused portion. (2.3)
- Do not add to other intravenous fluids. (2.3)

DOSAGE FORMS AND STRENGTHS

For injection: a lyophilized powder in a single-dose amber-colored vial containing Levothyroxine Sodium available in the following strengths: (3)

- 100 mcg per vial
- 200 mcg per vial
- 500 mcg per vial

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

- Excessive bolus doses of Levothyroxine Sodium for Injection (greater than 500 mcg) are associated with cardiac complications, particularly in the elderly and in patients with an underlying cardiac condition. Initiate therapy with doses at the lower end of the recommended range. (5.1)
- Close observation of the patient following the administration of Levothyroxine Sodium for Injection is advised. (5.1)
- Levothyroxine Sodium for Injection therapy for patients with previously undiagnosed endocrine disorders, including adrenal insufficiency, hypopituitarism, and diabetes insipidus, may worsen symptoms of these endocrinopathies. (5.2)

ADVERSE REACTIONS

Excessive doses of L-thyroxine can predispose to signs and symptoms compatible with hyperthyroidism. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Many drugs affect thyroid hormone pharmacokinetics and metabolism (e.g., absorption, synthesis, secretion, catabolism, protein binding, and target tissue response) and may alter the therapeutic response to Levothyroxine Sodium for Injection. (7, 12.3)

USE IN SPECIFIC POPULATIONS

Elderly and those with underlying cardiovascular disease should receive doses at the lower end of the recommended range. (8.5)

Revised: 7/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: NOT FOR TREATMENT OF OBESITY OR FOR WEIGHT LOSS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosage
- 2.2 Dosing in the Elderly and in Patients with Cardiovascular Disease
- 2.3 Reconstitution Directions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Risk of Cardiac Complications in Elderly and in Patients with Cardiovascular Disease
- 5.2 Need for Concomitant Glucocorticoids and Monitoring for Other Diseases in Patients with Endocrine Disorders
- 5.3 Not Indicated for Treatment of Obesity

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

- 7.1 Antidiabetic Therapy
- 7.2 Oral Anticoagulants
- 7.3 Digitalis Glycosides
- 7.4 Antidepressant Therapy
- 7.5 Ketamine
- 7.6 Sympathomimetics

7.7 Drug-Laboratory Test Interactions

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use and Patients with Underlying Cardiovascular Disease

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and Pharmacology

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: NOT FOR TREATMENT OF OBESITY OR FOR WEIGHT LOSS

Thyroid hormones, including Levothyroxine Sodium for Injection, should not be used for the treatment of obesity or for weight loss. (5.3)

Larger doses may produce serious or even life threatening manifestations of toxicity. (6)

1 INDICATIONS AND USAGE

Levothyroxine Sodium for Injection is indicated for the treatment of myxedema coma.

Important Limitations of Use: The relative bioavailability between Levothyroxine Sodium for Injection and oral levothyroxine products has not been established. Caution should be used when switching patients from oral levothyroxine products to Levothyroxine Sodium for Injection as accurate dosing conversion has not been studied.

2 DOSAGE AND ADMINISTRATION

2.1 Dosage

An initial intravenous loading dose of Levothyroxine Sodium for Injection between 300 to 500 mcg, followed by once daily intravenous maintenance doses between 50 and 100 mcg, should be administered, as clinically indicated, until the patient can tolerate oral therapy. The age, general physical condition, cardiac risk factors, and clinical severity of myxedema and duration of myxedema symptoms should be considered when determining the starting and maintenance dosages of Levothyroxine Sodium for Injection.

Levothyroxine Sodium for Injection produces a gradual increase in the circulating concentrations of the hormone with an approximate half-life of 9 to 10 days in hypothyroid patients. Daily administration of Levothyroxine Sodium for Injection should be maintained until the patient is capable of tolerating an oral dose and is clinically stable. For chronic treatment of hypothyroidism, an oral dosage form of levothyroxine should be used to maintain a euthyroid state. Relative bioavailability between Levothyroxine Sodium for Injection and oral levothyroxine products has not been established. Based on medical practice, the relative bioavailability between oral and intravenous administration of Levothyroxine Sodium for Injection is estimated to be from 48 to 74%. Due to differences in absorption characteristics of patients and the oral levothyroxine product formulations, TSH and thyroid hormone levels should be measured a few weeks after initiating oral levothyroxine and dose adjusted accordingly.

2.2 Dosing in the Elderly and in Patients with Cardiovascular Disease

Intravenous levothyroxine may be associated with cardiac toxicity-including arrhythmias, tachycardia, myocardial ischemia and infarction, or worsening of congestive heart failure and death-in the elderly and in those with underlying cardiovascular disease. Therefore,

cautious use, including doses in the lower end of the recommended range, may be warranted in these populations.

2.3 Reconstitution Directions

Reconstitute the lyophilized Levothyroxine Sodium for Injection by aseptically adding 5 mL of 0.9% Sodium Chloride Injection, USP only. Shake vial to ensure complete mixing. The resultant solution will have a final concentration of approximately 20 mcg per mL, 40 mcg per mL and 100 mcg per mL for the 100 mcg, 200 mcg and 500 mcg vials, respectively. Reconstituted drug product is preservative free and is stable for 4 hours. Discard any unused portion. **DO NOT ADD LEVOTHYROXINE SODIUM FOR INJECTION TO OTHER INTRAVENOUS FLUIDS.** Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

3 DOSAGE FORMS AND STRENGTHS

For injection: a lyophilized powder in a single-dose amber-colored vial containing Levothyroxine Sodium available in the following strengths:

- 100 mcg per vial
- 200 mcg per vial
- 500 mcg per vial

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Cardiac Complications in Elderly and in Patients with Cardiovascular Disease

Excessive bolus dosing of Levothyroxine Sodium for Injection (greater than 500 mcg) are associated with cardiac complications, particularly in the elderly and in patients with an underlying cardiac condition. Adverse events that can potentially be related to the administration of large doses of Levothyroxine Sodium for Injection include arrhythmias, tachycardia, myocardial ischemia and infarction, or worsening of congestive heart failure and death. Cautious use, including doses in the lower end of the recommended range, may be warranted in these populations. Close observation of the patient following the administration of Levothyroxine Sodium for Injection is advised.

5.2 Need for Concomitant Glucocorticoids and Monitoring for Other Diseases in Patients with Endocrine Disorders

Occasionally, chronic autoimmune thyroiditis, which can lead to myxedema coma, may occur in association with other autoimmune disorders such as adrenal insufficiency,

pernicious anemia, and insulin-dependent diabetes mellitus. Patients should be treated with replacement glucocorticoids prior to initiation of treatment with Levothyroxine Sodium for Injection, until adrenal function has been adequately assessed. Failure to do so may precipitate an acute adrenal crisis when thyroid hormone therapy is initiated, due to increased metabolic clearance of glucocorticoids by thyroid hormone. With initiation of Levothyroxine Sodium for Injection, patients with myxedema coma should also be monitored for previously undiagnosed diabetes insipidus.

5.3 Not Indicated for Treatment of Obesity

Thyroid hormones, including Levothyroxine Sodium for Injection, either alone or with other therapeutic agents, should not be used for the treatment of obesity or for weight loss. In euthyroid patients, doses within the range of daily hormonal requirements are ineffective for weight reduction. Larger doses may produce serious or even life threatening manifestations of toxicity, particularly when given in association with sympathomimetic amines such as those used for their anorectic effects [*see Adverse Reactions (6) and Overdosage (10)*].

6 ADVERSE REACTIONS

Excessive doses of levothyroxine can predispose to signs and symptoms compatible with hyperthyroidism. The signs and symptoms of thyrotoxicosis include, but are not limited to: exophthalmic goiter, weight loss, increased appetite, palpitations, nervousness, diarrhea, abdominal cramps, sweating, tachycardia, increased pulse and blood pressure, cardiac arrhythmias, angina pectoris, tremors, insomnia, heat intolerance, fever, and menstrual irregularities.

7 DRUG INTERACTIONS

Many drugs affect thyroid hormone pharmacokinetics and metabolism (e.g., synthesis, secretion, catabolism, protein binding, and target tissue response) and may alter the therapeutic response to Levothyroxine Sodium for Injection. In addition, thyroid hormones and thyroid status have varied effects on the pharmacokinetics and actions of other drugs (see Section 12.3).

7.1 Antidiabetic Therapy

Addition of levothyroxine to antidiabetic or insulin therapy may result in increased antidiabetic agent or insulin requirements. Careful monitoring of diabetic control is recommended, especially when thyroid therapy is started, changed, or discontinued.

7.2 Oral Anticoagulants

Levothyroxine increases the response to oral anticoagulant therapy. Therefore, a decrease in the dose of anticoagulant may be warranted with correction of the hypothyroid state or when the Levothyroxine Sodium for Injection dose is increased. Prothrombin time should be closely monitored to permit appropriate and timely dosage adjustments.

7.3 Digitalis Glycosides

The therapeutic effects of digitalis glycosides may be reduced by levothyroxine. Serum digitalis glycoside levels may be decreased when a hypothyroid patient becomes euthyroid, necessitating an increase in the dose of digitalis glycosides.

7.4 Antidepressant Therapy

Concurrent use of tricyclic (e.g., amitriptyline) or tetracyclic (e.g., maprotiline) antidepressants and levothyroxine may increase the therapeutic and toxic effects of both drugs, possibly due to increased receptor sensitivity to catecholamines. Toxic effects may include increased risk of cardiac arrhythmias and CNS stimulation; onset of action of tricyclics may be accelerated. Administration of sertraline in patients stabilized on levothyroxine may result in increased levothyroxine requirements.

7.5 Ketamine

Concurrent use may produce marked hypertension and tachycardia; cautious administration to patients receiving thyroid hormone therapy is recommended.

7.6 Sympathomimetics

Concurrent use may increase the effects of sympathomimetics or thyroid hormone. Thyroid hormones may increase the risk of coronary insufficiency when sympathomimetic agents are administered to patients with coronary artery disease.

7.7 Drug-Laboratory Test Interactions

Changes in thyroxine binding globulin (TBG) concentration must be considered when interpreting levothyroxine and triiodothyronine values, which necessitates measurement and evaluation of unbound (free) hormone and/or determination of the free levothyroxine index. Pregnancy, infectious hepatitis, estrogens, estrogen containing oral contraceptives, and acute intermittent porphyria increase TBG concentrations. Decreases in TBG concentrations are observed in nephrosis, severe hypoproteinemia, severe liver disease, acromegaly, and after androgen or corticosteroid therapy. Familial hyper or hypo thyroxine binding globulinemias have been described, with the incidence of TBG deficiency approximating 1 in 9,000.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category A – There are no reported cases of Levothyroxine Sodium for Injection used to treat myxedema coma in patients who were pregnant or lactating. Studies in pregnant women treated with oral levothyroxine to maintain a euthyroid state have not shown an increased risk of fetal abnormalities. Therefore, pregnant patients who develop myxedema should be treated with Levothyroxine Sodium for Injection as the risk of

nontreatment is associated with a high probability of significant morbidity or mortality to the maternal patient and the fetus.

8.2 Labor and Delivery

Patients in labor who develop myxedema have not been reported in the literature. However, patients should be treated with Levothyroxine Sodium for Injection as the risk of nontreatment is associated with a high probability of significant morbidity or mortality to the maternal patient and the fetus.

8.3 Nursing Mothers

Adequate replacement doses of thyroid hormones are required to maintain normal lactation. There are no reported cases of Levothyroxine Sodium for Injection used to treat myxedema coma in patients who are lactating. However, such patients should be treated with Levothyroxine Sodium for Injection as the risk of nontreatment is associated with a high probability of significant morbidity or mortality to the nursing patient.

8.4 Pediatric Use

Myxedema coma is a disease of the elderly. An approved, oral dosage form of levothyroxine should be used in the pediatric patient population for maintaining a euthyroid state in non-complicated hypothyroidism.

8.5 Geriatric Use and Patients with Underlying Cardiovascular Disease

See Section 2, Dosage and Administration, for full prescribing information in the geriatric patient population. Because of the increased prevalence of cardiovascular disease in the elderly, cautious use of Levothyroxine Sodium for Injection in the elderly and in patients with known cardiac risk factors is advised. Atrial fibrillation is a common side effect associated with levothyroxine treatment in the elderly [*see Dosage and Administration (2) and Warnings and Precautions (5)*].

10 OVERDOSAGE

In general, the signs and symptoms of overdosage with levothyroxine are those of hyperthyroidism [*see Warnings and Precautions (5) and Adverse Reactions (6)*]. In addition, confusion and disorientation may occur. Cerebral embolism, shock, coma, and death have been reported. Excessive doses of Levothyroxine Sodium for Injection (greater than 500 mcg) are associated with cardiac complications in patients with underlying cardiac disease.


Treatment of Overdosage

Levothyroxine Sodium for Injection should be reduced in dose or temporarily discontinued if signs or symptoms of overdose occur. To obtain up-to-date information about the treatment of overdose, a good resource is the certified Regional Poison Control Center. In managing overdose, consider the possibility of multiple drug overdoses, interaction among drugs, and unusual drug kinetics in the patient.

In the event of an overdose, appropriate supportive treatment should be initiated as dictated by the patient's medical status.

11 DESCRIPTION

Levothyroxine Sodium for Injection contains synthetic crystalline levothyroxine (L-thyroxine) sodium salt. Levothyroxine sodium has an empirical formula of $C_{15}H_{10}I_4NNaO_4$, a molecular weight of 798.85 g/mol (anhydrous), and the following structural formula:



Levothyroxine Sodium for Injection is a sterile, preservative-free lyophilized powder. Each vial contains: 100 mcg, 200 mcg or 500 mcg of levothyroxine sodium (the active ingredient), and the excipients dibasic sodium phosphate heptahydrate 0.944 mg; mannitol 3.0 mg; sodium hydroxide 0.35 mg; and sodium hydroxide for pH adjustment at the range of pH value 9.5-11.5.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Thyroid hormones exert their physiologic actions through control of DNA transcription and protein synthesis. Triiodothyronine (T_3) and levothyroxine (T_4) diffuse into the cell nucleus and bind to thyroid receptor proteins attached to DNA. This hormone nuclear receptor complex activates gene transcription and synthesis of messenger RNA and cytoplasmic proteins.

The physiological actions of thyroid hormones are produced predominantly by T_3 , the majority of which (approximately 80%) is derived from T_4 by deiodination in peripheral tissues.

12.2 Pharmacodynamics

Thyroid hormone synthesis and secretion is regulated by the hypothalamic pituitary-thyroid axis. Thyrotropin releasing hormone (TRH) released from the hypothalamus stimulates secretion of thyrotropin stimulating hormone (TSH) from the anterior pituitary. TSH, in

turn, is the physiologic stimulus for the synthesis and secretion of thyroid hormones, T_4 and T_3 , by the thyroid gland. Circulating serum T_3 and T_4 levels exert a feedback effect on both TRH and TSH secretion. When serum T_3 and T_4 levels increase, TRH and TSH secretion decrease. When thyroid hormone levels decrease, TRH and TSH secretion increases. TSH is used for the diagnosis of hypothyroidism and evaluation of levothyroxine therapy adequacy with other laboratory and clinical data [see Dosage (2.1)]. There are drugs known to affect thyroid hormones and TSH by various mechanisms and those examples are diazepam, ethionamide, lovastatin, metoclopramide, 6-mercaptopurine, nitroprusside, perphenazine, and thiazide diuretics. Some drugs may cause a transient decrease in TSH secretion without hypothyroidism and those drugs (dose) are dopamine (greater than 1 mcg per kg per min), glucocorticoids (hydrocortisone greater than 100 mg per day or equivalent) and octreotide (greater than 100 mcg per day).

Thyroid hormones regulate multiple metabolic processes and play an essential role in normal growth and development, and normal maturation of the central nervous system and bone. The metabolic actions of thyroid hormones include augmentation of cellular respiration and thermogenesis, as well as metabolism of proteins, carbohydrates and lipids. The protein anabolic effects of thyroid hormones are essential to normal growth and development.

12.3 Pharmacokinetics

Absorption – Levothyroxine Sodium for Injection is administered via the intravenous route. Following administration, the synthetic levothyroxine cannot be distinguished from the natural hormone that is secreted endogenously.

Distribution – Circulating thyroid hormones are greater than 99% bound to plasma proteins, including thyroxine binding globulin (TBG), thyroxine binding prealbumin (TBPA), and albumin (TBA), whose capacities and affinities vary for each hormone. The higher affinity of both TBG and TBPA for T_4 partially explains the higher serum levels, slower metabolic clearance, and longer half life of T_4 compared to T_3 . Protein bound thyroid hormones exist in reverse equilibrium with small amounts of free hormone. Only unbound hormone is metabolically active. Many drugs and physiologic conditions affect the binding of thyroid hormones to serum proteins [see Warnings and Precautions (5) and Drug Interactions (7)]. Thyroid hormones do not readily cross the placental barrier [see Warnings and Precautions (5) and Use in Specific Populations (8)].

Metabolism – T_4 is slowly eliminated. The major pathway of thyroid hormone metabolism is through sequential deiodination. Approximately eighty percent of circulating T_3 is derived from peripheral T_4 by monodeiodination. The liver is the major site of degradation for both T_4 and T_3 , with T_4 deiodination also occurring at a number of additional sites,

including the kidney and other tissues. Approximately 80% of the daily dose of T₄ is deiodinated to yield equal amounts of T₃ and reverse T₃ (r T₃). T₃ and r T₃ are further deiodinated to diiodothyronine. Thyroid hormones are also metabolized via conjugation with glucuronides and sulfates and excreted directly into the bile and gut where they undergo enterohepatic recirculation.

Elimination – Thyroid hormones are primarily eliminated by the kidneys. A portion of the conjugated hormone reaches the colon unchanged, where it is hydrolyzed and eliminated in feces as the free hormones. Urinary excretion of T₄ decreases with age.

Table 1: Pharmacokinetic Parameters of Thyroid Hormones in Euthyroid Patients

Hormone	Ratio in Thyroglobulin	Biologic Potency	Half-Life (Days)	Protein Binding (%) ²
T ₄	10 to 20	1	6 to 8 ¹	99.96
T ₃	1	4	≤ 2	99.5

T₄: Levothyroxine

T₃: Liothyronine

¹ 3 to 4 days in hyperthyroidism, 9 to 10 days in hypothyroidism.

² Includes TBG, TBPA, and TBA.

Drug Interactions

A listing of drug interaction with T₄ is provided in the following tables, although it may not be comprehensive due to the introduction of new drugs that interact with the thyroidal axis or the discovery of previously unknown interactions. The prescriber should be aware of this fact and should consult appropriate reference sources (e.g., package inserts of newly approved drugs, medical literature) for additional information if a drug-drug interaction with levothyroxine is suspected.

Table 2: Drugs That May Alter T₄ and T₃ Serum Transport Without Affecting free T₄ Concentration (Euthyroidism)

Drugs That May Increase Serum TBG Concentration	Drugs That May Decrease Serum TBG Concentration
Clofibrate Estrogen-containing oral contraceptives Estrogens (oral) Heroin/Methadone 5-Fluorouracil Mitotane Tamoxifen	Androgens/Anabolic Steroids Asparaginase Glucocorticoids Slow-Release Nicotinic Acid

Drugs That May Cause Protein-Binding Site Displacement	
Potential impact: Administration of these agents with levothyroxine results in an initial transient increase in FT ₄ . Continued administration results in a decrease in serum T ₄ and normal FT ₄ and TSH concentrations and, therefore, patients are clinically euthyroid.	
Salicylates (greater than 2 g/day)	Salicylates inhibit binding of T ₄ and T ₃ to TBG and transthyretin. An initial increase in serum FT ₄ is followed by return of FT ₄ to normal levels with sustained therapeutic serum salicylate concentrations, although total-T ₄ levels may decrease by as much as 30%.
Other drugs: Furosemide (greater than 80 mg IV) Heparin Hydantoins Non-Steroidal Anti-inflammatory Drugs - Fenamates - Phenylbutazone	

Table 3: Drugs That May Alter Hepatic Metabolism of T₄ (Hypothyroidism)

Potential impact: Stimulation of hepatic microsomal drug-metabolizing enzyme activity may cause increased hepatic degradation of levothyroxine, resulting in increased levothyroxine requirements.

Drug or Drug Class	
Carbamazepine Hydantoins	Phenytoin and carbamazepine reduce serum protein binding of levothyroxine, and total- and free-T ₄ may be reduced by 20% to 40%, but most patients have normal serum TSH levels and are clinically euthyroid.
Other drugs: Phenobarbital Rifampin	

Table 4: Drugs That May Decrease Conversion of T₄ to T₃

Potential impact: Administration of these enzyme inhibitors decreases the peripheral conversion of T₄ to T₃, leading to decreased T₃ levels. However, serum T₄ levels are usually normal but may occasionally be slightly increased.

Drug or Drug Class	Effect
---------------------------	---------------

Beta-adrenergic antagonists (e.g., Propranolol greater than 160 mg/day)	In patients treated with large doses of propranolol (greater than 160 mg/day), T ₃ and T ₄ levels change slightly, TSH levels remain normal, and patients are clinically euthyroid. It should be noted that actions of particular beta-adrenergic antagonists may be impaired when the hypothyroid patient is converted to the euthyroid state.
Glucocorticoids (e.g., Dexamethasone greater than or equal to 4 mg/day)	Short-term administration of large doses of glucocorticoids may decrease serum T ₃ concentrations by 30% with minimal change in serum T ₄ levels. However, long-term glucocorticoid therapy may result in slightly decreased T ₃ and T ₄ levels due to decreased TBG production (see above).
Other drug: Amiodarone	

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Animal studies have not been performed to evaluate the carcinogenic potential, mutagenic potential or effects on fertility of Levothyroxine Sodium for Injection.

13.2 Animal Toxicology and Pharmacology

No animal toxicology studies have been conducted with Levothyroxine Sodium for Injection.

14 CLINICAL STUDIES

No clinical studies have been conducted with Levothyroxine Sodium for Injection in patients with myxedema coma. However, data from published literature support the intravenous use of levothyroxine sodium for the treatment of myxedema coma.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Levothyroxine Sodium for Injection, a lyophilized powder in a single-dose amber-colored vial containing Levothyroxine Sodium available in the following strengths:

Product Code	Strength	Reconstituted Concentration	Each
506107	100 mcg per vial	20 mcg per mL	NDC 63323-649-07 One Single-Dose Vial per Carton
506247	200 mcg per vial	40 mcg per mL	NDC 63323-647-10 One Single-Dose Vial per Carton
506248	500 mcg per vial	100 mcg per mL	NDC 63323-648-10 One Single-Dose Vial per Carton

16.2 Storage and Handling

Protect from light and store dry product at 20°C to 25°C (68°F to 77°F) [see USP Controlled Room Temperature]. Reconstituted drug product is preservative free and stable for 4 hours. Discard any unused portion.

This container closure is not made with natural rubber latex.



Lake Zurich, IL 60047

www.fresenius-kabi.com/us

Package Insert Part Number Pending

NDC 63323-649-07 506107

Levothyroxine Sodium for Injection

100 mcg per vial

For Intravenous Use

Single-Dose Vial

Discard any unused portion.

Rx only

Sterile, Lyophilized

Preservative Free

Each vial contains: Levothyroxine sodium 100 mcg; inactive ingredients: see carton label.

Recommended Dosage: See Prescribing Information.

Store dry product at 20°C to 25°C (68°F to 77°F).

PROTECT FROM LIGHT.

Reconstitution Directions: Reconstitute the lyophilized levothyroxine sodium by aseptically adding 5 mL of 0.9% Sodium Chloride Injection, USP. Shake vial to ensure complete mixing. The resultant solution will have a final concentration of 20 mcg/mL. Use immediately after reconstitution. Discard any unused portion.

Fresenius Kabi

Lake Zurich, IL 60047



3 63323-649-07 4

XXXXXX

LOT/EXP

LOT: LLLLLL
EXP: YYYY-MM



32"



SN: L11111
LOT: L11111
EXP: YYYY-MM



GTIN: 00363323649074

100 mcg per vial

Levothyroxine Sodium
for Injection
NDC 63323-649-07

NDC 63323-649-07 506107

Preservative Free

Sterile, Lyophilized

Each vial contains:

Levothyroxine sodium 100 mcg;
dibasic sodium phosphate
heptahydrate 0.944 mg;
mannitol 3.0 mg; sodium
hydroxide 0.35 mg; and sodium
hydroxide for pH adjustment at the
range of pH value 9.5-11.5.

Recommended Dosage: See
Prescribing Information.

This container closure is
not made with natural rubber latex.



Lake Zurich, IL 60047
www.fresenius-kabi.com/us

Store dry product at
20°C to 25°C (68°F to 77°F)
[see USP Controlled Room
Temperature].

PROTECT FROM LIGHT.

Reconstitution Directions:
Reconstitute the lyophilized
levothyroxine sodium by
aseptically adding 5 mL
of 0.9% Sodium Chloride
Injection, USP. Shake vial
to ensure complete mixing.
The resultant solution will
have a final concentration
of 20 mcg/mL. Use imme-
diately after reconstitution.
Discard any unused portion.



NDC 63323-649-07 506107

**Levothyroxine
Sodium
for Injection**


100 mcg per vial

For Intravenous Use

Single-Dose Vial

Discard any unused
portion.

Rx only



3 63323-649-07 4

XXXXXX



1/32"

NDC 63323-647-10 506247

Levothyroxine Sodium for Injection

200 mcg per vial

For Intravenous Use

Single-Dose Vial

Discard any unused portion.

Rx only

Sterile, Lyophilized

Preservative Free

Each vial contains: Levothyroxine sodium
200 mcg; inactive ingredients: see carton label.

Recommended Dosage: See Prescribing
Information.

Store dry product at 20°C to 25°C (68°F to 77°F).

PROTECT FROM LIGHT.

Reconstitution Directions: Reconstitute the
lyophilized levothyroxine sodium by aseptically
adding 5 mL of 0.9% Sodium Chloride Injection,
USP. Shake vial to ensure complete mixing.

The resultant solution will have a final
concentration of 40 mcg/mL. Use immediately
after reconstitution. Discard any unused portion.

Fresenius Kabi

Lake Zurich, IL 60047



3

63323-647-10

0

XXXXXXXX

LOT/EXP

LOT: LLLLLL
EXP: YYYY-MM



EXP: LOT: YYY-MM
NDC 63323-647-10



GTIN: 00363323647100

200 mcg per vial

Levothyroxine Sodium
for Injection
NDC 63323-647-10

NDC 63323-647-10 506247

Preservative Free

Sterile, Lyophilized

Each vial contains:

Levothyroxine sodium 200 mcg;
dibasic sodium phosphate
heptahydrate 0.944 mg;
mannitol 3.0 mg; sodium
hydroxide 0.35 mg; and sodium
hydroxide for pH adjustment at the
range of pH value 9.5-11.5.

Recommended Dosage: See
Prescribing Information.

This container closure is
not made with natural rubber latex.

 **FRESENIUS
KABI**
Lake Zurich, IL 60047
www.fresenius-kabi.com/us

Store dry product at
20°C to 25°C (68°F to 77°F)
[see USP Controlled Room
Temperature].

PROTECT FROM LIGHT.

Reconstitution Directions:
Reconstitute the lyophilized
levothyroxine sodium by
aseptically adding 5 mL of
0.9% Sodium Chloride
Injection, USP. Shake vial
to ensure complete mixing.
The resultant solution will
have a final concentration of
40 mcg/mL. Use immediately
after reconstitution. Discard
any unused portion.

 **FRESENIUS
KABI**


NDC 63323-647-10 506247

**Levothyroxine
Sodium**
for Injection

200 mcg per vial

For Intravenous Use
Single-Dose Vial
Discard any unused
portion.
Rx only

 **FRESENIUS
KABI**



XXXXXX

NDC 63323-648-10 506248

Levothyroxine Sodium for Injection

500 mcg per vial

For Intravenous Use

Single-Dose Vial

Discard any unused portion.

Rx only

Sterile, Lyophilized

Preservative Free

Each vial contains: Levothyroxine sodium
500 mcg; inactive ingredients: see carton label.

Recommended Dosage: See Prescribing
Information.

Store dry product at 20°C to 25°C (68°F to 77°F).
PROTECT FROM LIGHT.

Reconstitution Directions: Reconstitute
the lyophilized levothyroxine sodium by
aseptically adding 5 mL of 0.9% Sodium
Chloride Injection, USP. Shake vial to ensure
complete mixing. The resultant solution will
have a final concentration of 100 mcg/mL. Use
immediately after reconstitution. Discard any
unused portion.

Fresenius Kabi

Lake Zurich, IL 60047



3

63323-648-10

7

XXXXXXXX

LOT/EXP

LOT: LLLLLL
EXP: YYYY-MM



EXP: YYY-YY-MM

SN: LLLLLL
LOT: LLLLLL



#####

GTIN: 0036323648107

500 mcg per vial

Levothyroxine Sodium
for Injection
NDC 63323-648-10

NDC 63323-648-10 506248

Preservative Free

Sterile, Lyophilized

Each vial contains:
Levothyroxine sodium 500 mcg;
dibasic sodium phosphate
heptahydrate 0.944 mg;
mannitol 3.0 mg; sodium
hydroxide 0.35 mg; and sodium
hydroxide for pH adjustment at the
range of pH value 9.5-11.5.

Recommended Dosage: See
Prescribing Information.

This container closure is
not made with natural rubber latex.



Lake Zurich, IL 60047
www.fresenius-kabi.com/us

Store dry product at
20°C to 25°C (68°F to 77°F)
[see USP Controlled Room
Temperature].

PROTECT FROM LIGHT.

Reconstitution Directions:
Reconstitute the lyophilized
levothyroxine sodium by
aseptically adding 5 mL of
0.9% Sodium Chloride
Injection, USP. Shake vial
to ensure complete mixing.
The resultant solution will
have a final concentration of
100 mcg/mL. Use immediately
after reconstitution. Discard
any unused portion.



NDC 63323-648-10 506248

**Levothyroxine
Sodium**
for Injection


500 mcg per vial

For Intravenous Use

Single-Dose Vial

Discard any unused
portion.

Rx only



3 63323-648-10 7

XXXXXX



Ramesh
Raghavachari

Digitally signed by Ramesh Raghavachari
Date: 7/06/2026 06:21:50PM
GUID: 502d0913000029f375128b0de8c50020