

NDC 76329-3869-2

For use in the Nose Only

166



REXTOVY™ Naloxone HCl
Nasal Spray 4 mg



3 76329386928

Distributed by:
IMS, LIMITED 76386908/2/26
So. El Monte, CA 91733, U.S.A.

XXXXXX
EXP YYYY-MM



INTERNATIONAL MEDICATION SYSTEMS, LIMITED
So. El Monte, CA 91733, U.S.A.
An Amphoslar Pharmaceuticals Company
872890C/2/26

REXTOVY™
Naloxone HCl Nasal Spray 4 mg
Emergency Treatment of Opioid Overdose

- For use in the nose only
- Do not test nasal spray device before use
- 1 nasal spray device contains 1 dose of medicine
- Each device sprays 1 time only
- Do not use if the blister is open or torn, or if the device appears damaged

NDC 76329-3869-2 STOCK NO. 3869

Read enclosed Directions

- Store at 20°C to 25°C (68°F to 77°F)
- Do not freeze
- Avoid excessive heat above 40°C (104°F)
- Protect from light

Lot: XXXXXX

Exp: YYYY-MM

Unfold for Directions

Directions REXTOVY (rex toe' vee) Naloxone Hydrochloride Nasal Spray 4 mg

Emergency Treatment of Opioid Overdose

Important:

- For use in the **nose only**
- **Do not** test nasal spray device before use
- 1 nasal spray device contains 1 dose of medicine
- Each device sprays 1 time only
- **Do not** use if the blister is open or torn, or if the device appears damaged




1) CHECK



Step 1: CHECK if you suspect an overdose:

- **CHECK** for a suspected overdose: the person will not wake up or is very sleepy or not breathing well
 - yell "Wake up!"
 - shake the person gently
 - if the person is not awake, go to Step 2

2) GIVE



Step 2: GIVE 1st dose in the nose

- **HOLD** the nasal spray device with your thumb on the bottom of the green plunger and fingers on the finger tabs
- **INSERT** the nozzle **all the way** into either NOSTRIL
- **PRESS** the green plunger all the way down until it stops to give the 1st dose
- 1 nasal spray device contains 1 dose

3) CALL



Step 3: CALL 911

- **CALL 911** immediately after giving the 1st dose

4) WATCH/GIVE



Step 4: WATCH & GIVE

- **WAIT** 2-3 minutes after the 1st dose to give the medicine time to work
- if the person wakes up: Go to Step 5
- if the person does not wake up:
 - **CONTINUE TO GIVE** doses every 2-3 minutes until the person wakes up
 - it is safe to keep giving doses

5) STAY



Step 5: STAY

- **STAY** until ambulance arrives: even if the person wakes up
- **GIVE** another dose if the person becomes very sleepy again
- You may need to give all the doses in the pack

For opioid emergencies, call 911.

For more information, go to www.amphastar.com or call 1-800-423-4136.



Distributed by International Medication Systems, Limited, So. El Monte, CA 91733, U.S.A.
An Amphastar Pharmaceuticals Company
7038690E/3-26




Drug Facts (continued)

Directions



Step 1: CHECK if you suspect an overdose:

- **CHECK** for a suspected overdose; the person will not wake up or is very sleepy or not breathing well
- yell "Wake up!"
- shake the person gently
- if the person is not awake, go to Step 2



Step 2: GIVE 1st dose in the nose

- **HOLD** the nasal spray device with your thumb on the bottom of the green plunger and fingers on the finger tabs
- **INSERT** the nozzle all the way into either NOSTRIL
- **PRESS** the green plunger all the way down until it stops to give the 1st dose
- 1 nasal spray device contains 1 dose



Step 3: CALL 911

- **CALL 911** immediately after giving the 1st dose



Step 4: WATCH & GIVE

- **WAIT** 2-3 minutes after the 1st dose to give the medicine time to work
- if the person **wakes up**: Go to Step 5
- if the person **does not wake up**:
 - **CONTINUE TO GIVE** doses every 2-3 minutes until the person wakes up
 - it is safe to keep giving doses



Step 5: STAY

- **STAY** until ambulance arrives: even if the person wakes up
- **GIVE** another dose if the person becomes very sleepy again
- You may need to give all the doses in the pack

Drug Facts (continued)

Warning
When using this product some people may experience symptoms when they wake up, such as shaking, sweating, nausea, or feeling angry. This is to be expected.

Other information

- store at 20°C to 25°C (68°F to 77°F)
- do not freeze
- avoid excessive heat above 40°C (104°F)
- protect from light
- the product is packaged in individual sealed blisters
- do not use if the blister is open or torn, or if the device appears damaged

Inactive ingredients
sodium chloride, sodium hydroxide to adjust pH, and water for injection USP

Questions?
call 1-800-423-4136 or go to www.amphastar.com
Mon.- Fri. 7 am - 5 pm PST

Safe to Use
Even if Opioids are Not Present



NDC 76329-3869-2
INTERNATIONAL MEDICATION SYSTEMS, LIMITED
So. El Monte, CA 91733, U.S.A.
An Amphastar Pharmaceuticals Company
www.amphastar.com
5638690F3-26 Stock No. 3869

THIS AREA REPRESENTS WHERE THE SERIAL NUMBER, LOT NUMBER AND EXPIRATION DATE (IN THE FORMAT: EXP YYYY-MM) ARE PLACED.

REXTOVY™
Naloxone HCl Nasal Spray 4 mg

Emergency Treatment of Opioid Overdose

For use in nose only

Easy to Use

Can Save a Life



2 Single-Dose Nasal Spray Devices
4 mg per Device
0.008 fl. oz. (0.25 mL) each



Do not test nasal spray device before use

1 nasal spray device contains 1 dose of medicine

Each device sprays 1 time only

Read Directions Before Use ▶

Drug Facts

Active ingredient (in each spray)	Purpose
Naloxone hydrochloride 4 mg	Emergency treatment of opioid overdose

Uses

- to "revive" someone during an overdose from many prescription pain medications or street drugs such as heroin
- this medicine can save a life

2.25"

UNUSED

5.625"

9.6875"

1.75"