

298376



Read directions and warnings before use. Keep this carton. It has important information.

Drug Facts

Active ingredient (in each tablet)
Levocetirizine dihydrochloride 5 mg.....Antihistamine

Purpose

Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies:
■ runny nose ■ itchy, watery eyes
■ sneezing ■ itching of the nose or throat

Warnings

Do not use
■ if you have kidney disease
■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have
■ ever had trouble urinating or emptying your bladder

When using this product
■ drowsiness may occur
■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
■ in rare cases, you may experience itching after stopping levocetirizine. Consult your healthcare provider if this happens.

Stop use and ask a doctor if
■ you have trouble urinating or emptying your bladder
■ an allergic reaction to this product occurs. Seek medical help right away.

Drug Facts (continued)

If pregnant or breast-feeding:
■ if breast-feeding: not recommended
■ if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

adults 65 years of age and older
■ ask a doctor

adults and children 12-64 years of age
■ take 1 tablet (5 mg) once daily in the evening
■ do not take more than 1 tablet (5 mg) in 24 hours
■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms

children 6-11 years of age
■ take ½ tablet (2.5 mg) once daily in the evening
■ do not take more than ½ tablet (2.5 mg) in 24 hours

children under 6 years of age
■ do not use
consumers with kidney disease
■ do not use

Drug Facts (continued)

Other information

■ store between 20° and 25°C (68° and 77°F)
■ safety sealed: do not use if carton was opened or if individual blister unit is open or torn

Drug Facts (continued)
Inactive ingredients
Colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide
Questions or comments? call 1-866-844-2798

SAMPLE NOT FOR RETAIL SALE



ORIGINAL PRESCRIPTION STRENGTH

XYZAL

ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

5 TABLETS

CLINICALLY PROVEN RELIEF IN 60 MINUTES

The trade dress of this Xyzal® package is subject to trademark protection.

Distributed by Chatterm, Inc., Chattanooga, TN 37409 ©2026

Origin: India 943862 NDC 41167-3510-6



943862

658576

CLINICALLY PROVEN RELIEF FOR 24 HOURS

The trade dress of this Xyzal® package is subject to trademark protection.

Distributed by Chatterm, Inc., Chattanooga, TN 37409 ©2025

Origin: India 943859 NDC 41167-3510-0



Drug Facts

Active ingredient (in each tablet) Purpose
Levocetirizine dihydrochloride 5 mg...Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other respiratory allergies:
■ runny nose ■ itchy, watery eyes
■ sneezing ■ itching of the nose or throat

Warnings

Do not use
■ if you have kidney disease
■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have
■ ever had trouble urinating or emptying your bladder

When using this product

■ drowsiness may occur
■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
■ in rare cases, you may experience itching after stopping levocetirizine. Consult your healthcare provider if this happens.

Stop use and ask a doctor if

■ you have trouble urinating or emptying your bladder
■ an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

■ if breast-feeding: not recommended
■ if pregnant: ask a health professional before use
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions

adults 65 years of age and older ■ ask a doctor
adults and children 12-64 years of age ■ take 1 tablet (5 mg) once daily in the evening
■ do not take more than 1 tablet (5 mg) in 24 hours
■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms
children 6-11 years of age ■ take ½ tablet (2.5 mg) once daily in the evening
■ do not take more than ½ tablet (2.5 mg) in 24 hours
children under 6 years of age ■ do not use
consumers with kidney disease ■ do not use

Other information

■ store between 20° and 25°C (68° and 77°F)
■ safety sealed: do not use if carton was opened or if individual blister unit is open or torn

Inactive ingredients

colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide

Questions or comments?

call 1-866-844-2798

Read directions and warnings before use. Keep this carton. It has important information.

CLINICALLY PROVEN RELIEF FROM INDOOR AND OUTDOOR ALLERGIES

CLINICALLY PROVEN RELIEF IN 60 MINUTES



ORIGINAL PRESCRIPTION STRENGTH

XYZAL®

ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

10 TABLETS



Actual Size

24 HOUR RELIEF OF
○ SNEEZING
○ RUNNY NOSE
○ ITCHY NOSE OR THROAT
○ ITCHY, WATERY EYES



0 41167 35100 0

LOT:

EXP:

943859

943863

CLINICALLY PROVEN RELIEF FOR 24 HOURS

The trade dress of this Xyzal® package is subject to trademark protection.

Distributed by
Chattam, Inc.
Chattanooga, TN 37409
©2025

Origin: India

943863
NDC 41167-3511-2

Certified
B
Corporation

EPC
Sustainable Forestry
Call for details of custody
www.forest.org

Drug Facts

Active ingredient (in each tablet)	Purpose
Levocetirizine dihydrochloride 5 mg...Antihistamine	

Uses
temporarily relieves these symptoms due to hay fever or other respiratory allergies:

■ runny nose	■ itchy, watery eyes
■ sneezing	■ itching of the nose or throat

Warnings
Do not use

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have

- ever had trouble urinating or emptying your bladder

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- in rare cases, you may experience itching after stopping levocetirizine. Consult your healthcare provider if this happens.

Stop use and ask a doctor if

- you have trouble urinating or emptying your bladder
- an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions

adults 65 years of age and older	■ ask a doctor
adults and children 12-64 years of age	■ take 1 tablet (5 mg) once daily in the evening
	■ do not take more than 1 tablet (5 mg) in 24 hours
children 6-11 years of age	■ ½ tablet (2.5 mg) once daily in the evening
	■ may be appropriate for less severe symptoms
children under 6 years of age	■ do not use
	■ do not use

Other information

- store between 20° and 25°C (68° and 77°F)
- safety sealed: do not use if carton was opened or if individual blister unit is open or torn

Inactive ingredients
colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide

Questions or comments?
call 1-866-844-2798

Read directions and warnings before use. Keep this carton. It has important information.

CLINICALLY PROVEN RELIEF FROM INDOOR AND OUTDOOR ALLERGIES

0 41167 35112 3

CLINICALLY PROVEN RELIEF IN 60 MINUTES



ORIGINAL PRESCRIPTION STRENGTH

XYZAL®

ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

20 TABLETS



Actual Size

LOT:

EXP:

943863

998576

CLINICALLY PROVEN RELIEF
IN 60 MINUTES

The trade dress of this Xyzal® package
is subject to trademark protection.

Distributed by
Chattem, Inc.,
Chattanooga, TN 37409
©2025

Origin: India
943866
NDC 41167-3510-1

Actual Size



24 HOUR RELIEF OF
○ SNEEZING
○ RUNNY NOSE
○ ITCHY NOSE OR THROAT
○ ITCHY, WATERY EYES

35
TABLETS



Actual Size

LOT: _____
EXP: _____



0 41167 35101 7

CLINICALLY
PROVEN
RELIEF
FOR 24
HOURS



Drug Facts

Active ingredient (in each tablet)
Levocetirizine dihydrochloride 5 mg... Antihistamine

Purpose

Uses
temporarily relieves these symptoms due to hay fever or other respiratory allergies:
■ runny nose ■ itchy, watery eyes
■ sneezing ■ itching of the nose or throat

Warnings
Do not use
■ if you have kidney disease
■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have
■ ever had trouble urinating or emptying your bladder

When using this product
■ drowsiness may occur
■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
■ in rare cases, you may experience itching after stopping levocetirizine. Consult your healthcare provider if this happens.

Stop use and ask a doctor if
■ you have trouble urinating or emptying your bladder
■ an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:
■ if breast-feeding: not recommended
■ if pregnant: ask a health professional before use

Drug Facts (continued)

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

adults 65 years of age and older ■ ask a doctor
■ take 1 tablet (5 mg) once daily in the evening
■ do not take more than 1 tablet (5 mg) in 24 hours
adults and children 12-64 years of age ■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms
children 6-11 years of age ■ take ½ tablet (2.5 mg) once daily in the evening
■ do not take more than ½ tablet (2.5 mg) in 24 hours
children under 6 years of age ■ do not use
consumers with kidney disease ■ do not use

Other information

■ store between 20° and 25°C (68° and 77°F)
■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients

colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide

Questions or comments?

call 1-866-844-2798

Read directions and warnings before use. Keep this carton. It has important information.

943866

251776

The trade dress of this Xyzal® package is subject to trademark protection.

Distributed by
Chatterm, Inc.,
Chattanooga, TN 37409
©2025

Origin: India
944157
NDC 41167-3510-2

Actual Size

CLINICALLY PROVEN RELIEF IN 60 MINUTES

CLINICALLY PROVEN RELIEF FOR 24 HOURS

ORIGINAL PRESCRIPTION STRENGTH

XYZAL®

ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

55 TABLETS

Actual Size

Certified Corporation

PAPER BOX PLASTIC BOTTLE

SUSTAINABLE FORESTRY
EPO
Certified Chain of Custody
Producing Sustainable Forestry
www.forests.org
50.XXXXX

Drug Facts

Active ingredient (in each tablet)	Purpose
Levocetirizine dihydrochloride 5 mg... Antihistamine	

Uses
temporarily relieves these symptoms due to hay fever or other respiratory allergies:
 ■ runny nose ■ itchy, watery eyes
 ■ sneezing ■ itching of the nose or throat

Warnings
Do not use
 ■ if you have kidney disease
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have
 ■ ever had trouble urinating or emptying your bladder

When using this product
 ■ drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery
 ■ in rare cases, you may experience itching after stopping levocetirizine. Consult your healthcare provider if this happens.

Stop use and ask a doctor if
 ■ you have trouble urinating or emptying your bladder
 ■ an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:
 ■ if breast-feeding: not recommended
 ■ if pregnant: ask a health professional before use

Drug Facts (continued)
Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

adults 65 years of age and older	■ ask a doctor
adults and children 12-64 years of age	■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours
children 6-11 years of age	■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms
children under 6 years of age	■ take ¼ tablet (2.5 mg) once daily in the evening ■ do not take more than ½ tablet (2.5 mg) in 24 hours
consumers with kidney disease	■ do not use

Other information
 ■ store between 20° and 25°C (68° and 77°F)
 ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients
 Colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide

Questions or comments?
 call 1-866-844-2798

Read directions and warnings before use. Keep this carton. It has important information.

LOT: _____

EXP: _____

0 41167 35102 4

944157

LOT
EXP

XYZAL[®]

ALLERGY 24HR

CLINICALLY PROVEN RELIEF
IN 60 MINUTES

CLINICALLY PROVEN RELIEF
FOR 24 HOURS

Drug Facts

Active ingredient (in each tablet) **Purpose**
Levocetirizine dihydrochloride 5 mg..... Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose ■ itchy, watery eyes
- sneezing ■ itching of the nose or throat

Warnings

Do not use

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have

- ever had trouble urinating or emptying your bladder

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- in rare cases, you may experience itching after stopping levocetirizine. Consult your healthcare provider if this happens.

Stop use and ask a doctor if

- you have trouble urinating or emptying your bladder
- an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). ▶

The trade dress of this Xyzal[®] package is subject to trademark protection.
Distributed by Chatterm, Inc., Chattanooga, TN 37409 ©2025
Origin: India 950476



Drug Facts (continued)

Directions

adults 65 years of age and older	<ul style="list-style-type: none"> ■ ask a doctor
adults and children 12-64 years of age	<ul style="list-style-type: none"> ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms
children 6-11 years of age	<ul style="list-style-type: none"> ■ take ½ tablet (2.5 mg) once daily in the evening ■ do not take more than ½ tablet (2.5 mg) in 24 hours
children under 6 years of age	<ul style="list-style-type: none"> ■ do not use
consumers with kidney disease	<ul style="list-style-type: none"> ■ do not use

Other information

- store between 20° and 25°C (68° and 77°F)
- safety sealed: do not use if sealed package was torn or opened, or if printed foil inner seal on bottle is torn or missing

Inactive ingredients

colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide

Questions or comments?

call toll-free 1-866-844-2798

Read directions and warnings before use. Keep this carton. It has important information.





944379
Distributed by Chatterm, Inc.,
Chattanooga, TN 37409
©2025 Origin: India

24 HOUR
RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

ORIGINAL
PRESCRIPTION STRENGTH

XYZAL®
ALLERGY 24HR
levocetirizine dihydrochloride tablet
5 mg / antihistamine

35
TABLETS

PEEL HERE ▶

Active ingredient (in each tablet)
Levocetirizine dihydrochloride 5 mg..... Antihistamine
Use: temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat **Warnings Do not use** ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine **Ask a doctor before use** ■ if you have ■ ever had trouble urinating or emptying your bladder **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping levocetirizine. Consult your healthcare provider if this happens. **Stop use and ask a doctor if** ■ you have trouble urinating or emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). **Directions adults 65 years of age and older** ▶

■ ask a doctor **adults and children 12-64 years of age** ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms **children 6-11 years of age** ■ take ½ tablet (2.5 mg) once daily in the evening ■ do not take more than ½ tablet (2.5 mg) in 24 hours **children under 6 years of age** ■ do not use **consumers with kidney disease** ■ do not use **Other information** ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing **inactive ingredients** Colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide **Questions or comments?** call 1-866-844-2798

ORIGINAL
PRESCRIPTION STRENGTH

XYZAL®
ALLERGY 24HR
levocetirizine dihydrochloride tablet
5 mg / antihistamine

35
TABLETS

Distributed by Chatterm, Inc.,
Chattanooga, TN 37409
©2025 Origin: India



944383
 Distributed by Chattem, Inc.,
 Chattanooga, TN 37409
 ©2025 Origin: India

24 HOUR
 RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

ORIGINAL
 PRESCRIPTION STRENGTH

XYZAL®
ALLERGY 24HR

levocetirizine dihydrochloride tablet
 5 mg / antihistamine

55
 TABLETS

PEEL HERE ▶

Active ingredient (in each tablet)

Levocetirizine dihydrochloride 5 mg..... Antihistamine
Use: temporarily relieves these symptoms due to hay fever or other
 respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes
 ■ itching of the nose or throat **Warnings Do not use** ■ if you have
 kidney disease ■ if you have ever had an allergic reaction to this
 product or any of its ingredients or to an antihistamine containing
 cetirizine **Ask a doctor before use** if you have ■ ever had trouble
 urinating or emptying your bladder **When using this product**
 ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives,

Purpose

and tranquilizers may increase drowsiness ■ be careful when driving a
 motor vehicle or operating machinery ■ in rare cases, you may experience
 itching after stopping levocetirizine. Consult your healthcare provider if this
 happens. **Stop use and ask a doctor if** ■ you have trouble urinating or
 emptying your bladder ■ an allergic reaction to this product occurs. Seek
 medical help right away. **If pregnant or breast-feeding:** ■ if
 breast-feeding: not recommended ■ if pregnant: ask a health professional
 before use **Keep out of reach of children.** In case of overdose, get
 medical help or contact a Poison Control Center right away
 (1-800-222-1222). **Directions adults 65 years of age and older** ▶

■ ask a doctor **adults and children 12-64 years of age** ■ take 1 tablet (5 mg) once
 daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ ½ tablet (2.5
 mg) once daily in the evening may be appropriate for less severe symptoms **children 6-11**
years of age ■ take ½ tablet (2.5 mg) once daily in the evening ■ do not take more than
 ½ tablet (2.5 mg) in 24 hours **children under 6 years of age** ■ do not use **consumers**
with kidney disease ■ do not use **Other information** ■ store between 20° and 25°C
 (68° and 77°F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal
 on bottle is torn or missing **inactive ingredients** Colloidal anhydrous silica, hypromellose,
 lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol
 400, titanium dioxide **Questions or comments?** call 1-866-844-2798

Distributed by Chattem, Inc.,
 Chattanooga, TN 37409
 ©2025 Origin: India

ORIGINAL
 PRESCRIPTION STRENGTH

XYZAL®
ALLERGY 24HR

levocetirizine dihydrochloride tablet
 5 mg / antihistamine

55
 TABLETS



944385
 Distributed by Chattem, Inc.,
 Chattanooga, TN 37409
 ©2025 Origin: India

24 HOUR
 RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

PEEL HERE ▶

ORIGINAL
 PRESCRIPTION STRENGTH

XYZAL®
ALLERGY 24HR

levocetirizine dihydrochloride tablet
 5 mg / antihistamine

80
 TABLETS

Active ingredient (in each tablet)

Levocetirizine dihydrochloride 5 mg..... Antihistamine
Use: temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat. **Warnings Do not use** ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine. **Ask a doctor before use** if you have ■ ever had trouble urinating or emptying your bladder. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives,

Purpose

and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ in rare cases, you may experience itching after stopping levocetirizine. Consult your healthcare provider if this happens. **Stop use and ask a doctor if** ■ you have trouble urinating or emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). **Directions adults 65 years of age and older** ▶

■ ask a doctor **adults and children 12-64 years of age** ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ **½ tablet (2.5 mg) once daily** in the evening may be appropriate for less severe symptoms **children 6-11 years of age** ■ take ½ tablet (2.5 mg) once daily in the evening ■ do not take more than ½ tablet (2.5 mg) in 24 hours **children under 6 years of age** ■ do not use **consumers with kidney disease** ■ do not use. **Other information** ■ store between 20° and 25° C (68° and 77° F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing. **Inactive ingredients** Colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide. **Questions or comments?** call 1-866-844-2798

Distributed by Chattem, Inc.,
 Chattanooga, TN 37409
 ©2025 Origin: India

ORIGINAL
 PRESCRIPTION STRENGTH

XYZAL®
ALLERGY 24HR

levocetirizine dihydrochloride tablet
 5 mg / antihistamine

80
 TABLETS

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

DOROTHY N CHANG
05/20/2026 09:41:31 AM