

80.3 [3.164]

19 [0.749]

100 [3.940]

1Y94KFAV1-T	
<p>Dissolve Tabs Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p> <p>1. SEPARATE AT PERFORATION. 2. PEEL AT ARROW.</p>	<p>START HERE</p> <p>Exp YYYY/MMM Batch No. 000000000</p> <p>PEEL</p>
<p>Dissolve Tabs Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p> <p>1. SEPARATE AT PERFORATION. 2. PEEL AT ARROW.</p>	<p>Exp YYYY/MMM Batch No. 000000000</p> <p>PEEL</p>
<p>Dissolve Tabs Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p> <p>1. SEPARATE AT PERFORATION. 2. PEEL AT ARROW.</p>	<p>Exp YYYY/MMM Batch No. 000000000</p> <p>PEEL</p>
<p>Place tablet on tongue; tablet disintegrates, with or without water. Do not crush or chew tablets. One 14-day course of treatment. Take 1 tablet every day for 14 days. Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.</p>	<p>PEEL</p> <p>Exp YYYY/MMM Batch No. 000000000</p>
	<p>Dissolve Tabs Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg acid reducer</p> <p>Dexcel Pharma Technologies Ltd 10 Hakidma Street, Yokneam 2069200, Israel</p> <p>1. SEPARATE AT PERFORATION. 2. PEEL AT ARROW.</p>

↔
Unsealed area

Batch No.
EXP
1974 FA VI-T 060425

Drug Facts

Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer

Purpose ■ treats frequent heartburn (occurs *2 or more* days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings ■ rash ■ skin redness ■ blisters ■ if an allergic reaction occurs, stop use and seek medical help right away.

Allergy alert: ■ do not use if you are allergic to omeprazole. ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ heartburn with **lightheadedness, sweating or dizziness** ■ chest or shoulder pain with shortness of breath; sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ chest or shoulder pain with shortness of breath; sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Ask a doctor before use if you have: ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. ■ **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue, tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age; ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, flavoc, hypromellose, hypromellose phthalate, maize malectan, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Made in Israel

Manufactured by:
Celexel Pharma Technologies Ltd
10 Hakdama Street, Yokneam 2089200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Location for lot number and expiration date

NOT FOR RESALE

Safety Feature –
Do not use if printed tablet blister unit is open or torn.

Compare to Prilosec OTC®*



Treats FREQUENT Heartburn! **24HR**

DISSOLVE TABS

Omeprazole

Delayed Release Orally Disintegrating Tablets 20 mg
Acid Reducer

Melts in your mouth
Dissolves without water



14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

FORMAT: STANDARD

TYPE SIZES:

TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, ferrous oxide, flavor, hypromellose, hypromellose phthalate, maize methylcellulose, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours ■ this product is to be used once a day every 24 hours, every day for 14 days

Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: skin reddening ■ blisters ■ rash ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts (continued) ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day every 24 hours, every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

4 months unless directed by a doctor ■ do not take for more than 14 days ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

If pregnant or breast-feeding, ask a health professional before use. ■ you develop a rash or joint pain ■ you need to take more than 1 course of treatment every 4 months ■ you need to take this product for more than 14 days

Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you get diarrhea ■ you develop a rash or joint pain ■ you need to take more than 1 course of treatment every 4 months

Ask a doctor before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ frequent heartburn, particularly with heartburn ■ heartburn with lightheadedness, sweating or dizziness

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: skin reddening ■ blisters ■ rash ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

4 months unless directed by a doctor ■ do not take for more than 14 days ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day every 24 hours, every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: skin reddening ■ blisters ■ rash ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

4 months unless directed by a doctor ■ do not take for more than 14 days ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day every 24 hours, every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: skin reddening ■ blisters ■ rash ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

4 months unless directed by a doctor ■ do not take for more than 14 days ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day every 24 hours, every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: skin reddening ■ blisters ■ rash ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Location for lot number and expiration date

1Y974 FA C6-T 060425



Safety Feature – Do not use if printed tablet blister unit is open or torn.

NDC XXXX-XXX-XX



Treats FREQUENT Heartburn!

24 HR

Compare to Prilosec OTC*

DISSOLVE TABS

Omeprazole
Delayed Release Orally Disintegrating Tablets 20 mg
Acid Reducer



Strawberry Flavor

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Actual size

EFB 1369

FORMAT: STANDARD

TYPE SIZES:

TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed

- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscarmellose, ferrous oxide, flavor, hypromellose, hypromellose phthalate, maize methylcellulose, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days ■ do not take for more than 14 months unless directed by a doctor

4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

Drug Facts (continued)

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: rash ■ skin redness ■ blisters ■ rash ■ if an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with **lightheadedness, sweating or dizziness** ■ chest or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent heartburn ■ frequent heartburn ■ frequent heartburn ■ frequent heartburn

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. ■ you get diarrhea ■ you develop a rash or joint pain

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts ■ **Active ingredient (in each tablet)** Omeprazole 20 mg. ■ **Purpose** Acid reducer

Location for lot number and expiration date

1Y974 FA C6-TA 061825



Safety Feature –
Do not use if printed tablet blister unit is open or torn.

NDC XXXX-XXX-XX



Treats FREQUENT Heartburn!

24 HR

Compare to Prilosec OTC*
FDA Approved

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer



Strawberry Flavor

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Actual size

EFB 1369

FORMAT: STANDARD

TYPE SIZES:

TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed

- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, ferrous oxide, flavor, hypromellose, hypromellose phthalate, maize methylcellulose, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ do not take more than 1 tablet a day ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days ■ do not take for more than 14 months unless directed by a doctor

4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

Drug Facts (continued) ■ **Purpose** Acid reducer ■ **Active ingredient (in each tablet)** Omeprazole 20 mg ■ **Use** ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings ■ **Do not use if you have:** ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with **lightheadedness, sweating or dizziness** ■ chest or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain ■ **Ask a doctor before use if you have:** ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ **Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you develop a rash or joint pain ■ you get diarrhea

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Location for lot number and expiration date

1Y974 FA C6-TB 061925



Safety Feature –
Do not use if printed tablet blister unit is open or torn.

NDC XXXX-XXX-XX



Treats FREQUENT Heartburn!

24 HR

Compare to Prilosec OTC*
Child-resistant

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer



Strawberry Flavor

14 TABLETS

One 14-day course of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Actual size

EFB 1369

FORMAT: STANDARD

TYPE SIZES:

TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the “Drug Facts” box

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed

- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscarmellose, ferrous oxide, flavor, hypromellose, hypromellose phthalate, maize methylcellulose, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days ■ do not take for more than 14 months unless directed by a doctor

4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Drug Facts (continued)

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash ■ an allergic reaction occurs; stop use and seek medical help right away. If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with **lightheadedness, sweating or dizziness** ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent heartburn ■ frequent chest pain ■ frequent chest pain ■ frequent chest pain ■ frequent chest pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. ■ you develop a rash or joint pain ■ you get diarrhea

Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Active ingredient (in each tablet) Omeprazole 20 mg; Acid reducer

Purpose Acid reducer

Location for lot number and expiration date

1Y974 FA C6-TK 061925



Safety Feature – Do not use if printed tablet blister unit is open or torn.

NDC XXXX-XXX-XX



Treats FREQUENT Heartburn!



Compare to Prilosec OTC*

DISSOLVE TABS

Omeprazole
Delayed Release Orally Disintegrating Tablets 20 mg
Acid Reducer




Strawberry Flavor

14 TABLETS


One 14-day course of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



EFB 1369

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the “Drug Facts” box



Location for lot number and expiration date

1Y930 FA C6-T 060425

NDC XXXXX-XXX-XX

Treats FREQUENT Heartburn!

24 HR



Compare to Prilosec OTC*

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer



28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



Location for lot number and expiration date

1Y930 FA C6-TA 061225

NDC XXXXX-XXX-XX

Treats FREQUENT Heartburn!

24 HR



Compare to Prilosec OTC*
FDA Approved

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer



Strawberry Flavor

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect


Melts in your mouth
Dissolves without water



Actual size

EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



Location for lot number and expiration date

1Y930 FA C6-TB 061225

NDC XXXXX-XXX-XX



Treats FREQUENT Heartburn!

24 HR

Compare to Prilosec OTC*
Child-resistant

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer



Strawberry Flavor

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect


Melts in your mouth
Dissolves without water



Actual size

EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



Location for lot number and expiration date

1Y930 FA C6-TK 061225

NDC XXXXX-XXX-XX



Compare to Prilosec OTC*

Treats FREQUENT Heartburn!

24 HR

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer



Strawberry Flavor

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect


Melts in your mouth
Dissolves without water



Actual size

EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



Location for lot number and expiration date

1Y930 FA C6-TC 061225

Drug Facts
Active ingredient (in each tablet) Omeprazole 20 mg; Acid reducer

Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 Allergy alert: ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin reddening ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin reddening ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin reddening

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with **lightheadedness, sweating or dizziness** ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief or symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age; ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize malto-dextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
 Melts In Your Mouth

Treats FREQUENT Heartburn! **24 HR**

Compare to Prilosec OTC*
Buy One 14 Count, Get One 14 Count Free!

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

28 TABLETS

Two 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size

EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.
 10 Hakidma Street, Yokneam 2069200, Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd

Drug Facts
Active ingredient (in each tablet) Omeprazole 20 mg; Acid reducer

Use ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 Allergy alert: ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ rash ■ blisters ■ rash
 Do not use if you are allergic to omeprazole.

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor before use if you have:
 These may be signs of a serious condition. See your doctor.
 ■ frequent chest pain
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions ■ for adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief or symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture
 ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize malto-dextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Location for lot number and expiration date

1Y930 FA C6-TD 061225



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

Treats FREQUENT Heartburn!

24 HR

Compare to Prilosec OTC®*
 100% More Free!
 14 Free Tablets!

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer



28 TABLETS

Two 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water



EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



Location for lot number and expiration date

1Y930 FA C6-TF 061225

Drug Facts
Active ingredient (in each tablet) Omeprazole 20 mg; Acid reducer

Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 Allergy alert: ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ rash ■ blisters ■ rash
 Do not use if you are allergic to omeprazole.

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Other information ■ read the directions and warnings before use ■ keep the carton, it contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize malto-dextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age; ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief or symptoms within 24 hours



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
 Melts In Your Mouth

Treats FREQUENT Heartburn!

24 HR

Compare to Prilosec OTC®*
Bonus Size!
14 Tablets Free

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

Strawberry Flavor

28 TABLETS

Two 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Actual size

EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



Location for lot number and expiration date

Drug Facts
Active ingredient (in each tablet) Omeprazole 20 mg; Acid reducer

Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 Allergy alert: ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ rash ■ blisters ■ rash
 Do not use if you are allergic to omeprazole.

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief or symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize malto-dextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
 Melts In Your Mouth

Treats FREQUENT Heartburn!

24 HR

Compare to Prilosec OTC*
Bonus Size!
14 Free

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

28 TABLETS

Two 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size

EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



Location for lot number and expiration date

1Y930 FA C6-TH 061225

NDC XXXXX-XXX-XX

Treats FREQUENT Heartburn!

24 HR

MELTech®
Melts In Your Mouth

Compare to Prilosec OTC*
Bonus!
+ 14 Free

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer



Strawberry Flavor

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water



Actual size

EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



Drug Facts (continued)

Active ingredient (in each tablet) Omeprazole 20 mg; Acid reducer

Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ rash ■ blisters ■ rash
 ■ do not use if you are allergic to omeprazole.
Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea ■ you develop a rash or joint pain
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Other information ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture
 ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize malto-dextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ children under 18 years of age; ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg

1Y930 FA C6-TJ 061225

Location for lot number and expiration date



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
 Melts In Your Mouth

Compare to Prilosec OTC*
Bonus!
14 Tablets FREE

Treats FREQUENT Heartburn! **24 HR**

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

28 TABLETS

Two 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size

EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

10 Hakidma Street, Yokneam 2069200, Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 Made in Israel

5:00 PM EST
Questions or comments? 1-800-719-9260; weekdays 7:30 AM to

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

If pregnant or breast-feeding, ask a health professional before use. ■ you get diarrhea ■ you develop a rash or joint pain ■ you need to take more than 1 course of treatment every 4 months ■ you need to take this product for more than 14 days ■ your heartburn continues or worsens ■ you need to ask a doctor if:

Stop use and ask a doctor if:
 Acid reducers may interact with certain prescription drugs.

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ frequent weight loss ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with **lightheadedness, sweating or dizziness** ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Warnings:
 ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ rash ■ skin redness ■ blisters ■ rash ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ do not use if you are allergic to omeprazole. ■ Allergy alert:

Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Active ingredient (in each tablet) Omeprazole 20 mg; Acid reducer

Drug Facts (continued)

Directions ■ For adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief or symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age; ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopollose, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize malto-dextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Location for lot number and expiration date


1Y930 FA C6-TM 061225

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature – Do not use if printed tablet blister unit is open or torn.



MELTech®
Melts In Your Mouth

Compare to Prilosec OTC®*
Value 2 Pack

Treats **FREQUENT** Heartburn!

24 HR

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

28 TABLETS

Two 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water



Strawberry Flavor



Actual size

EFB 1393

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-T 060425

Manufactured in Israel
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating, or pain spreading to arms, neck or shoulders, or lightheadedness Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients ■ read the directions and warnings before use ■ alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST</p>
--	--

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
 Melts in Your Mouth

Compare to Prilosec OTC®*

Treats FREQUENT Heartburn!

24 HR

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

Strawberry Flavor

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-TA 061325

Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ rash ■ skin redness ■ blisters ■ hives If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. ■ do not crush or chew tablets The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients ■ read the directions and warnings before use alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>
---	--

DISSOLVE TABS
Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
Melts in Your Mouth

Compare to Prilosec OTC®*
FDA Approved

Treats FREQUENT Heartburn! **24 HR**

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water

Strawberry Flavor

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-TB 061325

Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. ■ do not crush or chew tablets The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>
---	---

DISSOLVE TABS
Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
Melts in Your Mouth

Compare to Prilosec OTC®*
Child-resistant

Treats FREQUENT Heartburn! **24 HR**

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water

Strawberry Flavor

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-TK 061325

Manufactured in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. ■ do not crush or chew tablets The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST</p>
---	--

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
 Melts in Your Mouth

Compare to Prilosec OTC®*

Treats FREQUENT Heartburn! **24 HR**

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Orally Disintegrating Tablets

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-TE 061325

Manufactured in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.



<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ rash ■ skin redness ■ blisters ■ hives If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders, or lightheadedness Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. ■ do not crush or chew tablets The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>
---	--

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.



NDC XXXXX-XXX-XX

MELTech®
 Melts in Your Mouth

Compare to Prilosec OTC®*
Bonus Pack: 28 + 14 Free

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-TF 061325

Manufactured in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.



<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders, or lightheadedness requent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ do not crush or chew tablets ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>
---	--

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.



NDC XXXXX-XXX-XX

MELTech®
 Melts in Your Mouth

Compare to Prilosec OTC®*
Bonus Size!
14 Tablets Free

24 HR

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-TG 061325

Manufactured in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin redness If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders, or lightheadedness Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ do not crush or chew tablets ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients ■ read the directions and warnings before use ■ alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>
---	--

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
 Melts in Your Mouth

Compare to Prilosec OTC®*
Bonus Size!
14 Free

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-TH 061325

Manufactured in Israel
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.



<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs <i>2 or more</i> days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ rash ■ skin redness ■ blisters ■ If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating, or pain spreading to arms, neck or shoulders, or lightheadedness Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. ■ do not crush or chew tablets The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients ■ read the directions and warnings before use ■ alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>
---	--

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.



NDC XXXXX-XXX-XX

MELTech®
 Melts in Your Mouth

Compare to Prilosec OTC®*
Bonus!
+ 14 Free

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-TJ 061325

Manufactured in Israel
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin redness If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders, or lightheadedness Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ do not crush or chew tablets ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>
---	--

DISSOLVE TABS
Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
 Melts in Your Mouth

Compare to Prilosec OTC®*
Bonus!
14 Tablets FREE

DISSOLVE TABS

Omeprazole
 Delayed Release Orally
 Disintegrating Tablets 20 mg
 Acid Reducer

42 TABLETS

Three 14-day courses of treatment
 May take 1 to 4 days for full effect

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Location for lot number and expiration date

1Y955 FA C6-TL 061325

Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg.....Acid reducer</p> <p>Use ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ do not use if you are allergic to omeprazole ■ omeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued)</p> <p>Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. ■ do not crush or chew tablets The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.</p> <p>Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolvidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST</p>
---	---

DISSOLVE TABS
Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg



- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

Safety Feature –
Do not use if printed tablet blister unit is open or torn.

NDC XXXXX-XXX-XX

MELTech®
Melts in Your Mouth

Compare to Prilosec OTC®*
Value 3 Pack

Treats FREQUENT Heartburn! **24 HR**

DISSOLVE TABS

Omeprazole
Delayed Release Orally
Disintegrating Tablets 20 mg
Acid Reducer

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water

Strawberry Flavor

Actual size

EFB 1945

FORMAT: STANDARD	
TYPE SIZES:	
TITLE	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued)	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS	6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT	6 pt Helvetica Neue LT Std 57 Condensed
LEADING	6.5 pt
BULLETS	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
TELEPHONE #	6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES	1.5 pt
HAIRLINES	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
03/11/2026 09:34:59 AM