















Do not use if seal is broken or damaged



pyrukynd®
(mitapivat) tablets

50 mg and 20 mg TAPER PACK

14 tablets

Contains two 7-day blister wallets with 7 tablets per wallet;

7 film-coated tablets of 50 mg per tablet (one tablet per day during week 1)
7 film-coated tablets of 20 mg per tablet (one tablet per day during week 2)

7 film-coated tablets of 20 mg per tablet (one tablet per day during week 2)

Rx Only

Week 1

50 mg per tablet

Week 2

20 mg per tablet

Swallow tablets whole.
Do Not split, crush, chew, or dissolve the tablets.



[imitapivat) tablets

50 mg and 20 mg TAPER PACK



pyrantel
(mitapivat) tablets

Each film-coated tablet contains:
50 mg mitapivat, provided as 58.5
20 mg mitapivat, provided as 23.4

Keep out of the reach of children.

Storage: Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].

Recommended Dosage: See prescribing information.

For more information, visit www.pyrukynd.com or call 1-833-228-8474

Manufactured for and distributed by:
Agios Pharmaceuticals, Inc.
Cambridge, MA 02139

Product of Switzerland

© 2021 Agios Pharmaceuticals, Inc.
PYRUYND® is a registered trademark of Agios Pharmaceuticals, Inc.

2D
CODE

[imitapivat) tablets

50 mg and 20 mg TAPER PACK





David
Lewis

Digitally signed by David Lewis

Date: 4/08/2026 09:05:30AM

GUID: 508da72000029f287fa31e664741b577