

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use INLURIYO safely and effectively. See full prescribing information for INLURIYO.

INLURIYO (imlunestrant) tablets, for oral use

Initial U.S. Approval: 2025

RECENT MAJOR CHANGES

Indication and Usage (1) 09/2026

INDICATIONS AND USAGE

INLURIYO® is an estrogen receptor antagonist indicated:

- as monotherapy for the treatment of adults with ER-positive, HER2-negative, *ESR1*-mutated advanced or metastatic breast cancer, as detected by an FDA-authorized test, with disease progression following at least one line of endocrine therapy (1)
- in combination with abemaciclib for the treatment of adults with ER-positive, HER2-negative, *ESR1*-mutated advanced or metastatic breast cancer, as detected by an FDA-authorized test, with disease progression following at least one line of endocrine therapy (1)

DOSAGE AND ADMINISTRATION

- Select patients for treatment based on the presence of *ESR1* mutations. (2.1)
- The recommended dosage is 400 mg orally once daily, on an empty stomach at least 2 hours before food or 1 hour after food. (2.2)
- See Full Prescribing Information for dosage modifications. (2.3, 2.4, 2.5)
- For patients with moderate or severe hepatic impairment, the recommended dosage is 200 mg orally once daily. (2.4, 8.6)

DOSAGE FORMS AND STRENGTHS

Tablets: 200 mg of imlunestrant. (3)

CONTRAINDICATIONS

- None. (4)

WARNINGS AND PRECAUTIONS

- **Embryo-Fetal Toxicity:** INLURIYO can cause fetal harm. Advise of the potential risk to a fetus and to use effective contraception. (5.1, 8.1, 8.3)

ADVERSE REACTIONS

The most common (incidence $\geq 10\%$) adverse reactions, including laboratory abnormalities, for INLURIYO monotherapy were: decreased hemoglobin, musculoskeletal pain, decreased calcium, decreased neutrophils, increased AST, fatigue, diarrhea, increased ALT, increased triglycerides, nausea, decreased platelets, constipation, increased cholesterol, and abdominal pain. (6.1)

The most common (incidence $\geq 10\%$) adverse reactions, including laboratory abnormalities, for INLURIYO in combination with abemaciclib were decreased neutrophils, diarrhea, decreased hemoglobin, decreased lymphocytes, nausea, decreased platelets, fatigue, increased triglycerides, infections, increased AST, increased creatinine, increased ALT, vomiting, musculoskeletal pain, abdominal pain, decreased appetite, increased cholesterol, rash, cough, headache, and decreased weight. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Eli Lilly and Company at 1-800-LillyRx (1-800-545-5979) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- **Strong CYP3A Inhibitors:** Avoid concomitant use with INLURIYO. If concomitant use cannot be avoided, decrease the INLURIYO dosage. (2.5, 7.1)
- **Strong CYP3A Inducers:** Avoid concomitant use with INLURIYO. If concomitant use cannot be avoided, increase the INLURIYO dosage. (2.5, 7.1)

USE IN SPECIFIC POPULATIONS

- **Lactation:** Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 09/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
 - 2.1 Patient Selection
 - 2.2 Recommended Dosage and Administration
 - 2.3 Dosage Modifications for Adverse Reactions
 - 2.4 Dosage in Patients with Hepatic Impairment
 - 2.5 Dosage Modifications for Drug Interactions
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
 - 5.1 Embryo-Fetal Toxicity
- 6 ADVERSE REACTIONS
 - 6.1 Clinical Trial Experience
- 7 DRUG INTERACTIONS
 - 7.1 Effects of Other Drugs on INLURIYO
 - 7.2 Effects of INLURIYO on Other Drugs
- 8 USE IN SPECIFIC POPULATIONS
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.3 Females and Males of Reproductive Potential
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 8.6 Hepatic Impairment

- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
 - 13.2 Animal Toxicology and/or Pharmacology
- 14 CLINICAL STUDIES
- 16 HOW SUPPLIED/STORAGE AND HANDLING
- 17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

INLURIYO is indicated:

- as monotherapy, for the treatment of adults with estrogen receptor (ER)-positive, human epidermal growth factor receptor 2 (HER2)-negative, *estrogen receptor-1 (ESR1)*-mutated advanced or metastatic breast cancer, as detected by an FDA-authorized test, with disease progression following at least one line of endocrine therapy [see *Dosage and Administration (2.1)*].
- in combination with abemaciclib for the treatment of adults with ER-positive, HER2-negative, *ESR1*-mutated advanced or metastatic breast cancer, as detected by an FDA-authorized test, with disease progression following at least one line of endocrine therapy [see *Dosage and Administration (2.1)*].

2 DOSAGE AND ADMINISTRATION

2.1 Patient Selection

Select patients for treatment of ER-positive, HER2-negative advanced or metastatic breast cancer with INLURIYO based on the presence of *ESR1* mutation(s) in a plasma specimen using an FDA-authorized test [see *Indications and Usage (1) and Clinical Studies (14)*].

Information on FDA-authorized tests for the detection of *ESR1* mutations in breast cancer is available at: <https://www.fda.gov/CompanionDiagnostics>.

2.2 Recommended Dosage and Administration

The recommended dosage of INLURIYO, as monotherapy or in combination with abemaciclib, is 400 mg orally once daily until disease progression or unacceptable toxicity.

Take on an empty stomach at least 2 hours before food or 1 hour after food [see *Clinical Pharmacology (12.3)*]. Take INLURIYO tablets at approximately the same time daily. Swallow the tablets whole. Do not split, crush, or chew the tablets.

If patient misses a dose by 6 or more hours or vomits, instruct the patient to take the next dose the following day at its scheduled time.

Refer to the Prescribing Information for the recommended dosage of abemaciclib when used in combination with INLURIYO.

2.3 Dosage Modifications for Adverse Reactions

The recommended INLURIYO dosage modifications for adverse reactions are provided in Tables 1 and 2.

The recommended dose reduction is to 200 mg once daily.

Permanently discontinue INLURIYO in patients who are unable to tolerate 200 mg once daily.

Refer to the Prescribing Information for dosage modification guidelines and other relevant safety information for abemaciclib when used in combination with INLURIYO.

Table 1: INLURIYO Dosage Modifications - Adverse Reactions (except hepatotoxicity)

Grade	INLURIYO Dosage Modifications
Persistent or recurrent Grade 2 that does not resolve with maximal supportive measures within 7 days to baseline or Grade 1	Withhold until toxicity resolves to baseline or $\leq$ Grade 1. Resume INLURIYO at the same dose.
Grade 3 or 4 (except non-hepatic asymptomatic laboratory changes)	Withhold until toxicity resolves to baseline or $\leq$ Grade 1. Resume INLURIYO at next lower dose.

Table 2: INLURIYO Dosage Modification - Hepatotoxicity

Monitor ALT/AST during imlunestrant therapy as clinically indicated.	
Liver Transaminase	INLURIYO Dosage Modifications
Persistent or Recurrent: AST/ALT >3.0-5.0×ULN	Withhold until toxicity resolves to baseline or to >ULN-3.0×ULN. Resume INLURIYO at the same dose.
If AST/ALT at baseline is within the normal range: AST/ALT >5.0-20×ULN Or If AST/ALT at baseline is above ULN: AST/ALT ≥3 × baseline (if AST/ALT ≥1.5 × ULN at baseline) Or AST/ALT >8 × ULN (whichever is the lower threshold)	Withhold until toxicity resolves to baseline or to >ULN-3.0×ULN. Resume INLURIYO at next lower dose or discontinue if receiving 200 mg daily.
AST/ALT >20.0×ULN Or ALT or AST ≥ 3 × ULN concurrent with TBL ≥ 2 × ULN (if ALT or AST < 1.5 × ULN at baseline), in the absence of cholestasis Or ALT or AST ≥ 2 × baseline concurrent with TBL ≥ 2 × ULN (if ALT or AST ≥ 1.5 × ULN at baseline), in the absence of cholestasis	Permanently discontinue INLURIYO.

Abbreviation: ALT = alanine aminotransferase, AST = aspartate aminotransferase, TBL=total bilirubin, ULN = upper limit of normal.

2.4 Dosage in Patients with Hepatic Impairment

The recommended dosage of INLURIYO for patients with moderate (Child-Pugh B) or severe (Child-Pugh C) hepatic impairment is 200 mg once daily. Monitor for increased adverse reactions [see *Clinical Pharmacology* (12.3)].

Refer to the Prescribing Information for dosage modification for severe hepatic impairment for abemaciclib when used in combination with INLURIYO.

2.5 Dosage Modifications for Drug Interactions

Strong CYP3A Inhibitors

Avoid concomitant use with strong CYP3A inhibitors. If concomitant use cannot be avoided, decrease the INLURIYO dosage to 200 mg once daily [see *Drug Interactions* (7.1)].

Refer to the Prescribing Information for dosage modification for strong CYP3A Inhibitors for abemaciclib when used in combination with INLURIYO.

Strong CYP3A Inducers

Avoid concomitant use with strong CYP3A inducers. If concomitant use cannot be avoided, increase the INLURIYO dosage to 600 mg once daily [see *Drug Interactions* (7.1)].

3 DOSAGE FORMS AND STRENGTHS

INLURIYO tablets contain 200 mg imlunestrant and are white, film coated-capsule-shaped tablets, with “LILLY” on one side and “1717” and elongated 4-point starburst on the other side.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Embryo-Fetal Toxicity

Based on findings in animals and its mechanism of action, INLURIYO can cause fetal harm when administered to a pregnant woman. In an animal reproduction study, oral administration of imlunestrant to pregnant rats during the period of organogenesis led to embryo-fetal mortality and structural abnormalities at maternal exposures that were below the human exposure at the recommended dose based on AUC.

Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with INLURIYO and for 1 week after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with INLURIYO and for 1 week after the last dose [see *Use in Specific Populations (8.1, 8.3) and Clinical Pharmacology (12.1)*].

6 ADVERSE REACTIONS

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

ER+, HER2-, *ESR1*-mutated Advanced or Metastatic Breast Cancer

INLURIYO as Monotherapy

The safety of INLURIYO was evaluated in 651 patients with ER+, HER2- locally advanced or metastatic breast cancer previously treated with endocrine therapy with or without a prior CDK4/6 inhibitor in EMBER-3 [see *Clinical Studies (14)*]. Patients received INLURIYO 400 mg orally, once daily (n=327), or standard of care (n=324) consisting of either fulvestrant (n=292) or exemestane (n=32). Among patients who were treated with INLURIYO, the median duration of exposure was 5.6 months (range: 0.2 to 28.6 months) in EMBER-3.

Serious adverse reactions occurred in 10% of patients who received INLURIYO. Serious adverse reactions in > 1% of patients included pleural effusion (1.2%). Fatal adverse reactions occurred in 1.8% of patients who received INLURIYO, including cardiac arrest, acute myocardial infarction, right ventricular failure, hypovolemic shock, and upper gastrointestinal hemorrhage (each 0.3%).

Permanent treatment discontinuation of INLURIYO due to an adverse reaction occurred in 4.6% of patients. Adverse reactions which resulted in permanent discontinuation of INLURIYO included increased alanine aminotransferase (ALT) (0.9%), abdominal pain, fatigue, fractured sacrum, hepatotoxicity, neuropathy peripheral, and pyrexia (each 0.3%).

Dosage interruption of INLURIYO due to an adverse reaction occurred in 10% of patients. Adverse reactions which required dosage interruption in >0.5% were vomiting (1.5%); increased aspartate aminotransferase (AST), and COVID-19 (each 0.9%); and increased ALT, anemia, diarrhea, decreased neutrophil count, and pyrexia (each 0.6%).

Dose reductions of INLURIYO due to an adverse reaction occurred in 2.4% of patients. Adverse reactions which required dose reductions were increased AST (0.6%); and increased ALT, anemia, fatigue, interstitial lung disease, nausea, neutropenia, and vomiting (each 0.3%).

The most common (≥10%) adverse reactions, including laboratory abnormalities, were decreased hemoglobin, musculoskeletal pain, decreased calcium, decreased neutrophils, increased AST, fatigue, diarrhea, increased ALT, increased triglycerides, nausea, decreased platelets, constipation, increased cholesterol, and abdominal pain.

Table 3 summarizes the adverse reactions in EMBER-3.

Table 3: Adverse Reactions (≥10%) in Patients Who Received INLURIYO in EMBER-3

Adverse Reaction ^a	INLURIYO N=327		Fulvestrant or Exemestane N=324	
	All Grades %	Grades 3 or 4 %	All Grades %	Grades 3 or 4 %
Musculoskeletal Disorders				
Musculoskeletal Pain	30	3.7	29	1.9
General Disorders and Administration Site Conditions				
Fatigue ^b	23	0.3	14	0.6
Gastrointestinal Disorders				
Diarrhea	22	0.6	12	0.0
Nausea	17	0.3	13	0.0
Constipation	10	0	6	0.3
Abdominal pain ^b	10	0.3	6	0.6

^a Adverse reactions were graded using NCI CTCAE version 5.0.

^b Includes other related terms

Clinically relevant adverse reactions (<10%) in patients who received INLURIYO included: vomiting (9%), headache (9%), cough (9%), decreased appetite (8%), hot flush (7%), pruritus (3.7%), dyspepsia (2.8%), and stomatitis (2.4%).

Table 4 summarizes the laboratory abnormalities in EMBER-3.

Table 4: Select Laboratory Abnormalities (≥10%) That Worsened from Baseline in Patients Who Received INLURIYO in EMBER-3

Lab Abnormality ^a	INLURIYO ^b		Fulvestrant or Exemestane ^b	
	All Grades %	Grades 3 or 4 %	All Grades %	Grades 3 or 4 %
Hematology				
Hemoglobin decreased	30	1.2	35	3.4
Neutrophils decreased	26	4	29	4.7
Platelets decreased	16	1.8	14	1.3
Chemistry				
Calcium decreased	26	0	19	0.6
AST increased	25	1.9	27	2.3
ALT increased	21	1.3	23	1.0
Triglycerides increased	21	0	22	1.2
Cholesterol increased	10	0	12	0

^a Graded according to NCI CTCAE version 5.0

^b The denominator used to calculate the rate varied from 252 to 325 based on the number of patients with a baseline value and at least one post-treatment value.

INLURIYO in Combination with Abemaciclib

The safety of INLURIYO in combination with abemaciclib was evaluated in 535 patients with ER+, HER2- locally advanced or metastatic breast cancer previously treated with endocrine therapy with or without a prior CDK4/6 inhibitor in EMBER-3 [see *Clinical Studies (14)*]. Patients received INLURIYO 400 mg orally once daily in combination with abemaciclib 150 mg twice daily (N=208), or INLURIYO monotherapy 400 mg, once daily (n=327). Among patients who were treated with INLURIYO in combination with abemaciclib, the median duration of exposure was 7.7 months (range 1 day to 25.3 months).

Serious adverse reactions occurred in 21% of patients who received INLURIYO in combination with abemaciclib. Serious adverse reactions in >1% of patients included pneumonia (2.4%), abdominal pain, and renal failure (each 1.4%). Fatal adverse reactions occurred in 3.8% of patients who received INLURIYO in combination with abemaciclib, including pneumonia (1.4%), myocardial infarction, interstitial lung disease, and sepsis (0.5% each).

Permanent treatment discontinuation of INLURIYO due to an adverse reaction occurred in two patients (1%). The adverse reactions which required permanent discontinuation were increased ALT and anemia (0.5% each).

Dosage interruptions of INLURIYO due to an adverse reaction occurred in 50% of patients. Adverse reactions which required dosage interruptions of INLURIYO in ≥2% of patients included diarrhea (14%), neutropenia (9%), decreased neutrophil count (6%), nausea (6%), anemia (6%), increased ALT (2.4%), and vomiting (2.4%).

Dose reductions of INLURIYO due to an adverse reaction occurred in 18% of patients. Adverse reactions which required dose reductions of INLURIYO in ≥ 1% of patients included diarrhea (6%), neutropenia (1.9%), nausea (1.9%), increased ALT (1.9%), and decreased neutrophil count (1.4%).

The most common (≥10%) adverse reactions, including laboratory abnormalities, were decreased neutrophils, diarrhea, decreased hemoglobin, decreased lymphocytes, nausea, decreased platelets, fatigue, increased triglycerides, infections, increased AST, increased creatinine, increased ALT, vomiting, musculoskeletal pain, abdominal pain, decreased appetite, increased cholesterol, rash, cough, headache, and decreased weight.

Table 5 summarizes the adverse reactions in EMBER-3.

Table 5: Adverse Reactions (≥10%) in Patients Who Received INLURIYO in Combination with Abemaciclib in EMBER-3

Adverse Reaction ^a	INLURIYO plus Abemaciclib N=208		INLURIYO N=327	
	All Grades %	Grades 3 or 4 %	All Grades %	Grades 3 or 4 %
Gastrointestinal Disorders				
Diarrhea	86	9	23	0.3
Nausea	51	1.9	18	0.3
Vomiting	32	0.5	10	0.6
Abdominal pain ^b	24	1.9	11	0.3
General Disorders and Administration Site Conditions				
Fatigue ^b	41	5	24	0.3
Infections and Infestations				
Infections ^b	36	7	25	3.4
Musculoskeletal Disorders				
Musculoskeletal pain ^b	25	1.4	31	2.8
Metabolism and Nutrition Disorders				
Decreased appetite ^b	22	1	9	0.3
Skin and Subcutaneous Tissue Disorders				
Rash ^b	14	1.4	4.9	0
Respiratory, Thoracic and Mediastinal Disorders				
Cough ^b	13	0	9	0
Nervous System Disorders				
Headache ^b	10	0.5	9	0
Investigations				
Weight decreased	10	0	4.6	0

^a Adverse reactions were graded using NCI CTCAE version 5.0.

^b Includes other related terms

Clinically relevant adverse reactions (<10%) in patients who received INLURIYO in combination with abemaciclib included: constipation (9%), stomatitis (9%), fever (9%), urinary tract infection (8%), dizziness (8%), hot flush (6%), insomnia (6%), arrhythmia (6%), alopecia (5%), back pain (4.8%), venous thromboembolic event (4.8%) and interstitial lung disease or pneumonitis (2.9%).

Table 6 summarizes the laboratory abnormalities in EMBER-3.

Table 6: Select Laboratory Abnormalities (≥10%) That Worsened from Baseline in Patients Who Received INLURIYO in Combination with Abemaciclib in EMBER-3

Lab Abnormality ^a	INLURIYO plus Abemaciclib ^b		INLURIYO ^b	
	All Grades %	Grades 3 or 4 %	All Grades %	Grades 3 or 4 %
Hematology				
Neutrophils decreased	86	21	28	4.3
Hemoglobin decreased	81	9	34	2.2
Lymphocytes decreased	58	9	29	2.2
Platelets decreased	48	2.4	18	2.8
Chemistry				
Triglycerides increased	40	2.3	22	0
AST increased	36	2.5	27	1.9
Creatinine increased	36	1.1	6	0.3
ALT increased	33	5	22	1.3
Cholesterol increased	18	0	12	0.4

^a Graded according to NCI CTCAE version 5.

^b The denominator used to calculate the frequency varied from 178 to 205 based on the number of patients with a baseline value and at least one post-treatment value.

7 DRUG INTERACTIONS

7.1 Effects of Other Drugs on INLURIYO

Strong CYP3A Inhibitors

Avoid concomitant use of INLURIYO with strong CYP3A inhibitors. If concomitant use cannot be avoided, reduce the dosage of INLURIYO [see *Dosage and Administration (2.5)*, *Clinical Pharmacology (12.3)*].

Imlunestrant is a CYP3A substrate. Concomitant use of a strong CYP3A inhibitor increases imlunestrant exposure [see *Clinical Pharmacology (12.3)*], which may increase the risk of INLURIYO-associated adverse reactions.

Strong CYP3A Inducers

Avoid concomitant use of INLURIYO with strong CYP3A inducers. If concomitant use cannot be avoided, increase the dosage of INLURIYO [see *Dosage and Administration (2.5)*, *Clinical Pharmacology (12.3)*].

Imlunestrant is a CYP3A substrate. Concomitant use of a strong CYP3A inducer decreases imlunestrant exposure [see *Clinical Pharmacology (12.3)*], which may reduce effectiveness of INLURIYO.

7.2 Effects of INLURIYO on Other Drugs

P-gp or BCRP Substrates

Avoid concomitant use unless otherwise recommended in the Prescribing Information for P-gp or BCRP substrates where minimal concentration changes may lead to serious adverse reactions.

Imlunestrant inhibits both P-gp and BCRP. Imlunestrant increases exposure of P-gp and BCRP substrates, which may increase the risk of adverse reactions related to these substrates [see *Clinical Pharmacology (12.3)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Refer to the Prescribing Information of abemaciclib when administered in combination with INLURIYO for pregnancy information.

Based on findings in animals and its mechanism of action, INLURIYO can cause fetal harm when administered to a pregnant woman [see *Clinical Pharmacology* (12.1)]. There are no available human data on INLURIYO use in pregnant women to inform the drug-associated risk. In an animal reproduction study, oral administration of imlunestrant to pregnant rats during the period of organogenesis led to embryo-fetal mortality and structural abnormalities at maternal exposures below the human exposure at the recommended dose based on AUC (see *Data*). Advise pregnant women and females of reproductive potential of the potential risk to a fetus.

The background risk of major birth defects and miscarriage for the indicated population is unknown. However, the background risk in the U.S. general population of major birth defects is 2 to 4% and of miscarriage is 15 to 20% of clinically recognized pregnancies.

Data

Animal Data

In an embryo-fetal development study, imlunestrant was administered orally to pregnant rats during the period of organogenesis from gestation day 6 to 17 at doses of 0.3, 3, and 30 mg/kg/day. Imlunestrant caused embryo-fetal mortality (increased resorption, reduced number of live fetuses) at ≥ 0.3 mg/kg/day, approximately 0.1 times the human AUC at the recommended dose. Early delivery, fetal malformations (including small jaw, protruding tongue, malrotated, and hyperextended hindlimb) and fetal variations (edema localized subcutis) were observed at ≥ 3 mg/kg/day, approximately 1 time the human AUC at the recommended dose.

8.2 Lactation

Refer to the Prescribing Information of abemaciclib when administered in combination with INLURIYO for lactation information. When used in combination, advise patients to not breastfeed during treatment and for the longest post-treatment duration recommended in the Prescribing Information of the individual products.

Risk Summary

There are no data on the presence of imlunestrant or its metabolites in human milk, its effects on the breastfed child, or on milk production. Because of the potential for serious adverse reactions in the breastfed child, advise lactating women to not breastfeed during treatment with INLURIYO and for 1 week after the last dose.

8.3 Females and Males of Reproductive Potential

Refer to the Prescribing Information of abemaciclib when administered in combination with INLURIYO for contraception and infertility information. When used in combination, advise patients to use effective contraception during treatment and for the longest post-treatment duration recommended in the Prescribing Information of the individual products.

INLURIYO can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations* (8.1)].

Pregnancy Testing

Verify the pregnancy status in females of reproductive potential prior to initiating INLURIYO.

Contraception

Females

Advise females of reproductive potential to use effective contraception during treatment with INLURIYO and for 1 week after the last dose.

Males

Advise male patients with female partners of reproductive potential to use effective contraception during treatment with INLURIYO and for 1 week after the last dose.

Infertility

Based on findings in animals, INLURIYO may impair fertility in females and males of reproductive potential. Findings in animals were reversible [see *Nonclinical Toxicology* (13.1)].

8.4 Pediatric Use

The safety and effectiveness of INLURIYO have not been established in pediatric patients.

8.5 Geriatric Use

Of 327 patients who received INLURIYO monotherapy in the EMBER-3 study, 118 patients were ≥ 65 years of age and 37 patients were ≥ 75 years of age. No overall differences in safety or effectiveness of INLURIYO monotherapy have been observed between patients 65 years of age and older and younger adult patients.


Of the 208 patients who received INLURIYO in combination with abemaciclib in the EMBER-3 study, 90 patients were ≥ 65 years of age and 27 patients were ≥ 75 years of age. There was a higher rate of nausea in patients aged 65 years or older (59%) compared with younger patients (44%). The EMBER-3 study did not include sufficient numbers of patients 65 years of age and older with *ESR1* mutations treated with INLURIYO in combination with abemaciclib to determine whether they respond differently than younger patients.

8.6 Hepatic Impairment

Reduce the dose of INLURIYO in patients with moderate (Child-Pugh B) or severe (Child-Pugh C) hepatic impairment. No dosage modification is recommended for patients with mild hepatic impairment (Child-Pugh A) [see *Dosage and Administration* (2.4) and *Clinical Pharmacology* (12.3)].

11 DESCRIPTION

INLURIYO tablets contain imlunestrant, an estrogen receptor antagonist. The chemical name for imlunestrant tosylate is (5R)-5-(4-(2-(3-(fluoromethyl)azetidin-1-yl)ethoxy)phenyl)-8-(trifluoromethyl)-5H-(1)benzopyrano(4,3-c)quinolin-2-ol, tosylate salt (1:1). Imlunestrant tosylate is a white to practically white to yellow powder with the empirical formula $C_{29}H_{24}F_4N_2O_3 \cdot C_7H_8O_3S$ and a molecular weight 696.71 g/mol. The aqueous solubility of imlunestrant tosylate is slightly soluble at low pH, insoluble at neutral pH, and sparingly soluble at high pH. The chemical structure of imlunestrant tosylate is shown below:



INLURIYO tablets are for oral administration. Each INLURIYO tablet is available as capsule-shaped, film-coated tablet that contains 200 mg imlunestrant (equivalent to 265.66 mg imlunestrant tosylate). The tablet contains the following inactive ingredients: croscarmellose sodium, hydroxypropyl cellulose, magnesium stearate, and microcrystalline cellulose. The tablets are coated using a common white coating, which consists of polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Imlunestrant is an estrogen receptor (ER) antagonist that binds to ER α . In vitro, imlunestrant induced degradation of ER α , leading to inhibition of ER-dependent gene transcription and cellular proliferation in ER+ breast cancer cells. Imlunestrant demonstrated in vitro and in vivo anti-tumor activity in ER+ breast cancer xenograft models, including models with *ESR1* mutations.

12.2 Pharmacodynamics

Imlunestrant exposure-response relationships and the time course of pharmacodynamics have not been fully characterized.

Cardiac Electrophysiology

At 2 times the mean maximum concentration observed with the approved recommended dose, a mean increase in the QTc interval >20 msec was not observed.

12.3 Pharmacokinetics

Imlunestrant pharmacokinetics were observed at steady state at the approved recommended dosage and are presented as mean (%CV) unless otherwise specified. The maximum concentration ($C_{\max}$) of imlunestrant is 141 ng/mL (45%) and the area under the concentration-time curve (AUC) is 2,400 ng·h/mL (46%). Imlunestrant $C_{\max}$ and AUC increase in a dose proportional manner over a dosage range of 200 mg to 1,200 mg (0.5 to 3 times the approved recommended dosage) once daily. Steady-state is reached in approximately 6 days and the accumulation is 2.3-fold based on AUC.

Absorption

Imlunestrant absolute oral bioavailability after a single oral 400 mg dose is 10% (32%). Imlunestrant median (min, max) time to maximum plasma concentration ($T_{\max}$) is 4 (2, 8) hours.

Effect of Food

Imlunestrant AUC increased 2-fold and $C_{\max}$ increased 3.6-fold following administration with a low-fat meal (approximately 475 calories with 13% fat, 16% protein, and 71% carbohydrates). The effect of high-fat meal (approximately 800-1,000 calories with 500-600 calories from fat) on imlunestrant exposures is unknown.

Distribution

The apparent (oral) volume of distribution is 8,120 L (69%). Imlunestrant protein binding is >99% and is not concentration dependent.

Elimination

Imlunestrant elimination half-life is 30 hours with an estimated apparent clearance of 166 L/h (51%).

Metabolism

Imlunestrant is metabolized by sulfation, CYP3A4, and direct glucuronidation (UGT1A1, 1A3, 1A8, 1A9, 1A10).

Excretion

After a single dose of radiolabeled imlunestrant 400 mg to healthy subjects, 97% of the dose was recovered in feces (62% unchanged) and 0.3% in urine.

Specific Populations

No clinically significant differences in the pharmacokinetics of imlunestrant based on age (28 to 95 years), race (64% White, 23% Asian, and 5% Black or African American), ethnicity (74% non-Hispanic/Latino, 17% Hispanic/Latino), body weight (36 to 145 kg), mild to moderate (eGFR 30 to 89 mL/min, estimated by CKD-EPI equation) renal impairment, or UGT1A1 genetic polymorphisms (e.g., UGT1A1*1/*28 or UGT1A1*28/*28). The effect of severe (eGFR 15 to 29 mL/min) renal impairment and renal impairment requiring dialysis on imlunestrant pharmacokinetics is unknown.

Patients with Hepatic Impairment

Imlunestrant AUC increased 2.2-fold in subjects with moderate hepatic impairment (Child-Pugh B) and 3.1-fold in subjects with severe hepatic impairment (Child-Pugh C). No clinically significant differences in the pharmacokinetics of imlunestrant were observed in subjects with mild hepatic impairment (Child-Pugh A).

Drug Interaction Studies

Clinical Studies

Strong CYP3A Inhibitors: Imlunestrant AUC increased 2.1-fold and $C_{\max}$ increased 1.9-fold following concomitant use of itraconazole (strong CYP3A inhibitor) for multiple days.

Strong CYP3A Inducers: Imlunestrant AUC decreased by 42% and $C_{\max}$ decreased by 29% following concomitant use of carbamazepine (strong CYP3A inducer) for multiple days.

P-gp Substrates: Digoxin (P-gp substrate) AUC increased 1.4-fold and $C_{\max}$ increased 1.6-fold following concomitant use of imlunestrant.

BCRP Substrates: Rosuvastatin (BCRP substrate) AUC increased 1.5-fold and $C_{\max}$ increased 1.6-fold following concomitant use with imlunestrant.

Other Drugs: No clinically significant differences in the pharmacokinetics of imlunestrant were observed when used concomitantly with omeprazole (gastric acid-reducing agent) or quinidine (P-gp inhibitor).

No clinically significant differences in the pharmacokinetics of midazolam (CYP3A substrate), repaglinide (CYP2C8 substrate), omeprazole (CYP2C19 substrate), or dextromethorphan (CYP2D6 substrate) were observed when used concomitantly with imlunestrant.

In Vitro Studies

CYP Enzymes: Imlunestrant is an inhibitor of CYP2B6 and CYP2C9 but is not an inhibitor of CYP1A2. Imlunestrant is not an inducer of CYP1A2, CYP2B6, or CYP2C9.

Transporter Systems: Imlunestrant is not a substrate of BCRP, OCT1, OATP1B1, or OATP1B3.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Carcinogenicity studies have not been conducted with imlunestrant.

Mutagenesis

Imlunestrant was not mutagenic in the bacterial reverse mutation (Ames) assay. Imlunestrant was clastogenic in an in vitro human lymphocyte micronucleus assay. Imlunestrant was not genotoxic in an in vivo rat bone marrow micronucleus test and did not induce DNA breaks in the liver and duodenum comet assays.

Impairment of Fertility

Fertility studies with imlunestrant in animals have not been conducted. In repeat-dose toxicity studies up to 6 months in rats and 3 months in cynomolgus monkeys, oral administration of imlunestrant resulted in follicular cysts in the ovary and atrophy in the vagina, cervix, and uterus at doses ≥ 10 mg/kg/day in rats (≥ 4 times the human AUC at the recommended dose) and ≥ 15 mg/kg/day in cynomolgus monkeys (≥ 1 times the human AUC at the recommended dose). Decreased sperm and cellular debris in the epididymis and spermatid retention in the testis were observed in male rats at ≥ 10 mg/kg/day. The effects of imlunestrant on male and female reproductive organs were reversible in rats following a 3-month recovery period. Reversibility was not assessed in cynomolgus monkeys.

13.2 Animal Toxicology and/or Pharmacology

In a 6-month repeat-dose toxicity study, oral administration of imlunestrant to rats resulted in epithelial hyperplasia in the urinary bladder, granulosa cell hyperplasia in the ovary, and hyperplasia in the mammary gland at doses ≥ 10 mg/kg/day (≥ 4 times the human AUC at the recommended dose). These effects, except urinary bladder effects, were reversible after a 3-month recovery period.

14 CLINICAL STUDIES

ER+, HER2-, *ESR1*-mutated Locally Advanced or Metastatic Breast Cancer

The efficacy of INLURIYO was evaluated in EMBER-3 (NCT04975308), a randomized, open-label, active-controlled, multicenter trial that enrolled 874 adult patients with ER+, HER2- locally advanced or metastatic breast cancer of which 323 patients had *ESR1* mutations (*ESR1m*), who were previously treated with an aromatase inhibitor either alone or in combination with a CDK4/6 inhibitor. Patients were excluded if they were eligible to receive a PARP inhibitor. Patients were required to have progressed:

- Within 12 months of completing neoadjuvant or adjuvant aromatase inhibitor therapy with no systemic treatment for recurrent disease or
- Greater than 12 months after neoadjuvant or adjuvant endocrine therapy or de novo metastatic disease and had progressed on only one line of aromatase inhibitor therapy.

Patients were randomized 1:1:1 to

- INLURIYO 400 mg orally once daily; or
- investigator's choice of endocrine therapy [fulvestrant 500 mg intramuscularly on days 1, 15, 29, and once monthly thereafter or exemestane 25 mg orally once daily]; or
- INLURIYO 400 mg once daily in combination with abemaciclib 150 mg orally twice daily.

Randomization was stratified by previous treatment with CDK4/6 inhibitor (yes vs no), presence of visceral metastasis (yes vs no), and region (East Asia vs North America/Western Europe vs Others). *ESR1m* status was determined by blood circulating tumor deoxyribonucleic acid (ctDNA) analysis using the Guardant360 CDx assay and was limited to specific *ESR1* mutations in the ligand binding domain. Patients were treated until disease progression or unacceptable toxicity.

INLURIYO as Monotherapy

The major efficacy outcomes for INLURIYO monotherapy were investigator-assessed progression-free survival (PFS) according to RECIST v1.1 in the overall population and in the population of patients whose tumors had an *ESR1m*. Other efficacy measures included overall survival (OS), blinded independent review committee (BIRC)-assessed PFS, and objective response rate (ORR). A statistically significant difference in PFS in the overall population was not observed; however, there was a statistically significant difference in PFS in the *ESR1m* population for INLURIYO monotherapy compared to investigator's choice of endocrine therapy (fulvestrant or exemestane) indicating the PFS benefit of INLURIYO monotherapy was limited to the *ESR1m* population.

Among the patients on the INLURIYO arm or investigator's choice of endocrine therapy who were positive for *ESR1m* (N=256), the median age was 61 years (range: 28-85 years); all patients were female, of which 11% were pre/perimenopausal; 61% were White, 26% Asian, 4% Black, 4% were American Indian or Alaskan Native, 4.7% missing, 0.8% multiple, 19% were Hispanic/Latino; and baseline ECOG performance status was 0 (63%) or 1 (37%). Most patients had visceral metastasis (59%) at baseline. Of the patients enrolled, 21% had received no endocrine therapy and 79% had received one line of endocrine therapy in the advanced or metastatic setting. Overall, 70% of patients were treated with a prior CDK4/6 inhibitor, 2.3% treated in the adjuvant setting and 67% treated in the advanced or metastatic setting.

Efficacy results for the *ESR1m* population are summarized in Table 7 and Figure 1. PFS assessment in the *ESR1m* population based on a BIRC was consistent with the investigator assessment. At the time of PFS analysis, overall survival data was immature with 31% of deaths in the *ESR1m* population.

Table 7: Efficacy Results for INLURIYO Monotherapy Compared with Fulvestrant/Exemestane in Patients with *ESR1m* in EMBER-3

	INLURIYO N=138	Fulvestrant or Exemestane N=118
Progression-free Survival (PFS) ^{a, b}		
Number of PFS Events, n (%)	109 (79)	102 (86)
Median in months (95% CI)	5.5 (3.9, 7.4)	3.8 (3.7, 5.5)
Hazard Ratio (95% CI) ^c	0.62 (0.46, 0.82)	
p-value ^d	0.0008	
Objective Response Rate ^b		
Patients with Measurable Disease	112	91
ORR (95% CI)	14% (8, 21)	8% (2, 13)
Complete response rate	0.9%	0
Partial response rate	13%	8%


^a Investigator Assessed

^b per RECIST v1.1

^c Based on the stratified Cox proportional hazard model

^d Two-sided p-value based on stratified log-rank test (compared to a significance level of 0.04)

Figure 1: Kaplan-Meier Plot of Investigator-Assessed PFS for Patients with *ESR1m*, Treated with INLURIYO Monotherapy or Fulvestrant/Exemestane in EMBER-3



INLURIYO in Combination with Abemaciclib

The major efficacy outcome for INLURIYO in combination with abemaciclib was investigator-assessed PFS according to RECIST v1.1 in the overall population. Other efficacy measures included OS, BIRC-assessed PFS, and ORR. While the comparison of PFS between INLURIYO in combination with abemaciclib and INLURIYO monotherapy in the overall population was statistically significant, a PFS improvement was not demonstrated for INLURIYO monotherapy compared to investigator's choice of endocrine therapy in either the overall or *ESR1m* not detected populations, indicating that the benefit for INLURIYO monotherapy was only observed in the *ESR1m* population. Therefore, exploratory analyses of PFS, OS and ORR in the *ESR1m* population comparing INLURIYO in combination with abemaciclib and INLURIYO monotherapy were performed.

Among the patients on the INLURIYO in combination with abemaciclib arm or the INLURIYO monotherapy arm who were positive for *ESR1m* (n=159), the median age was 62 years (range: 31-85 years); all patients were female, of which 12% were pre/perimenopausal; 60% were White, 27% Asian, 5% Black, 0.6% were American Indian or Alaskan Native, 0.6% multiple races, 7% missing, 12% were Hispanic/Latino, 75% not Hispanic/Latino, 13% not reported/missing; and all patients had a baseline ECOG performance status of 0 (66%) or 1 (34%). Most patients had visceral metastasis (58%) at baseline. Of the patients enrolled, all received a prior line of endocrine therapy and 81% had received one line of endocrine therapy in the advanced or metastatic setting. Overall, 79% of patients were treated with a prior CDK4/6 inhibitor, 4.4% were treated in the adjuvant setting and 74% were treated in the advanced or metastatic setting. Among the patients who had received prior CDK4/6 inhibitors, the specific CDK4/6 inhibitor received was palbociclib 65%, ribociclib 28%, and abemaciclib 6%.

The exploratory analyses results in the *ESR1m* population are summarized in Table 8 and Figure 2. PFS assessment in the *ESR1m* population based on a BIRC was consistent with the investigator assessment. At the time of interim analysis, overall survival data was immature with 35% of deaths in the *ESR1m* population.

Table 8: Efficacy Results for INLURIYO in Combination with Abemaciclib Compared with INLURIYO Monotherapy in Patients with *ESR1m* in EMBER-3

	INLURIYO + Abemaciclib N=67	INLURIYO N=92
Progression-free Survival (PFS) ^{a,b}		


Number of PFS Events, n (%)	36 (54)	71 (77)
Median in months (95% CI)	11.1 (7.4, 13.7)	5.5 (3.8, 7.2)
Hazard Ratio (95% CI) ^c	0.53 (0.35, 0.80)	
Objective Response Rate ^{a, b}		
Patients with Measurable Disease	54	74
ORR (95% CI)	35% (22, 48)	15% (7, 23)
Complete response rate	1.9%	1.4%
Partial response rate	33%	14%

^a Investigator Assessed

^b per RECIST v1.1

^c Based on the stratified Cox proportional hazard model

Figure 2: Kaplan-Meier Plot of Investigator-Assessed PFS for Patients with *ESR1m*, Treated with INLURIYO in Combination with Abemaciclib or INLURIYO Monotherapy in EMBER-3



16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

INLURIYO 200 mg tablets are white, film coated, capsule-shaped tablets with “LILLY” on one side and “1717” and an elongated 4-point starburst on the other side.

INLURIYO 200 mg tablets are supplied in either a 28-count or 56-count bottle configuration.

- 28-count: NDC 0002-1717-28
- 56-count: NDC 0002-1717-56

Storage and Handling

Store at 20°C to 25°C (68°F to 77°F). Excursions between 15°C to 30°C (59°F to 86°F) are permitted [see *USP Controlled Room Temperature*].

Disposal

Dispose unused medication via a take-back option if available. Otherwise, follow FDA instructions for disposing medication in the household trash, www.fda.gov/drugdisposal. Do NOT flush down the toilet.

17 PATIENT COUNSELING INFORMATION

Advise patients to read the FDA-approved patient labeling (Patient Information).

Dyslipidemia

Advise patients that hypercholesterolemia and hypertriglyceridemia may occur while taking INLURIYO. Inform patients that lipid profile monitoring will be performed prior to starting and periodically while taking INLURIYO [see *Adverse Reactions* (6.1)].

Embryo-Fetal Toxicity

Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females to inform their healthcare provider of a known or suspected pregnancy [see *Warnings and Precautions* (5.1) and *Use in Specific Populations* (8.1)].

Advise females of reproductive potential to use effective contraception during treatment with INLURIYO and for 1 week after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with INLURIYO and for 1 week after the last dose [see *Use in Specific Populations* (8.3)].

Refer to the Prescribing Information of abemaciclib when administered in combination with INLURIYO for pregnancy and contraception information. When used in combination, advise patients to use effective contraception during treatment and for the longest post-treatment duration recommended in the Prescribing Information of the individual products.

Lactation

Advise women not to breastfeed during treatment with INLURIYO and for 1 week after the last dose [see *Use in Specific Populations* (8.2)].

Refer to the Prescribing Information of abemaciclib when administered in combination with INLURIYO for lactation information. When used in combination, advise patients to not breastfeed during treatment and for the longest post-treatment duration recommended in the Prescribing Information of the individual products.

Infertility

Advise males and females of reproductive potential that INLURIYO may impair fertility [see *Use in Specific Populations* (8.3)].

Drug Interactions

Advise patients to inform their healthcare providers of all concomitant medications, including prescription medicines, over the counter drugs, and herbal products [see *Drug Interactions* (7.1, 7.2)].

Dosing Instructions

Instruct patients to take INLURIYO at approximately the same time each day and to swallow the tablet(s) whole. Tablets should not be chewed, crushed, or split prior to swallowing [see *Dosage and Administration* (2.2)].

Advise patients to take INLURIYO without food, either 2 hours before food or 1 hour after food [see *Dosage and Administration* (2.2)].

Instruct patient that if a dose of INLURIYO is missed by more than 6 hours or vomiting occurs, skip the dose and take the next dose the following day at its regularly scheduled time [see *Dosage and Administration* (2.2)].

Marketed by: Lilly USA, LLC, Indianapolis, IN 46285, USA

Copyright © 2025, YYYY, Eli Lilly and Company. All rights reserved.

Pat.: www.lilly.com/patents

A4.01-INL-000X-USPI-YYYYMMDD

PATIENT INFORMATION
INLURIYO® (en-loo-ree-yoh)
imlunestrant
tablets

What is INLURIYO?

INLURIYO is a prescription medicine that is used alone or in combination with the medicine abemaciclib to treat adults with estrogen receptor (ER)-positive, human epidermal growth factor receptor 2 (HER2)-negative, *estrogen receptor-1 (ESR1)*-mutated advanced breast cancer or breast cancer that has spread to other parts of the body (metastatic), **and** whose disease has progressed after at least 1 line of endocrine therapy.

Your healthcare provider will perform a test to make sure that INLURIYO is right for you.

When INLURIYO is used in combination with abemaciclib, read the Patient Information for abemaciclib.

It is not known if INLURIYO is safe and effective in children.

Before taking INLURIYO, tell your healthcare provider about all of your medical conditions including if you:

- have liver problems.
- are pregnant or plan to become pregnant. INLURIYO can harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with INLURIYO.
- Use effective birth control (contraception) during treatment with INLURIYO and for 1 week after the last dose.
- Your healthcare provider will tell you how long you should use birth control (contraception) after combination treatment with INLURIYO and abemaciclib.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with INLURIYO.

Males with female partners who are able to become pregnant:

- Use effective birth control (contraception) during treatment with INLURIYO and for 1 week after the last dose.
- are breastfeeding or plan to breastfeed. It is not known if INLURIYO passes into breastmilk.
 - Do not breastfeed during treatment with INLURIYO and for 1 week after the last dose.
 - Your healthcare provider will tell you how long you should not breastfeed after combination treatment with INLURIYO and abemaciclib.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. INLURIYO and other medicines may affect each other causing side effects. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

How should I take INLURIYO?

- Take INLURIYO exactly as your healthcare provider tells you.
- Do not stop taking INLURIYO or change the dose unless your healthcare provider tells you to.
- Take INLURIYO 1 time a day at about the same time each day.
- Take INLURIYO on an empty stomach, at least 2 hours before food or 1 hour after food.
 - Swallow INLURIYO tablets whole. Do not chew, crush, or split the tablets.
- If you miss a dose of INLURIYO by 6 or more hours or vomit after taking INLURIYO, do not take another dose on that day. Take your next dose the following day at your regularly scheduled time.

What are the possible side effects of INLURIYO?

The most common side effects of INLURIYO when taken alone include:

- decreased hemoglobin (anemia)
- muscle and joint pain
- decreased white blood cell counts
- increased liver function tests
- decreased calcium blood levels
- tiredness
- diarrhea
- increased triglyceride and cholesterol blood levels. Your healthcare provider will check your blood levels before and during treatment with INLURIYO.
- nausea
- decreased platelet counts
- constipation
- stomach-area (abdominal) pain

The most common side effects of INLURIYO when taken with abemaciclib include:

- decreased white blood cell counts
- diarrhea
- decreased hemoglobin (anemia)
- nausea
- decreased platelet counts
- tiredness
- increased triglycerides and cholesterol blood levels. Your healthcare provider will check your blood levels before and during treatment with INLURIYO.
- infections
- increased liver function tests
- increased creatinine blood levels
- vomiting
- muscle and joint pain
- stomach-area (abdominal) pain
- decreased appetite
- rash
- cough
- headache
- weight loss

Your healthcare provider may change your dose, temporarily stop, or completely stop treatment with INLURIYO if you develop certain side effects.

INLURIYO may affect fertility in males and in females who are able to become pregnant. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of INLURIYO.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store INLURIYO?

- Store INLURIYO at room temperature between 68°F to 77°F (20°C to 25°C).
- Throw away (dispose of) unused INLURIYO by a medicine take-back program if available. If no take-back program is available, visit www.fda.gov/drugdisposal for instructions on how to throw away medicines in the household trash.
- **Do not** flush INLURIYO down the toilet.

Keep INLURIYO and all medicines out of the reach of children.

General information about the safe and effective use of INLURIYO.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use INLURIYO for a condition for which it was not prescribed. Do not give INLURIYO to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for more information about INLURIYO.

What are the ingredients in INLURIYO?

Active ingredient: imlunestrant

Inactive ingredients: croscarmellose sodium, hydroxypropyl cellulose, magnesium stearate, and microcrystalline cellulose. The tablet coating contains polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide.

INLURIYO is a trademark of Eli Lilly and Company. All rights reserved.

Marketed by: Lilly USA, LLC, Indianapolis, IN 46285, USA

Copyright © 2025, YYYY, Eli Lilly and Company. All rights reserved.
For more information, go to www.INLURIYO.com or call 1-800-545-5979

This Patient Information has been approved by the U.S. Food and Drug Administration

Revised: 09/2026

A2.01-INL-000X-PPI-YYYYMMDD