


Drug Facts Format Information				
Drug Facts:	5.50 pt	Subheadings/Body Text:	4.50 pt	Barlines: 1.0 pt
Drug Facts (continued):	4.50 pt	Bullets:	3.50 pt	Horizontal Scale: 76%
Headings:	4.50 pt	Leading:	4.50 pt	Tracking: -20



Drug Facts Format Information					
Drug Facts:	5.50 pt	Subheadings/Body Text:	4.50 pt	Barlines:	1.0 pt
Drug Facts (continued):	4.50 pt	Bullets:	3.50 pt	Hairlines:	0.5 pt
Headings:	4.50 pt	Leading:	4.20/4.50 pt	Horizontal Scale:	76%
				Tracking:	-20





Net Content Statement

PDP Area: (width) x height

Net Content Size:

6.84 in²

0.125"

NET CONTENT

1/16 IN.

1/8 IN.

3/16 IN.

1/4 IN.

1/2 IN.

AREA OF PDP

5 SQ. IN. OR LESS

MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.

MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.

MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.

OVER 400 SQ. IN.

Drug Facts Format Information

Drug Facts: 9.00 pt | Subheadings/Body Text: 6.00 pt | Barlines: 1.5 pt | Horizontal Scale: 75% |

Drug Facts (continued): 8.00 pt | Bullets: 5.00 pt | Hairlines: 0.5 pt | Tracking: -25 |

Headings: 8.00 pt | Leading: 5.8/6.0 pt |



Net Content Statement	
PDP Area: (width x height)	3.08 In ²
Net Content Size:	0.0676"



TO OPEN: PUSH AND TURN CAP

NDC 05850-546-40

TYLENOL
with **NAPROXEN**

Acetaminophen 325mg,
Naproxen Sodium (NSAID) 180 mg
Tablets, Pain Reliever

LOT: XXXXXXX
EXP: YY/MM

40 Tablets

READ AND KEEP CAUTION FOR COMPLETE WARNINGS AND INFORMATION
(Do not use with other medications containing ASPIRIN; ANESTHESIA can cause heart danger)

Drug Facts

Active Ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever
Naproxen sodium 180 mg (NSAID)*	Pain reliever

Other ingredients: acetaminophen-inflammatory drug

Uses

- For the temporary relief of minor aches and pains due to:
 - headaches
 - arthritis
 - muscle aches
 - menstrual cramps
 - nerve pain of arthritis

Warnings

Aspirin-like heart disease warning:
The product contains aspirin/glycine. Severe side effects may occur if you take:
• more than three drugs containing aspirin or NSAIDs;
• more than 4,000 mg of acetaminophen in 24 hours;
• a low-calorie diet while using the product.

PEEL HERE →

Drug Facts (continued)	Drug Facts (continued)
Acetaminophen overdose may cause more serious reactions. Symptoms may include: nausea/vomiting, abdominal pain, loss of consciousness, and coma. If you suspect an overdose, call your doctor or poison control center immediately. NSAIDs (including aspirin) can increase your risk of a serious allergic reaction, which may include: hives, difficulty breathing, and swelling of the face, lips, tongue, and throat. Stop taking the medicine and call your doctor right away. If you already are allergic, you may not need to stop right away. NSAIDs can also bleeding warning. You may notice: bruising, bleeding, or blood in your stool or urine. If you have any of these symptoms, stop taking the medicine and call your doctor right away.	Heart attack and stroke warning. NSAIDs may cause heart attack, strokes, or death. These can occur before or after the first dose. The higher you take NSAIDs, the higher your risk. Stop taking NSAIDs if you have any of these symptoms: chest pain, shortness of breath, weakness, or dizziness. Stomach and intestinal bleeding warning. NSAIDs may cause stomach or intestinal bleeding, which may be fatal. The risk increases with longer use of NSAIDs. If you are taking aspirin, the risk is even higher. Stop taking NSAIDs if you have any of these symptoms: black or bloody stools, stomach pain, or vomiting blood. NSAIDs may cause kidney problems. NSAIDs may cause kidney problems, which may be fatal. Stop taking NSAIDs if you have any of these symptoms: change in the amount of urine you pass, swelling, or feeling tired. NSAIDs may cause liver problems. NSAIDs may cause liver problems, which may be fatal. Stop taking NSAIDs if you have any of these symptoms: yellowing of the skin or eyes, dark urine, or stomach pain.

Drug Facts (continued) When Using this product: Take with food or milk. Stir until just coats.	Drug Facts (continued) Directions: See and use with extra care. Read directions carefully.
Stop use and ask doctor if: you are pregnant or if the following signs occur:	Warnings: Do not use if you are allergic to any of the ingredients.
See your doctor if you have:	Do not use if you have:
Do not use if you have:	Do not use if you have:
Do not use if you have:	Do not use if you have:
Do not use if you have:	Do not use if you have:
Do not use if you have:	Do not use if you have:
Do not use if you have:	Do not use if you have:
Do not use if you have:	Do not use if you have:
Do not use if you have:	Do not use if you have:
Do not use if you have:	Do not use if you have:



Net Content Statement

PDP Area: (width x height) 3.09 in²

Net Content Size: 0.0671"

NET CONTENT

1/16 IN.

1/8 IN.

3/16 IN.

1/4 IN.

1/2 IN.

AREA OF PDP

5 SQ. IN. OR LESS

MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.

MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.

MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.

OVER 400 SQ. IN.

Drug Facts Format Information									
Drug Facts:	7.00 pt	Subheadings/Body Text:	5.00 pt	Barlines:	1.5 pt	Horizontal Scale:	85%		
Drug Facts (continued):	6.00 pt	Bullets:	4.00 pt	Hairlines:	0.5 pt	Tracking:	-12		
Headings:	6.00 pt	Leading:	5.50 pt						

LOT XXXXXX

EXP YYYMM

TO OPEN: PUSH AND TURN CAP

NDC 55540-546-80

TYLENOL[®]

WITH **NAPROXEN**

12hr

Acetaminophen 325 mg,
Naproxen Sodium (NSAID) 110 mg
Tablets, Pain Reliever

80 Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

Drug Facts

Active Ingredients (in each tablet)
Acetaminophen 325 mg
Naproxen sodium 110 mg (NSAID)
transdermal and inhaled drug

Pain Reliever
Pain Reliever

Drug Facts (continued)

Warnings
Acetaminophen liver damage warning:
This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ 3 or more alcoholic drinks every day while using this product
Acetaminophen allergy alert: may cause severe skin reactions.
Symptoms may include:
■ skin redness
■ hives
■ rash
If skin reaction occurs, stop use and seek medical help right away.

PEEL HERE

Drug Facts (continued)

NSAID allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives
■ facial swelling
■ asthma (wheezing)
■ shock
■ skin redness
■ rash
■ hives
If an allergic reaction occurs, stop use and seek medical help right away.
NSAID stomach bleeding warning:
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or blood drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. There can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
■ if you are allergic to acetaminophen, naproxen sodium, or any other pain reliever or any of its inactive ingredients in this product
■ right before or after heart surgery
Ask a doctor before use if
■ you have problems or serious side effects from taking pain relievers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because naproxen sodium may decrease the benefit of aspirin
■ taking the blood thinning drug warfarin
■ taking any other drug
When taking this product
■ take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint
■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke
■ chest pain
■ trouble breathing
■ weakness in one part or side of body
■ altered speech
■ leg swelling
■ pain gets worse or lasts more than 10 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ redness or swelling is present in the painful area
■ any new symptoms occur
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 30 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

DIRECTIONS
■ do not take more than directed (see overdose warnings)
■ do not use for more than 10 days unless directed by a doctor
■ drink a full glass of water with each dose
Adults and children 12 years and over
take 2 tablets every 12 hours while symptoms last
Children under 12 years
ask a doctor
■ do not take more than 7 tablets in 12 hours unless directed by a doctor
■ do not take more than 4 tablets in 24 hours unless directed by a doctor
Other information
■ each tablet contains sodium 10 mg
■ read full warnings and directions before use. Keep out of reach of children.
■ store between 20°-25°C (68°-77°F)
■ do not use if foil liner seal imprinted with "Sealed For Your Safety" is broken or missing.
Inactive ingredients: croscarmellose sodium, silicon dioxide, croscarmellose, hypromellose, lactose monohydrate, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polydextrose, croscarmellose, croscarmellose
Check for other? call 1-800-543-7865 (toll-free) or 215-273-4755 (in NJ)
Distributed by: Kaneva Brands LLC Summit, NJ 07901 © Kaneva Brands LLC 2026 Pat. www.kanevapats.com 3004438

Reference ID: 5840292

Net Content Statement

PDP Area: (width x height) 3.09 in²

Net Content Size: 0.0671"

NET CONTENT

1/16 IN.

1/8 IN.

3/16 IN.

1/4 IN.

1/2 IN.

AREA OF PDP

5 SQ. IN. OR LESS

MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.

MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.

MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.

OVER 400 SQ. IN.

Drug Facts Format Information									
Drug Facts:	7.00 pt	Subheadings/Body Text:	5.00 pt	Barlines:	1.5 pt	Horizontal Scale:	85%		
Drug Facts (continued):	6.00 pt	Bullets:	4.00 pt	Hairlines:	0.5 pt	Tracking:	-12		
Headings:	6.00 pt	Leading:	5.50 pt						

TO OPEN: PUSH AND TURN CAP

NO. XXXX-XXX-XX

TYLENOL

WITH NAPROXEN

12hr

Acetaminophen 325 mg, Naproxen Sodium (NSAID) 110 mg Tablets, Pain Reliever

90 Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

Drug Facts

Active ingredients (in each tablet)
Acetaminophen 325 mg
Naproxen sodium 110 mg (NSAID)
Nonsteroidal anti-inflammatory drug

Purpose
Pain reliever
Pain reliever

Elder
For the temporary relief of minor aches and pains due to:
backache
headache
muscle aches
menstrual cramps

Drug Facts (continued)

Warnings
Acetaminophen liver damage warning:
This product contains acetaminophen. Severe liver damage may occur if you take:
more than 4,000 mg of acetaminophen in 24 hours
5 or more alcoholic drinks every day while using this product
Acetaminophen allergy alert: may cause severe skin reactions.
Symptoms may include:
skin redness
hives
rash
if skin reaction occurs, stop use and seek medical help right away.

PEEL HERE

Drug Facts (continued)

NSAID allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives
facial swelling
asthma (wheezing)
shock
skin redness
itch
hives
If an allergic reaction occurs, stop use and seek medical help right away.
NSAID stomach bleeding warning:
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
if you are allergic to acetaminophen, naproxen sodium, or any other pain reliever or any of the inactive ingredients in this product
right before or after heart surgery
Ask a doctor before use if
you have liver disease
the stomach bleeding warning applies to you
you have problems or serious side effects from taking pain relievers
you have a history of stomach problems, such as heartburn
you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke
you are taking a diuretic
Ask a doctor or pharmacist before use if you are
under a doctor's care for any serious condition
taking aspirin for heart attack or stroke, because naproxen sodium may decrease the benefit of aspirin
taking the blood thinning drug warfarin
taking any other drug
When using this product
take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if
you experience any of the following signs of stomach bleeding:
fever
vomit blood
have bloody or black stools
have stomach pain that does not get better
you have symptoms of heart problems or stroke
chest pain
trouble breathing
weakness in one part or side of body
stomach upset
leg swelling
pain gets worse or lasts more than 10 days
you have difficulty swallowing
if it feels like the pill is stuck in your throat
redness or swelling is present in the painful area
any new symptoms occur
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 30 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Dosage form
do not take more than 10 days unless directed by a doctor
drink 8 full glasses of water with each dose
adults and children 12 years and over
take 2 tablets every 12 hours while symptoms last
children under 12 years
ask a doctor
do not take more than 7 tablets in 12 hours unless directed by a doctor
do not take more than 4 tablets in 24 hours unless directed by a doctor
Other information
each tablet contains sodium 10 mg
read all warnings and directions before use. Keep out of reach of children.
do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.
Inactive ingredients include silicon dioxide, croscarmellose, hydroxypropyl methylcellulose, magnesium stearate, medium chain triglycerides, polyvinylpyrrolidone, polyethylene glycol, polyethylene glycol, titanium dioxide
Check for use
call 1-800-343-7865 (toll-free) or 215-273-8765 (collect)

Reference ID: 5840292

Net Content Statement

PDP Area: (width x height) 6.89 in²
Net Content Size: 0.107"

NET CONTENT

1/8 IN.
1/8 IN.
3/16 IN.
1/4 IN.
1/2 IN.

AREA OF PDP
5 SQ. IN. OR LESS
MORE THAN 5 SQ. IN. BUT
NOT MORE THAN 25 SQ. IN.
MORE THAN 25 SQ. IN. BUT
NOT MORE THAN 100 SQ. IN.
MORE THAN 100 SQ. IN. BUT
NOT MORE THAN 400 SQ. IN.
OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	10.00 pt	Subheadings/Body Text:	6.00 pt	Badlines:	2.0 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Headlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.30 pt	Tracking:	0

EXP: YYYMM

LOT: XXXXXX

TO OPEN: PUSH AND TURN CAP

NO: 50590-346-14

TYLENOL[®]

WITH NAPROXEN

12

Acetaminophen 325 mg,
Naproxen Sodium (NSAID) 110 mg

Tablets, Pain Reliever

140 Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medications containing ACETAMINOPHEN; can cause liver damage.

Drug Facts

Active ingredients (in each tablet)

Acetaminophen 325 mg

Naproxen sodium 110 mg (NSAID)

nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

For the temporary relief of minor aches and pains due to:
■ headache ■ backache
■ muscular aches ■ toothache
■ menstrual cramps ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning
This product contains acetaminophen. Severe liver damage may occur if you take:
■ with other drugs containing acetaminophen
■ more than 4,000mg of acetaminophen in 24 hours
■ 3 or more alcoholic drinks every day while using this product

PEEL HERE

Drug Facts (continued)

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:
■ skin redness ■ hives ■ rash
If skin reaction occurs, stop use and seek medical help right away.
NSAID allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin redness ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
NSAID stomach bleeding warning:
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use:
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
■ if you are allergic to acetaminophen, naproxen sodium, or any other pain reliever or any of the inactive ingredients in this product
■ right before or after heart surgery
Ask a doctor before use if:
■ you have liver disease
■ the stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are:
■ under a care for any serious condition
■ taking aspirin for heart attack or stroke, because naproxen sodium may decrease the benefit of aspirin
■ taking the blood thinning drug warfarin
■ taking any other drug.

Drug Facts (continued)

When using this product ■ Take with food or milk if stomach upset occurs
Stop use and ask a doctor if:
■ you experience any of the following signs of stomach bleeding:
■ heart burn ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing ■ weakness in one part or side of body
■ slurred speech ■ big swelling
■ pain gets worse or lasts more than 10 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ redness or swelling is present in the painful area
■ any new symptoms occur
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
Overdose warnings: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions ■ do not take more than directed (see overdose warning)
■ do not use for more than 10 days unless directed by a doctor
■ drink a full glass of water with each dose
Take 2 tablets every 12 hours while symptoms last.
adults and children 12 years and over
children under 12 years ■ ask a doctor
■ do not take more than 7 tablets in 12 hours unless directed by a doctor
■ do not take more than 4 tablets in 24 hours unless directed by a doctor
Other information
■ each tablet contains: sodium 90 mg
■ store between 20°-25° (68°-77°)
■ read all warnings and directions before use. Keep carton.
Do not use if full inner seal imprinted with "Sealed For Your Safety" is broken or missing.
Inactive ingredients: cellular silicon dioxide, croscarmellose, hypromellose, lactose monohydrate, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polyvinyl alcohol, titanium dioxide
Questions? call 1-800-342-7668 (ask-free) or 216-273-8756 (collect).

Manufactured by: Kenvue Brands LLC, Summit, NJ 07901
© Kenvue Brands LLC 2018 P/N: www.tylenopain.com 30084230

Reference ID: 5840292

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	12.3 In ²	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.129"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	12.00 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.50 pt	Horizontal Scale:	90%
				Tracking:	-6



Net Content Statement

PDP Area: (width x height) 6.89 in²
Net Content Size: 0.107"

NET CONTENT

1/8 IN.
1/8 IN.
3/16 IN.
1/4 IN.
1/2 IN.

AREA OF PDP
5 SQ. IN. OR LESS
MORE THAN 5 SQ. IN. BUT
NOT MORE THAN 25 SQ. IN.
MORE THAN 25 SQ. IN. BUT
NOT MORE THAN 100 SQ. IN.
MORE THAN 100 SQ. IN. BUT
NOT MORE THAN 400 SQ. IN.
OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	10.00 pt	Subheadings/Body Text:	6.00 pt	Badlines:	2.0 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Headlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.30 pt	Tracking:	0

XXXXXX
LOT XXXXXX
EXP: YYYMM

TO OPEN: PUSH AND TURN CAP

NDC 50580-346-02

TYLENOL®

WITH NAPROXEN

12hr

Acetaminophen 325 mg,
Naproxen Sodium (NSAID) 110 mg
Tablets, Pain Reliever

200 Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medicines containing ACETAMINOPHEN, can cause liver damage.

Drug Facts

Active Ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain Reliever
Naproxen sodium 110 mg (NSAID) "nonsteroidal anti-inflammatory drug"	Pain Reliever

Uses

For the temporary relief of minor aches and pains due to:
■ headache
■ muscular aches
■ menstrual cramps
■ backache
■ toothache
■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning:
This product contains acetaminophen. Severe liver damage may occur if you take:
■ with other drugs containing acetaminophen
■ more than 4,000 mg of acetaminophen in 24 hours
■ 3 or more alcoholic drinks every day while using this product

PEEL HERE

Drug Facts (continued)

Acetaminophen allergy alert: may cause severe skin reactions:
Symptoms may include: ■ skin redness ■ blisters ■ rash
If skin reaction occurs, stop use and seek medical help right away.
NSAID allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin redness ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
NSAID stomach bleeding warning:
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use:
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen, naproxen sodium, or any other pain reliever or any of the inactive ingredients in this product.
■ right before or after heart surgery
Ask a doctor before use if:
■ you have liver disease
■ the stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver, kidney, kidney disease, asthma, or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are:
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because naproxen sodium may decrease this benefit of aspirin
■ taking the blood thinning drug warfarin
■ taking any other drug

Drug Facts (continued)

When using this product ■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if:
■ you experience any of the following signs of stomach bleeding:
■ hot heart ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing ■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts more than 10 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ redness or swelling is present in the painful area
■ any new symptoms occur
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 30 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions ■ do not take more than directed (see overdose warning)
■ do not use for more than 10 days unless directed by a doctor
■ Drink a full glass of water with each dose
adults and children 12 years and over take 2 tablets every 12 hours while symptoms last
Children under 12 years ask a doctor
■ do not take more than 2 tablets in 12 hours unless directed by a doctor
■ do not take more than 4 tablets in 24 hours unless directed by a doctor
Other information
■ each tablet contains sodium 10 mg
■ Read all warnings and directions before use. Keep carton.
■ Store between 20°-25°C (68°-77°F).
■ Do not use if full lower seal separated with "Sealed For Your Safety" is broken or missing.
Inactive ingredients: cellulose, silicon dioxide, croscarmellose, hypromellose, lactose monohydrate, magnesium stearate, sodium cholethrate, microcrystalline cellulose, polydextrated starch, titanium dioxide
Questions? call 1-800-343-7895 (toll free) or 215-273-8750 (toll-free)
© 2006 Beechler, a Sanofi-Schering-Plough company, KJ 07060
© 2006 Beechler, a Sanofi-Schering-Plough company, KJ 07060
© 2006 Beechler, a Sanofi-Schering-Plough company, KJ 07060

Reference ID: 5840292

Net Content Statement

PDP Area: (width x height) 6.89 in²
Net Content Size: 0.107"

NET CONTENT

1/8 IN.
3/16 IN.
1/4 IN.
1/2 IN.

AREA OF PDP

5 SQ. IN. OR LESS
MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	10.00 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.0 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Headlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.30 pt	Tracking:	0

EPD

LOT: XXXXXX
EXP: YYYMM

TO OPEN: PUSH AND TURN CAP

NDC XXXX-XXX-XX

TYLENOL®

WITH NAPROXEN

12hr

Acetaminophen 325 mg,
Naproxen Sodium (NSAID) 110 mg
Tablets, Pain Reliever

220 Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medicines containing ACETAMINOPHEN, can cause liver damage.

Drug Facts

Active Ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever
Naproxen sodium 110 mg (NSAID)	Pain reliever
*nonsteroidal anti-inflammatory drug	

Uses

■ For the temporary relief of minor aches and pains due to:
■ headache
■ muscular aches
■ menstrual cramps
■ backache
■ toothache
■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning:
This product contains acetaminophen. Severe liver damage may occur if you take:
■ with other drugs containing acetaminophen
■ more than 4,000 mg of acetaminophen in 24 hours
■ 3 or more alcoholic drinks every day while using this product

PEEL HERE

Drug Facts (continued)

Acetaminophen allergy alert: may cause severe skin reactions:
Symptoms may include: ■ skin redness ■ blisters ■ rash
If skin reaction occurs, stop use and seek medical help right away.
NSAID allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin redness ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
NSAID stomach bleeding warning:
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use:
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
■ if you are allergic to acetaminophen, naproxen sodium, or any other pain reliever or any of the inactive ingredients in this product
■ right before or after heart surgery
Ask a doctor before use if:
■ you have liver disease
■ the stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver, kidney, kidney disease, asthma, or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are:
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because naproxen sodium may decrease this benefit of aspirin
■ taking the blood thinning drug warfarin
■ taking any other drug

Drug Facts (continued)

When using this product ■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if:
■ you experience any of the following signs of stomach bleeding:
■ hot flat ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing ■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts more than 10 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ redness or swelling is present in the painful area
■ any new symptoms occur
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions ■ do not take more than directed (see overdose warning)
■ do not use for more than 10 days unless directed by a doctor
■ drink a full glass of water with each dose
adults and children 12 years and over take 2 tablets every 12 hours while symptoms last
Children under 12 years ask a doctor
■ do not take more than 2 tablets in 12 hours unless directed by a doctor
■ do not take more than 4 tablets in 24 hours unless directed by a doctor
Other information
■ each tablet contains sodium 10 mg
■ read all warnings and directions before use. Keep carton.
■ store between 50°-25° (58°-77°F)
■ do not use if full meter seal labeled with "Stand for Your Safety" is broken or missing.
Inactive ingredients: colloidal silicon dioxide, croscarmellose, hypromellose, lactose monohydrate, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polydextran starch, titanium dioxide
Questions? call 1-800-343-7895 (toll free) or 215-273-8758 (toll-free)
© 2009 Sunovion, Inc. Sunovion, Inc. is a trademark of Sunovion, Inc. All other trademarks are the property of their respective owners.

Reference ID: 5840292

