

Drug Facts Format Information				
Drug Facts:	5.50 pt	Subheadings/Body Text:	4.50 pt	Barlines: 1.0 pt
Drug Facts (continued):	4.50 pt	Bullets:	3.50 pt	Horizontal Scale: 76%
Headings:	4.50 pt	Leading:	4.50 pt	Tracking: -20

[illegible]

Reference ID: 5840292

Drug Facts Format Information				
Drug Facts:	5.50 pt	Subheadings/Body Text:	4.50 pt	Barlines: 1.0 pt
Drug Facts (continued):	4.50 pt	Bullets:	3.50 pt	Horizontal Scale: 76%
Headings:	4.50 pt	Leading:	4.20/4.50 pt	Hairlines: 0.5 pt
				Tracking: -20

[illegible]

Net Content Statement		NET CONTENT	AREA OF POP
PDP Area: (width x height)	6.84 in²	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.125"	1/8 IN.	MORE THAN 5 SQ. IN., BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN., BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN., BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information				
Drug Facts:	9.00 pt	Subheadings/Body Text:	6.00 pt	Barlines: 1.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Horizontal Scale: 75%
Headings:	8.00 pt	Leading:	5.9/6.0 pt	Tracking: -25



Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	6.84 in ²	1/8 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.125"	3/16 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		1/4 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/2 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	9.00 pt	Subheadings/Body Text:	6.00 pt	Badlines:	1.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Headlines:	0.5 pt
Headings:	8.00 pt	Leading:	5.8/5.9 pt	Tracking:	-15



Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (Width x height)	3.08 in ²	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.0676"	1/8 IN.	MORE THAN 5 SQ. IN., BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN., BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN., BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information				
Drug Facts:	7.00 pt	Subheadings/Body Text:	5.00 pt	Barlines: 1.5 pt
Drug Facts (continued):	6.00 pt	Bullets:	4.00 pt	Hairlines: 0.5 pt
Headings:	6.00 pt	Leading:	4.9/5.0 pt	Horizontal Scale: 77% Tracking: -20



LOT: XXXXXXXX
EXP: YYYYMM

TO OPEN, PUSH AND TURN CAP

NDC 50580-346-40




Acetaminophen 325 mg,
Naprofen Sodium (NSAID) 110 mg
Tablets, Pain Reliever

40 Tablets

READ AND KEEP CAUTION FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medicines containing **ACETAMINOPHEN**, as taking more **dangerous**

Drug Facts

Active Ingredients (in each tablet)	Purpose
Naprofen 110 mg (NSAID)	Pain reliever
Naprofen sodium 110 mg (NSAID)	Pain reliever
Acetaminophen 325 mg	Pain reliever

Uses

For the temporary relief of minor aches and pains due to:

- headaches • backaches • menstrual aches
- muscular cramps • minor pain of arthritis

Warnings

Naprofen has some drug warnings

This product contains acetaminophen. Serious liver damage may occur if you take:
 • more than 4 other drugs containing acetaminophen
 • more than 4,000 mg of acetaminophen in 24 hours
 • 3 or more alcoholic drinks every day while using this product

➔

PPE HERE

[illegible][illegible]

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height) Net Content Size:	7.44 in ² 0.1254"	1/16 IN.	5 SQ. IN. OR LESS
		1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information				
Drug Facts:	9.00 pt	Subheadings/Body Text:	6.00 pt	Barlines: 2.0 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Horizontal Scale: 78%
Headings:	8.00 pt	Leading:	5.9/6.0 pt	Tracking: -25



Net Content Statement

PDP Area: (width x height) 3.09 in²
Net Content Size: 0.0671"

NET CONTENT

1/16 IN.
1/8 IN.
3/16 IN.
1/4 IN.
1/2 IN.

AREA OF PDP

5 SQ. IN. OR LESS
MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
OVER 400 SQ. IN.

Drug Facts Format Information							
Drug Facts:	7.00 pt	Subheadings/Body Text:	5.00 pt	Barlines:	1.5 pt	Horizontal Scale:	85%
Drug Facts (continued):	6.00 pt	Bullets:	4.00 pt	Hairlines:	0.5 pt	Tracking:	-12
Headings:	6.00 pt	Leading:	5.50 pt				

TO OPEN: PUSH AND TURN CAP

NDK 59360-346-90

TYLENOL[®]

WITH NAPROXEN

12hr

Acetaminophen 325 mg,
Naproxen Sodium (NSAID) 110 mg

Tablets, Pain Reliever

80 Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

PEEL HERE

Drug Facts

Active ingredients (in each tablet)

Acetaminophen 325 mg
Naproxen sodium 110 mg (NSAID)
Nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

For the temporary relief of minor aches and pains due to:
Headache
Muscular aches
Menstrual cramps

For the temporary relief of minor aches and pains due to:
Backache
Nausea
Minor pain of arthritis

Drug Facts (continued)

Warnings

Acetaminophen liver damage warning
This product contains acetaminophen. Severe liver damage may occur if you take:
more than 4,000 mg of acetaminophen in 24 hours
or more alcohol
drinks every day while using this product.
Acetaminophen allergy alert
Acetaminophen allergy alert may cause severe skin reactions. Symptoms may include:
skin redness
hives
rash
If skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

NSAID allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives
facial swelling
asthma (wheezing)
shock
If an allergic reaction occurs, stop use and seek medical help right away.
NSAID stomach bleeding warning
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 65 or older
have had stomach ulcers or bleeding problems
take blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
If you are allergic to acetaminophen, naproxen sodium, or any other pain reliever or any of the inactive ingredients in this product
right before or after heart surgery
Ask a doctor before use if:
you have liver disease
you have problems or serious side effects from taking pain relievers
you have a history of stomach problems, such as heartburn
you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
you are taking a diuretic
Ask a doctor or pharmacist before use if you are:
under a doctor's care for any serious condition
taking aspirin for heart attack or stroke, because naproxen sodium may decrease the benefit of aspirin
taking the blood thinning drug warfarin
taking any other drug
When using this product
take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if:
you experience any of the following signs of stomach bleeding:
feet faint
vomit blood
have bloody or black stools
have stomach pain that does not get better
you have symptoms of heart problems or stroke
chest pain
trouble breathing
weakness in one part or side of body
slurred speech
leg swelling
pain gets worse or lasts more than 10 days
you have difficulty swallowing
it feels like the pill is stuck in your throat
redness or swelling is present in the painful area
any new symptoms occur
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 32 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions
Do not take more than directed (see overdose warning).
Do not use for more than 10 days unless directed by a doctor.
Adults and children 12 years and over
Take 2 tablets every 12 hours while symptoms last.
Children under 12 years
Ask a doctor.
Do not take more than 2 tablets in 12 hours unless directed by a doctor.
Do not take more than 4 tablets in 24 hours unless directed by a doctor.
Other information
each tablet contains sodium 10 mg
read all warnings and directions before use. Keep carton.
Store between 20-25°C (68-77°F).
Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.
Inactive ingredients
cellulose, silicon dioxide, croscarmellose, hypromellose, lactose monohydrate, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, titanium dioxide.
Overdose
Call 1-800-343-7005, ask your doctor, or call 1-800-477-4775.
Distributed by: Kennewick Brands LLC, Summit, NJ 07901 © Kennewick Brands LLC 2025 Pdf: www.kennewickbrands.com 3006459

Reference ID: 5840292

Drug Facts Format Information					
Drug Facts:	10.00 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.50 pt	Horizontal Scale:	80%
				Tracking:	-5




Drug Facts Format Information							
Drug Facts:	7.00 pt	Subheadings/Body Text:	5.00 pt	Barlines:	1.5 pt	Horizontal Scale:	85%
Drug Facts (continued):	6.00 pt	Bullets:	4.00 pt	Hairlines:	0.5 pt	Tracking:	-12
Headings:	6.00 pt	Leading:	5.50 pt				

Drug Facts (continued)		Drug Facts (continued)	
allergic allergy alert: naproxen might cause a worse allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • facial swelling • asthma (wheezing) • shock allergy alert: if you have had an allergic reaction to aspirin, you may have a more severe reaction to naproxen. if you are allergic to naproxen, stop use and seek medical help right away.		Do not use • with any other drug containing acetaminophen (paracetamol) or nonsteroidal anti-inflammatory drugs (NSAIDs). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are allergic to acetaminophen or NSAIDs, ask your doctor or pharmacist for more information. • with any other pain reliever or any of the inactive ingredients in this product right before or after heart surgery.	
Heartburn, stomach bleeding warning: Heartburn: heartburn or acid reflux may cause worse stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems in the past • take blood thinning anticoagulants or steroid drugs • take aspirin, aspirin-like drugs, prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take naproxen for a longer time (see directions) Heart attack and stroke warning: NSAIDs, aspirin, and aspirin-like drugs may increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed for aspirin, heart failure, and stroke.		Ask a doctor before use if: • you have liver problems or alcoholism • you have kidney problems or are on dialysis • you have stomach or heartburn before you use this product • you have had blood pressure, heart failure, low blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke • you are taking diuretics Ask a doctor or pharmacist before use if you are: • under a doctor's care for any serious condition • taking aspirin, heart attack or stroke, blood thinning aspirin may decrease this benefit • taking the blood-thinning drug warfarin • taking any other drugs	
When using this product • do not take more than directed for aspirin, heart attack, heart failure, and stroke		When using this product • do not have more than 1 stomach upset benefit	

Reference ID: 5840292

Drug Facts Format Information				
Drug Facts:	10.00 pt	Subheadings/Body Text:	6.00 pt	Barlines: 2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines: 0.5 pt
Headings:	8.00 pt	Leading:	5.9-6.0 pt	Horizontal Scale: 80% Tracking: -25



Drug Facts Format Information					
Drug Facts:	12.00 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.50 pt	Horizontal Scale:	90%
				Tracking:	-6

Reference ID: 5840292

Net Content Statement

PDP Area: (width) x (height)
Net Content Size: 6.89 in²
0.107"

NET CONTENT

1/16 IN.
1/8 IN.
3/16 IN.
1/4 IN.
1/2 IN.

AREA OF PDP
5 SQ. IN. OR LESS
MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	10.00 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.0 pt
Drug Facts (continued):	8.00 pt	Tablets:	5.00 pt	Barlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.30 pt	Tracking:	0
Horizontal Scale: 95%					

TO OPEN: PUSH AND TURN CAP

NDC 50580-346-02

LOT: XXXXXX
EXP: YYYYMM

TYLENOL[®]

WITH NAPROXEN[®]

12hr

Acetaminophen 325 mg,
Naproxen Sodium (NSAID) 110 mg
Tablets, Pain Reliever

200 Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medication containing ACETAMINOPHEN, can cause liver damage.

Drug Facts

Active ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever
Naproxen sodium 110 mg (NSAID)*	Pain reliever
*nonsteroidal anti-inflammatory drug	

Uses

For the temporary relief of minor aches and pains due to:
■ headache ■ backache
■ muscle aches ■ toothache
■ menstrual cramps ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning:
This product contains acetaminophen. Severe liver damage may occur if you take:
■ with other drugs containing acetaminophen
■ more than 4,000 mg of acetaminophen in 24 hours
■ 3 or more alcoholic drinks every day while using this product

PEEL HERE

Drug Facts (continued)

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash
If skin reaction occurs, stop use and seek medical help right away.
NSAID allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin redness ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
NSAID stomach bleeding warning:
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use:
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen, naproxen sodium, or any other pain reliever or any of the inactive ingredients in this product.
■ right before or after heart surgery
Ask a doctor before use if:
■ you have liver disease.
■ the stomach bleeding warning applies to you.
■ you have problems or serious side effects from taking pain relievers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are:
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because naproxen sodium may decrease this benefit of aspirin
■ taking the blood thinning drug warfarin
■ taking any other drug.

Drug Facts (continued)

When using this product:
■ Take with food or milk if stomach upset occurs.
Stop use and ask a doctor if:
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing
■ slurred speech ■ leg swelling
■ pain gets worse or lasts more than 10 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ redness or swelling is present in the painful area
■ any new symptoms occur.
These could be signs of a serious condition.
If pregnant or breast-feeding, stop use and ask your health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-522-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)


Directions ■ Do not take more than directed (see overdose warning)
■ do not use for more than 10 days unless directed by a doctor
■ drink a full glass of water with each dose.

Adults and children 12 years and over	Take 2 tablets every 12 hours while symptoms last.
Children under 12 years	Ask a doctor

■ do not take more than 2 tablets in 12 hours unless directed by a doctor
■ do not take more than 4 tablets in 24 hours unless directed by a doctor
Other information
■ each tablet contains: sodium 10 mg
■ read all warnings and directions before use. Keep carton.
■ store between 59°-77° F (15°-25° C).
■ do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.
Inactive ingredients: cellulose, dibasic sodium phosphate, croscarmellose, hypromellose, lactose monohydrate, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polyethylene glycol, polyethylene glycol, titanium dioxide
Questions? call 1-800-345-7989 (ask, free) or 219-273-8756 (collect)
© 2006 Novartis Pharmaceuticals Corporation, East Hanover, NJ 07923
® Novartis Pharmaceuticals Corporation 30064216

Reference ID: 5840292

Drug Facts Format Information				
Drug Facts:	12.00 pt	Subheadings/Body Text:	6.00 pt	Barlines: 2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Horizontal Scale: 90%
Headings:	8.00 pt	Leading:	6.50 pt	Tracking: -6



Net Content Statement

PDP Area: (width x height) 6.89 in²

Net Content Size: 0.107"

NET CONTENT

1 1/16 IN.
1 1/8 IN.
3 1/16 IN.
1 1/4 IN.
1 1/2 IN.

AREA OF PDP
3 SQ. IN. OR LESS
MORE THAN 3 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	10.00 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.0 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Barlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.30 pt	Tracking:	0

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

TYLENOL[®] WITH NAPROXEN[®] 12hr

Acetaminophen 325 mg, Naproxen Sodium (NSAID) 110 mg Tablets, Pain Reliever

220 Tablets

Drug Facts

Active ingredients (in each tablet)
Acetaminophen 325 mg
Naproxen sodium 110 mg (NSAID)¹
Nondrolool anti-inflammatory drug

Purpose
Pain reliever

Uses
For the temporary relief of minor aches and pains due to:
Headache
Muscular aches
Toothache
Menstrual cramps
Minor pain of arthritis

Warnings
Acetaminophen liver damage warning:
This product contains acetaminophen. Severe liver damage may occur if you take:
With other drugs containing acetaminophen.
More than 4,000 mg of acetaminophen in 24 hours.
3 or more alcoholic drinks every day while using this product.

PEEL HERE

Drug Facts (continued)

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash
If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin redness ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ take 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use:
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen, naproxen sodium, or any other pain reliever or any of the inactive ingredients in this product.
■ right before or after heart surgery

Ask a doctor before use if:
■ you have liver disease
■ the stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are:
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because naproxen sodium may decrease this benefit of aspirin
■ taking the blood thinning drug warfarin
■ taking any other drug

Drug Facts (continued)

When using this product: ■ Take with food or milk if stomach upset occurs

Stop use and ask a doctor if:
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing ■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ you get worse or last more than 10 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ nausea or swelling is present in the painful area
■ any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding: use a health professional before use. It is especially important not to use naproxen sodium at 30 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-322-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions: ■ Do not take more than directed (see overdose warning)
■ do not use for more than 10 days unless directed by a doctor
■ drink a full glass of water with each dose

Adults and children 12 years and over: take 2 tablets every 12 hours while symptoms last

Children under 12 years: ask a doctor

■ do not take more than 2 tablets in 12 hours unless directed by a doctor
■ do not take more than 4 tablets in 24 hours unless directed by a doctor

Other information
■ each tablet contains: sodium 10 mg
■ read all warnings and directions before use. Keep carton.
■ store between 59°-77° F (15°-25° C) (59°-77° F).

Do not use if full inner seal imprinted with "Sealed For Your Safety" is broken or missing.

Inactive ingredients: croscarmellose sodium, croscarmellose, hypromellose, lactose monohydrate, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions? call 1-800-343-7895 (toll free) or 312-873-8755 (collect)

© 2009 Bayer HealthCare LLC, Summit, NJ 07961
© Kenvue Brands LLC 2026 Pat. www.kenvuebrands.com 00000000

Reference ID: 5840292

Drug Facts Format Information					
Drug Facts:	12.00 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.50 pt	Horizontal Scale:	90%
				Tracking:	-6